

Clinical trials of omega-3 fatty acids for cardiovascular prevention in patients at high risk

TrialResults-center www.trialresultscenter.org

1 omega-3 Fatty acids

Trial	Treatments	Patients	Trials design and methods
diet vs control			
Burr (DART 2) , 2003 n=1571/1543 follow-up: 36-108 months	dietary advice (to eat more oily fish) versus No such dietary advice or capsules	men being treated for angina	Parallel groups open UK
Burr (DART) , 1989 n=1015/1018 follow-up: 24 months	dietary advice (to eat more oily fish) versus No such dietary advice or capsulesish)ag	post-MI	Parallel groups open UK
fish oil vs control			
Bemelmans , 2002 n=51/52 follow-up: 24 months	a-lin rich margarine (80% fat of which 15% was a-lin) versus linoleic rich margarine (80% fat of which 0.3% was a-lin), identical in taste and packaging	patients with multiple cardiovascular risk factors (10 yr IHD risk 20%)	Parallel groups double-blind the Netherlands
Brox , 2001 n=40/40 follow-up:	seal oil - 15 ml/d (2.6g EPA + DHA) versus no supplement	dyslipidaemia	open with blind assessment
Franzen , 1993 n=15/15 follow-up: 12 months	fish oil capsules, 9g/d (1.8g EPA + 1.4g DHA daily) versus olive oil capsules	people with angiographically determined CHDg	Parallel groups double-blind
Shimizu , 1995 n=29/16 follow-up: 12 months	EPA-ethyl capsules 3/d (0.9g/d EPA) versus no treatment	people with non-insulin dependant diabetes	Parallel groups open Japan
MaxEPA vs control			
Bellamy , 1992 n=60/60 follow-up: 7 months	MaxEPA capsules (3g/d EPA + DHA) versus no treatment	people referred for coronary angioplasty	Parallel groups NA UK
Dehmer , 1998 n=46/44 follow-up: 6 months	MaxEPA capsules, 18/d (5.4g EPA + DHA daily) versus no treatment	men undergoing coronary angioplasty imag	open US

continued...

Trial	Treatments	Patients	Trials design and methods
Kaul , 1992 n=58/49 follow-up: 6 months	MaxEPA capsules, 10/d (3g/d EPA + DHA) versus no treatment	people undergoing angioplasty	Parallel groups open India
Omacor vs control			
Eritslund , 1996 n=317/293 follow-up: 12 months	Omacor capsules, 4/d (3.3g EPA + DHA daily) versus no treatment	people admitted for coronary bypass grafting	Parallel groups open Norway
GISSI-P , 1999 n=5665/5668 follow-up: median 40 months	Omacor gelatine capsules, 1/d (0.9g/d EPA + DHA daily) versus no treatment	people with recent myocardial infarction	Parallel groups open Italy
omega-3 Fatty acids vs control			
OMEGA , 2009 [NCT00251134] n=1940/1911 follow-up: 1 year	omega-3 fatty acids 1g daily (and standard medical therapy) versus standard medical therapy alone	Patients within 3-14 days after a non-ST-elevation myocardial infarction (NSTEMI) or ST-elevation myocardial infarction (STEMI)	Parallel groups open Germany
Promega vs control			
Milner , 1989 n=100/100 follow-up: 6 months	Promega 9 capsules/d (4.5g EPA + DHA) versus no treatment	people about to undergo angioplasty	Parallel groups open with blind assessment US
alpha-linolenic acid vs placebo			
ALPHA OMEGA (ALA) , 2010 [NCT00127452] n=2409/2428 follow-up: 40 months	margarine supplemented with plant-derived alpha-linolenic acid (with a targeted additional daily intake of 2 g of ALA) versus placebo	men and women with a history of myocardial infarction	Factorial plan double-blind the Netherlands
Natvig , 1968 n=6716/6690 follow-up: 12 months	linseed oil, 10 ml /d (55% a-linolenic acid) versus placebo (sunflower oil, 10 ml/d (1.4% a-linolenic acid))	working men, though a few had had a previous MI or angina7ieq	Parallel groups double-blind Norway
Esapent vs placebo			
Maresta , 2002 n=169/170 follow-up: 7 months	Esapent capsules, 6/d for 2 mo, then 3/d (5.1g/d EPA + DHA initially, later 2.6g/d) versus placebo (identical olive oil capsules, 6/d for 2 mo, then 3/d)	undergoing planned PTCAB	Parallel groups double-blind Italy
Sirtori , 1998 n=470/465 follow-up: 6 months	Esapent fish oil capsules 3/d for first 2 mo, 2/d after that (2.6g/dEPA + DHA initially, then 1.8g/d) versus placebo (olive oil capsules 3/d for first 2 mo, 2/d after that)	people with raised triglycerides plus glucose intolerance, non-insulindependent diabetes or hypertension	Parallel groups double blind Italy

continued...

Trial	Treatments	Patients	Trials design and methods
Eskisol vs placebo			
Rossing , 1996 n=18/18 follow-up: 12 months	Eskisol fish oil emulsion 21 ml/d (4.6g EPA +DHA) versus placebo (olive oil emulsion 21 ml/d)	people with insulin dependant diabetes, diabetic nephropathy and normalBP	Parallel groups double blind Denmark
fish oil vs placebo			
Borchgrevink , 1966 n=100/100 follow-up: mean 10 months (range 3-16 mo)	linseed oil 10 ml/d initially, later raised to 20 or 30 ml/d (4.5g/d a-lin, later 9 or 13.5 g/d) versus placebo (corn oil, 10 ml/d initially, later raised to 20 or 30 ml/d)	men with impending or recent myocardial infarctionage/p	Parallel groups double-blind Norway
Leaf , 1994 n=275/276 follow-up: 6 months	fish oil concentrate capsules 10x1 g/d (6.9g/d EPA + DHA) versus placebo (corn oil capsules 10x1 g/d with 0.4% fish oil to maintain blinding (0.003g/d EPA + DHA))	people undergoing angioplasty	Parallel groups double blind US
Sacks (TOHP 1) , 1994 [NCT00000528] n=NA follow-up:	fish oil versus placebo	double blind	double-blind
von Schacky , 1999 n=112/111 follow-up: 24 months	concentrated fish oil capsules, 6/d for first 3 mo, 3/d for rest of study (4g/d EPA +DHA + DPA+ a-lin for first 3 mo, then 2g/d) versus placebo (capsules containing fat which replicated the fat composition of the average European diet, 6/d forfirst 3 mo, 3/d for rest of study, opaque soft gelatine capsules identical to fish capsules)	people with angiographically proven coronary artery disease	Parallel groups double blind Germany
HiEPA vs placebo			
Hawthorne , 1992 n=49/47 follow-up: 12 months	HiEPA oil, 10 ml x 2/d (5.6g/d EPA + DHA) versus placebo (olive oil, 10 ml x 2/d (0g/d EPA + DHA))	people with ulcerative colitis	Parallel groups double blind UK
MaxEPA vs placebo			
Bairati , 1992 n=107/98 follow-up: 7 months	MaxEPA, 15 capsules/d (4.5g EPA + DHA) versus placebo (olive oil, 15 capsules/d)	patients undergoing planned angioplasty	Parallel groups double blind Canada
Lau , 1993 n=32/32 follow-up: 12 months	MaxEPA 10x 1g capsules daily (2.8g/d EPA + DHA) versus placebo (air-filled capsules, 10/d)	people with rheumatoid arthritisp	Parallel groups double blind UK

continued...

Trial	Treatments	Patients	Trials design and methods
Lau , 1995 n=25/20 follow-up: 6 months	MaxEPA 10x 1g capsules daily (2.8g/d EPA + DHA) versus placebo (air-filled capsules, 10/d)	people with rheumatoid arthritis	Parallel groups double blind Hong Kong
Nye , 1990 n=36/37 follow-up: 12 months	MaxEPA capsules 12/d (2.2g EPA) versus placebo (olive oil capsules, 12/d, identical to MaxEPA)	people undergoing angioplasty	Parallel groups double blind New Zealand
Singh , 1997 n=122/118 follow-up: 12 months	MaxEPA fish oil capsules 6/d (1.8g EPA + DHA) versus placebo (aluminium hydroxide 100 mg/d)	people with suspected acute MI	Parallel groups double blind India
Omacor vs placebo			
Johansen , 1999 n=250/250 follow-up: 6.5 months	Omacor capsules, 6/d (5g EPA + DHA daily) versus placebo (corn oil capsules, 6/d)	people about to undergo elective coronary angioplasty	Parallel groups double blind Norway
Nilsen , 2001 n=150/150 follow-up: 24 months	Omacor capsules 4/d (3.5g EPA + DHA) versus placebo (corn oil capsules, 4/d)	people with acute myocardial infarction 4-8 days agoe/pj	Parallel groups double-blind Norway
omega-3 fatty acids vs placebo			
ALPHA OMEGA (EPA DHA) , 2010 [NCT00127452] n=2404/2433 follow-up: 40 months	400 mg per day supplement of the fish oil fatty acids EPA (eicosapentaenoic acid) and DHA (docosahexaenoic acid) via enriched margarines versus placebo	men and women with a history of myocardial infarction	Factorial plan double-blind the Netherlands
Risk and Prevention Study , 2013 [NCT00317707.] n=6244/6269 follow-up: 5 year (median)	n-3 fatty acids (1 g daily) versus placebo (olive oil)	men and women with multiple cardiovascular risk factors or atherosclerotic vascular disease but not myocardial infarction	double-blind
GISSI HF fatty acid , 2008 [NCT00336336.] n=3494/3481 follow-up: 3.9y median (IQR 3-4.4)	n-3 polyunsaturated fatty acids (PUFA) 1 g daily versus placebo	Patients with NYHA classes II to IV heart failure, whatever the cause and the LVEF and already receiving optimized recommended therapy with no clear indication or contraindication to cholesterol-lowering therapy	double blind Italy
n3-PUFA-HF ongoing [NCT00149409] n=NA follow-up:	Omega-3-Polyunsaturated Fatty-Acids (EPH/DHA 1:1.2) versus placebo	Patients With Severe Chronic Heart Failure	Parallel groups double blind
Pikazol vs placebo			

continued...

Trial	Treatments	Patients	Trials design and methods
Bonnema , 1995 n=14/14 follow-up: 24 months	Pikazol fish oil capsules, 6x1 g/d (3.3g EPA + DHA) versus placebo (olive oil capsules, 6x1 g/d)	people with insulin treated diabetes and microalbuminureakK	Parallel groups double-blind Denmark
Promega vs placebo			
Connor , 1993 n=8/8 follow-up: 6 months	Promega oil, 15g/d (6g/d EPA + DHA) versus placebo (Olive oil, 15g/d)	people with non-insulin dependant diabetes and hypertiglyceridaemia	Parallel groups double-blind US
Sacks (HARP) , 1995 n=41/39 follow-up: 29 months	Promega capsules 12x1 g/d (6.0g EPA + DHA + DPA) versus placebo (olive oil capsules, 12x1 g/d)	people with angiographically documented CHD DPA)	Parallel groups double-blind US
Super EPA vs placebo			
Reis , 1991 n=146/72 follow-up: 6 months	Super EPA capsules 12x1 g/d (7.0g EPA + DHA + a-lin) ORPromega capsules 12x1 g/d (6.0g EPA + DHA + a-lin) versus placebo (olive oil capsules, 12x1 g/d)	people undergoing angioplasty	Parallel groups double blind US

References

Burr (DART 2), 2003:

Ness AR, Gallacher JE, Bennett PD, Gunnell DJ, Rogers PJ, Kessler D, Burr ML Advice to eat fish and mood: a randomised controlled trial in men with angina. *Nutr Neurosci* 2003 Feb;6:63-5 [[12608739](#)]

Burr (DART), 1989:

Burr ML, Fehily AM, Gilbert JF, Rogers S, Holliday RM, Sweetnam PM, Elwood PC, Deadman NM Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: diet and reinfarction trial (DART). *Lancet* 1989 Sep 30;2:757-61 [[2571009](#)]

Ness AR, Whitley E, Burr ML, Elwood PC, Smith GD, Ebrahim S The long-term effect of advice to eat more fish on blood pressure in men with coronary disease: results from the diet and reinfarction trial. *J Hum Hypertens* 1999 Nov;13:729-33 [[10578215](#)]

Bemelmans, 2002:

Bemelmans WJ, Broer J, Feskens EJ, Smit AJ, Muskiet FA, Lefrandt JD, Bom VJ, May JF, Meyboom-de Jong B Effect of an increased intake of alpha-linolenic acid and group nutritional education on cardiovascular risk factors: the Mediterranean Alpha-linolenic Enriched Groningen Dietary Intervention (MARGARIN) study. *Am J Clin Nutr* 2002 Feb;75:221-7 [[11815311](#)]

Bemelmans WJ, Broer J, de Vries JH, Hulshof KF, May JF, Meyboom-De Jong B Impact of Mediterranean diet education versus posted leaflet on dietary habits and serum cholesterol in a high risk population for cardiovascular disease. *Public Health Nutr* 2000 Sep;3:273-83 [[10979147](#)]

Brox, 2001:

Brox J, Olaussen K, Osterud B, Elvevoll EO, Bjornstad E, Brattebog G, Iversen H A long-term seal- and cod-liver-oil supplementation in hypercholesterolemic subjects. *Lipids* 2001 Jan;36:7-13 [[11214732](#)]

Franzen, 1993:

Franzen D, Geisel J, Hpp HW, Oette K, Hilger HH [Long-term effects of low dosage fish oil on serum lipids and lipoproteins] *Med Klin (Munich)* 1993;88:134-8 [[8474402](#)]

Shimizu, 1995:

Shimizu H, Ohtani K, Tanaka Y, Sato N, Mori M, Shimomura Y Long-term effect of eicosapentaenoic acid ethyl (EPA-E) on albuminuria of non-insulin dependent diabetic patients. *Diabetes Res Clin Pract* 1995;28:35-40 [[7587910](#)]

Bellamy, 1992:

Bellamy CM, Schofield PM, Faragher EB, Ramsdale DR Can supplementation of diet with omega-3 polyunsaturated fatty acids reduce coronary angioplasty restenosis rate? *Eur Heart J* 1992 Dec;13:1626-31 [[1289091](#)]

Dehmer, 1998:

Dehmer GJ, Popma JJ, van den Berg EK, Eichhorn EJ, Prewitt JB, Campbell WB, Jennings L, Willerson JT, Schmitz JM Reduction in the rate of early restenosis after coronary angioplasty by a diet supplemented with n-3 fatty acids. *N Engl J Med* 1988 Sep 22;319:733-40 [[2842680](#)]

Kaul, 1992:

Kaul U, Sanghvi S, Bahl VK, Dev V, Wasir HS Fish oil supplements for prevention of restenosis after coronary angioplasty. *Int J Cardiol* 1992 Apr;35:87-93 [[1563884](#)]

Eritsland, 1996:

Eritsland J, Arnesen H, Gronseth K, Fjeld NB, Abdelnoor M Effect of dietary supplementation with n-3 fatty acids on coronary artery bypass graft patency. *Am J Cardiol* 1996 Jan 1;77:31-6 [[8540453](#)]

Eritsland J, Arnesen H, Seljeflot I, Hostmark AT Long-term metabolic effects of n-3 polyunsaturated fatty acids in patients with coronary artery disease. *Am J Clin Nutr* 1995 Apr;61:831-6 [[7702027](#)]

GISSI-P, 1999:

Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto miocardico. *Lancet* 1999 Aug 7;354:447-55 [[10465168](#)]

Marchioli R, Barzi F, Bomba E, Chieffo C, Di Gregorio D, Di Mascio R, Franzosi MG, Geraci E, Levantesi G, Maggioni AP, Mantini L, Marfisi RM, Mastrogiuseppe G, Mininni N, Nicolosi GL, Santini M, Schweiger C, Tavazzi L, Tognoni G, Tucci C, Valagussa F Early protection against sudden death by n-3 polyunsaturated fatty acids after myocardial infarction: time-course analysis of the results of the Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico (GISSI)-Prevenzione. *Circulation* 2002 Apr 23;105:1897-903 [[11997274](#)]

OMEGA, 2009:

Senges Randomized Trial of Omega-3 Fatty Acids on Top of Modern Therapy After Acute Myocardial Infarction: The OMEGA Trial ACC.09/i2, Orlando, FL, March 2009 [0]

Rauch B, Schiele R, Schneider S, Diller F, Victor N, Gohlke H, Gottwik M, Steinbeck G, Del Castillo U, Sack R, Worth H, Katus H, Spitzer W, Sabin G, Seneges J OMEGA, a randomized, placebo-controlled trial to test the effect of highly purified omega-3 fatty acids on top of modern guideline-adjusted therapy after myocardial infarction. *Circulation* 2010;122:2152-9 [[21060071](#)] [10.1161/CIRCULATIONAHA.110.948562](#)

Milner, 1989:

Milner MR, Gallino RA, Leffingwell A, Pichard AD, Brooks-Robinson S, Rosenberg J, Little T, Lindsay J Jr Usefulness of fish oil supplements in preventing clinical evidence of restenosis after percutaneous transluminal coronary angioplasty. *Am J Cardiol* 1989 Aug 1;64:294-9 [[2526993](#)]

ALPHA OMEGA (ALA), 2010:

Kromhout D, Giltay EJ, Geleijnse JM n-3 Fatty Acids and Cardiovascular Events after Myocardial Infarction. *N Engl J Med* 2010 Nov 18;363:2015-2026 [[20929341](#)] [10.1056/NEJ-Moa1003603](#)

Natvig, 1968:

Maresta, 2002:

Skali H, Solomon SD, Pfeffer MA Are we asking too much of our trials? *Am Heart J* 2002 Jan;143:1-3 [[11773904](#)]

Sirtori, 1998:

Sirtori CR, Crepaldi G, Manzato E, Mancini M, Rivellese A, Paoletti R, Pazzucconi F, Pamparana F, Stragliotto E One-year treatment with ethyl esters of n-3 fatty acids in patients with hypertriglyceridemia and glucose intolerance: reduced triglyceridemia, total cholesterol and increased HDL-C without glyceimic alterations. *Atherosclerosis* 1998;137:419-27 [[9622285](#)]

Rossing, 1996:

Rossing P, Hansen BV, Nielsen FS, Myrup B, Hlmer G, Parving HH Fish oil in diabetic nephropathy. *Diabetes Care* 1996;19:1214-9 [[8908382](#)]

Borchgrevink, 1966:

Borchgrevink CF, Skaga E, Berg KJ, Skjaeggstad O Absence of prophylactic effect of linolenic acid in patients with coronary heart-disease. *Lancet* 1966 Jul 23;2:187-9 [[4161161](#)]

Leaf, 1994:

Leaf A, Jorgensen MB, Jacobs AK, Cote G, Schoenfeld DA, Scheer J, Weiner BH, Slack JD, Kellett MA, Raizner AE Do fish oils prevent restenosis after coronary angioplasty? *Circulation* 1994 Nov;90:2248-57 [[7955181](#)]

Mehta VY, Jorgensen MB, Raizner AE, Wolde-Tsadik G, Mahrer PR, Mansukhani P Spontaneous regression of restenosis: an angiographic study. *J Am Coll Cardiol* 1995 Sep;26:696-702 [[7642861](#)]

Sacks (TOHP 1), 1994:

The effects of nonpharmacologic interventions on blood pressure of persons with high normal levels. Results of the Trials of Hypertension Prevention, Phase I. *JAMA* 1992 Mar 4;267:1213-20 [[1586398](#)]

von Schacky, 1999:

von Schacky C, Angerer P, Kothny W, Theisen K, Mudra H The effect of dietary omega-3 fatty acids on coronary atherosclerosis. A randomized, double-blind, placebo-controlled trial. *Ann Intern Med* 1999;130:554-62 [[10189324](#)]

Hawthorne, 1992:

Hawthorne AB, Daneshmend TK, Hawkey CJ, Belluzzi A, Everitt SJ, Holmes GK, Malkinson C, Shaheen MZ, Willars JE Treatment of ulcerative colitis with fish oil supplementation: a prospective 12 month randomised controlled trial. *Gut* 1992;33:922-8 [[1353742](#)]

Bairati, 1992:

Bairati I, Roy L, Meyer F Double-blind, randomized, controlled trial of fish oil supplements in prevention of recurrence of stenosis after coronary angioplasty. *Circulation* 1992 Mar;85:950-6 [[1537131](#)]

Lau, 1993:

Lau CS, Morley KD, Belch JJ Effects of fish oil supplementation on non-steroidal anti-inflammatory drug requirement in patients with mild rheumatoid arthritis—a double-blind placebo controlled study. *Br J Rheumatol* 1993 Nov;32:982-9 [[8220938](#)]

Lau, 1995:

Lau CS, McLaren M, Belch JJ Effects of fish oil on plasma fibrinolysis in patients with mild rheumatoid arthritis. *Clin Exp Rheumatol* 1995 Jan-Feb;13:87-90 [[7774110](#)]

Nye, 1990:

Nye ER, Ablett MB, Robertson MC, Ilsley CD, Sutherland WH Effect of eicosapentaenoic acid on restenosis rate, clinical course and blood lipids in patients after percutaneous transluminal coronary angioplasty. *Aust N Z J Med* 1990 Aug;20:549-52 [[2222347](#)]

Singh, 1997:

Singh RB, Niaz MA, Sharma JP, Kumar R, Rastogi V, Moshiri M Randomized, double-blind, placebo-controlled trial of fish oil and mustard oil in patients with suspected acute myocardial infarction: the Indian experiment of infarct survival-4. *Cardiovasc Drugs Ther* 1997;11:485-91 [[9310278](#)]

Johansen, 1999:

Johansen O, Brekke M, Seljeflot I, Abdelnoor M, Arnesen H N-3 fatty acids do not prevent restenosis after coronary angioplasty: results from the CART study. *Coronary Angioplasty Restenosis Trial. J Am Coll Cardiol* 1999;33:1619-26 [[10334433](#)]

Nilsen, 2001:

Nilsen DW, Albrektsen G, Landmark K, Moen S, Aarsland T, Woie L Effects of a high-dose concentrate of n-3 fatty acids or corn oil introduced early after an acute myocardial infarction on serum triacylglycerol and HDL cholesterol. *Am J Clin Nutr* 2001 Jul;74:50-6 [[11451717](#)]

ALPHA OMEGA (EPA DHA), 2010:

Kromhout D, Giltay EJ, Geleijnse JM n-3 Fatty Acids and Cardiovascular Events after Myocardial Infarction. *N Engl J Med* 2010 Nov 18;363:2015-2026 [[20929341](#)] [10.1056/NEJ-Moa1003603](#)

Risk and Prevention Study, 2013:

Roncaglioni MC, Tombesi M, Avanzini F, Barlera S, Caimi V, Longoni P, Marzona I, Milani V, Silletta MG, Tognoni G, Marchioli R n-3 fatty acids in patients with multiple cardiovascular risk factors. *N Engl J Med* 2013 May 9;368:1800-8 [23656645] [10.1056/NEJMoa1205409](https://doi.org/10.1056/NEJMoa1205409)

Efficacy of n-3 polyunsaturated fatty acids and feasibility of optimizing preventive strategies in patients at high cardiovascular risk: rationale, design and baseline characteristics of the Rischio and Prevenzione study, a large randomised trial in general practice. *Trials* 2010;11:68 [20509875]

GISSI HF fatty acid, 2008:

n3-PUFA-HF, :

Bonnema, 1995:

Bonnema SJ, Jespersen LT, Marving J, Gregersen G. Supplementation with olive oil rather than fish oil increases small arterial compliance in diabetic patients. *Diabetes, Nutrition and Metabolism Clinical and Experimental* 1995;8:81-87

Connor, 1993:

Connor WE, Prince MJ, Ullmann D, Riddle M, Hatcher L, Smith FE, Wilson D The hypotriglyceridemic effect of fish oil in adult-onset diabetes without adverse glucose control. *Ann N Y Acad Sci* 1993 Jun 14;683:337-40 [8352456]

Sacks (HARP), 1995:

Sacks FM, Stone PH, Gibson CM, Silverman DI, Rosner B, Pasternak RC Controlled trial of fish oil for regression of human coronary atherosclerosis. HARP Research Group. *J Am Coll Cardiol* 1995 Jun;25:1492-8 [7759696]

Reis, 1991:

Reis GJ, Boucher TM, Sipperly ME, Silverman DI, McCabe CH, Baim DS, Sacks FM, Grossman W, Pasternak RC Randomised trial of fish oil for prevention of restenosis after coronary angioplasty. *Lancet* 1989 Jul 22;2:177-81 [2568519]

2 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.