

# Clinical trials of angiotensin-receptor blockers for hypertension in all diseases requiring ACEi (HF, CHD, HT,...)

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 angiotensin receptor blocker

Trial	Treatments	Patients	Trials design and methods
<b>ARBs vs control</b>			
<b>Suzuki , 2008</b> n=183/183 follow-up:	ARBs (valsartan, candesartan, and losartan) versus no ARBs	patients with diabetes and chronic kidney disease on dialysis	Parallel groups open
<b>candesartan vs control</b>			
<b>Takahashi , 2006</b> n=43/37 follow-up: 19.4 months	candesartan versus control	patients on chronic haemodialysis in stable condition and with no clinical evidence of cardiac disorders	Parallel groups open
<b>candesartan vs conventional treatment</b>			
<b>E-COST , 2005</b> n=1053/995 follow-up:	candesartan, 2 to 12 mg daily versus conventional antihypertensive drugs other than angiotensin converting enzyme inhibitors or ARBs	Japanese essential hypertensive subjects (sitting blood pressure 140-180/90-110 mmHg) aged 35-79 years	Parallel groups single-blind Japan
<b>E-COST-R , 2005</b> n=69/72 follow-up:	candesartan versus conventional treatment	hypertensive subjects 60 to 75 years old with non-diabetic chronic renal insufficiency	Parallel groups open
<b>HIJ-CREATE , 2009</b> n=1024/1025 follow-up: 4.2 y (median)	angiotensin II receptor blocker-based therapy versus non-angiotensin II receptor blocker-based therapy	patients with angiographically documented coronary artery disease and hypertension	Parallel groups open Japan
<b>candesartan vs placebo</b>			
<b>SCOPE , 2003</b> n=2477/2460 follow-up: 3.7 y (mean)	candesartan, 816 mg once daily (target 160/90) versus placebo	patients aged 70-89 years, with systolic blood pressure 160-179 mmHg, and/or diastolic blood pressure 90-99 mmHg, and a Mini Mental State Examination (MMSE) test score >24	Parallel groups double-blind 15 countries
<b>irbesartan vs placebo</b>			
<b>IDNT (irbesartan vs pbo) , 2001</b> n=579/569 follow-up: 2.6 y	Irbesartan 300mg/d (target 135/85) versus placebo	hypertensive patients with nephropathy due to type 2 diabetes	Parallel groups double-blind worldwide

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>IRMA 2 , 2001</b> n=404/207 follow-up: 2 years	irbesartan 150 mg daily or 300 mg daily versus placebo	hypertensive patients with type 2 diabetes and microalbuminuria	Parallel groups double-blind multinational
<b>losartan vs placebo</b>			
<b>RENAAL , 2001</b> n=751/762 follow-up: 3.4 years	Losartan 50 to 100 mg once daily versus placebo	patients with type 2 diabetes and nephropathy	Parallel groups double-blind
<b>telmisartan vs placebo</b>			
<b>PROPHESSE , 2008</b> [NCT00153062] n=10146/10186 follow-up: 2.5 y	telmisartan 80 mg daily versus placebo	patients who recently had an ischemic stroke	Factorial plan double blind 35 countries
<b>Candesartan vs usual care</b>			
<b>HIJ-CREATE , 2009</b> n=1025/1024 follow-up: up to 60 months	cardesartan adjusted dose for target arterial pressure of <130/85 mmHg versus usual care (non-ARB-based pharmacotherapy including angiotensin-converting enzyme-inhibitors)	hypertension with angiographically documented coronary artery disease (acute or stable)	Parallel groups open japan
<b>candesartan vs amlodipine</b>			
<b>CASE-J , 2008</b> n=2354/2349 follow-up: 3.2 years	candesartan-based regimen versus amlodipine-based regimen	high-risk Japanese hypertensive patients	Parallel groups open (blinded assessment) Japan
<b>irbesartan vs amlodipine</b>			
<b>IDNT (irbesartan vs amlodipine) , 2001</b> n=579/567 follow-up: 26y	Irbesartan 300mg/d (with a target of 135/85) versus amlodipine 10mg/d (with a target of 135/85)	hypertensive patients with nephropathy due to type 2 diabetes	Parallel groups double-blind worldwide
<b>valsartan vs amlodipine</b>			
<b>VALUE , 2004</b> [NCT00129233] n=7649/7596 follow-up: 4.2 y (mean)	valsartan based regimen versus amlodipine based regimen	patients, aged 50 years or older with treated or untreated hypertension and high risk of cardiac events	Parallel groups Double blind 31 countries
<b>losartan vs atenolol</b>			
<b>LIFE , 2002</b> n=4605/4588 follow-up: 4.8 y (mean)	losartan versus atenolol	patients aged 55-80 years, with previously treated or untreated hypertension (sitting blood pressure 160/200/95/115 mm Hg) and ECG signs of LVH.	Parallel groups Double blind USA, Europe
<b>telmisartan vs enalapril</b>			
<b>DETAIL , 2004</b> n=120/130 follow-up: 5 year	telmisartan 80 mg daily versus enalapril 20 mg daily	subjects with type 2 diabetes and early nephropathy	Parallel groups double-blind

continued...

Trial	Treatments	Patients	Trials design and methods
<b>DETAIL , 2004</b> n=120/130 follow-up: 5 y	Telmisartan 80 mg daily versus Enalapril 20 mg daily	patients with type 2 diabetes and early nephropathy	Parallel groups double-blind
<b>candesartan vs hydrochlorothiazide</b>			
<b>ALPINE , 2003</b> n=197/196 follow-up: 1 year	candesartan versus hydrochlorothiazide	newly detected hypertensives	Parallel groups double-blind Sweden
<b>olmesartan 40 mg vs olmesartan 20 mg plus a calcium-channel blocker</b>			
<b>OSCAR , 2011</b> [NCT00134160] n=578/586 follow-up:	high-dose olmesartan 40 mg per day versus 20-mg/day olmesartan comined with standard dose of amlodipine or azelnidipine	high-risk elderly Japanese hypertension patients	Parallel groups Japan

## References

### Suzuki, 2008:

Suzuki H, Kanno Y, Sugahara S, Ikeda N, Shoda J, Takenaka T, Inoue T, Araki R Effect of angiotensin receptor blockers on cardiovascular events in patients undergoing hemodialysis: an open-label randomized controlled trial. *Am J Kidney Dis* 2008;52:501-6 [[18653268](#)] [10.1053/j.ajkd.2008.04.031](#)

### Takahashi, 2006:

Takahashi A, Takase H, Toriyama T, Sugiura T, Kurita Y, Ueda R, Dohi Y Candesartan, an angiotensin II type-1 receptor blocker, reduces cardiovascular events in patients on chronic haemodialysis—a randomized study. *Nephrol Dial Transplant* 2006;21:2507-12 [[16766543](#)] [10.1093/ndt/gff293](#)

### E-COST, 2005:

Suzuki H, Kanno Y Effects of candesartan on cardiovascular outcomes in Japanese hypertensive patients. *Hypertens Res* 2005;28:307-14 [[16138560](#)] [10.1291/hypres.28.307](#)

### E-COST-R, 2005:

Nakamura T, Kanno Y, Takenaka T, Suzuki H An angiotensin receptor blocker reduces the risk of congestive heart failure in elderly hypertensive patients with renal insufficiency. *Hypertens Res* 2005;28:415-23 [[16156505](#)] [10.1291/hypres.28.415](#)

### HIJ-CREATE, 2009:

Kasanuki H, Hagiwara N, Hosoda S, Sumiyoshi T, Honda T, Haze K, Nagashima M, Yamaguchi J, Origasa H, Urashima M, Ogawa H Angiotensin II receptor blocker-based vs. non-angiotensin II receptor blocker-based therapy in patients with angiographically documented coronary artery disease and hypertension: the Heart Institute of Japan Candesartan Randomized Trial for Evaluation in Coronary Artery Disease (HIJ-CREATE). *Eur Heart J* 2009;30:1203-12 [[19346521](#)] [10.1093/eurheartj/ehp101](#)

### SCOPE, 2003:

Saxby BK, Harrington F, Wesnes KA, McKeith IG, Ford GA Candesartan and cognitive decline in older patients with hypertension: a substudy of the SCOPE trial. *Neurology* 2008;70:1858-66 [[18458219](#)] [10.1212/01.wnl.0000311447.85948.78](#)

Lithell H, Hansson L, Skoog I, Elmfeldt D, Hofman A, Olofsson B, Trenkwalder P, Zanchetti A The Study on Cognition and Prognosis in the Elderly (SCOPE): principal results of a randomized double-blind intervention trial. *J Hypertens* 2003;21:875-86 [[12714861](#)] [10.1097/01.hjh.0000059028.82022.89](#)

### IDNT (irbesartan vs pbo), 2001:

Lewis EJ, Hunsicker LG, Clarke WR, Berl T, Pohl MA, Lewis JB, Ritz E, Atkins RC, Rohde R, Raz I Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001;345:851-60 [[11565517](#)]

Lewis EJ, Hunsicker LG, Clarke WR, Berl T, Pohl MA, Lewis JB, Ritz E, Atkins RC, Rohde R, Raz I Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001;345:851-60 [[11565517](#)]

### IRMA 2, 2001:

Parving HH, Lehnert H, Brchner-Mortensen J, Gomis R, Andersen S, Arner P The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. *N Engl J Med* 2001;345:870-8 [[11565519](#)] [10.1056/NEJMoa011489](#)

**RENAAL, 2001:**

Brenner BM, Cooper ME, de Zeeuw D, Keane WF, Mitch WE, Parving HH, Remuzzi G, Snapinn SM, Zhang Z, Shahinfar S Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001;345:861-9 [[11565518](#)]

**PROPELLS, 2008:**

Yusuf S, Diener HC, Sacco RL, Cotton D, Ounpuu S, Lawton WA, Palesch Y, Martin RH, Albers GW, Bath P, Bornstein N, Chan BP, Chen ST, Cunha L, Dahlf B, De Keyser J, Donnan GA, Estol C, Gorelick P, Gu V, Hermansson K, Hilbrich L, Kaste M, Lu C, Machnig T, — *N Engl J Med* 2008;359:1225-37 [[18753639](#)] [10.1056/NEJMoa0804593](#)

**HIJ-CREATE, 2009:**

Kasanuki H, Hagiwara N, Hosoda S, Sumiyoshi T, Honda T, Haze K, Nagashima M, Yamaguchi J, Origasa H, Urashima M, Ogawa H Angiotensin II receptor blocker-based vs. non-angiotensin II receptor blocker-based therapy in patients with angiographically documented coronary artery disease and hypertension: the Heart Institute of Japan Candesartan Randomized Trial for Evaluation in Coronary Artery Disease (HIJ-CREATE). *Eur Heart J* 2009;30:1203-12 [[19346521](#)] [10.1093/eurheartj/ehp101](#)

**CASE-J, 2008:**

Ogihara T, Nakao K, Fukui T, Fukiyama K, Ueshima K, Oba K, Sato T, Saruta T Effects of candesartan compared with amlodipine in hypertensive patients with high cardiovascular risks: candesartan antihypertensive survival evaluation in Japan trial. *Hypertension* 2008 Feb;51:393-8 [[18172059](#)] [10.1161/HYPERTENSIONAHA.107.098475](#)

**IDNT (irbesartan vs amlodipine), 2001:**

Lewis EJ, Hunsicker LG, Clarke WR, Berl T, Pohl MA, Lewis JB, Ritz E, Atkins RC, Rohde R, Raz I Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001;345:851-60 [[11565517](#)]

**VALUE, 2004:**

Julius S, Kjeldsen SE, Weber M, Brunner HR, Ekman S, Hansson L, Hua T, Laragh J, McInnes GT, Mitchell L, Plat F, Schork A, Smith B, Zanchetti A Outcomes in hypertensive patients at high cardiovascular risk treated with regimens based on valsartan or amlodipine: the VALUE randomised trial. *Lancet* 2004 Jun 19;363:2022-31 [[15207952](#)]

**LIFE, 2002:**

Dahlof B, Devereux RB, Kjeldsen SE, Julius S, Beevers G, de Faire U, Fyhrquist F, Ibsen H, Kristiansson K, Lederballe-Pedersen O, Lindholm LH, Nieminen MS, Omvik P, Oparil S, Wedel H Cardiovascular morbidity and mortality in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet* 2002 Mar 23;359:995-1003 [[11937178](#)]

**DETAIL, 2004:**

Barnett AH, Bain SC, Bouter P, Karlberg B, Madsbad S, Jervell J, Mustonen J Angiotensin-receptor blockade versus converting-enzyme inhibition in type 2 diabetes and nephropathy. *N Engl J Med* 2004;351:1952-61 [[15516696](#)] [10.1056/NEJMoa042274](#)

**DETAIL, 2004:**

Barnett AH, Bain SC, Bouter P, Karlberg B, Madsbad S, Jervell J, Mustonen J Angiotensin-receptor blockade versus converting-enzyme inhibition in type 2 diabetes and nephropathy. *N Engl J Med* 2004;351:1952-61 [[15516696](#)]

**ALPINE, 2003:**

Lindholm LH, Persson M, Alaupovic P, Carlberg B, Svensson A, Samuelsson O Metabolic outcome during 1 year in newly detected hypertensives: results of the Antihypertensive Treatment and Lipid Profile in a North of Sweden Efficacy Evaluation (ALPINE study). *J Hypertens* 2003;21:1563-74 [[12872052](#)] [10.1097/01.hjh.0000084723.53355.76](#)

**OSCAR, 2011:**

Ogawa H, Kim-Mitsuyama S, Jinnouchi T, Matsui K, Arakawa K Rationale, design and patient baseline characteristics of OlmeSartan and calcium antagonists randomized (OSCAR) study: a study comparing the incidence of cardiovascular events between high-dose angiotensin II receptor blocker (ARB) monotherapy and combination therapy of ARB with calcium channel blocker in Japanese elderly high-risk hypertensive patients (ClinicalTrials.gov no. NCT00134160). *Hypertens Res* 2009;32:575-80 [[19444280](#)] [10.1038/hr.2009.60](#)

## 2 ARBs vs ACEIs

Trial	Treatments	Patients	Trials design and methods
<b>Losartan vs Captopril</b>			
<b>ELITE-II , 2000</b> n=1578/1574 follow-up: 1.5 y	Losartan titrated to 50 mg once daily versus Captopril titrated to 50 mg three times daily	patients aged 60 years or older with New York Heart Association class II-IV heart failure and ejection fraction of 40% or less. Patients	Parallel groups double-blind
<b>OPTIMAAL , 2001</b> n=2744/2733 follow-up: 2.7 y	losartan (titrated to 50 mg once daily) versus Captopril (titrated to 50 mg three times daily)	patients 50 years of age or older, with confirmed acute myocardial infarction and heart failure during the acute phase or a new Q-wave anterior infarction or reinfarction	Parallel groups NA Europe (7 countries)
<b>ELITE , 1997</b> n=352/370 follow-up: 65279;1 y	65279;Losartan titrated to 50 mg once daily for 48 weeks versus 65279;Captopril titrated to 50 mg three times daily for 48 weeks	naive patients (aged 65 years or more) with NYHA class II-IV heart failure and ejection fractions of 40% or less	Parallel groups double-blind
<b>Valsartan vs Captopril</b>			
<b>VALIANT/Val , 2003</b> n=4909/4909 follow-up: 2.1 y	Valsartan versus Captopril	patients with myocardial infarction complicated by left ventricular systolic dysfunction, heart failure, or both	Parallel groups double-blind
<b>Telmisartan vs Enalapril</b>			
<b>DETAIL , 2004</b> n=120/130 follow-up: 5 year	telmisartan 80 mg daily versus enalapril 20 mg daily	subjects with type 2 diabetes and early nephropathy	Parallel groups double-blind
<b>DETAIL , 2004</b> n=120/130 follow-up: 5 y	Telmisartan 80 mg daily versus Enalapril 20 mg daily	patients with type 2 diabetes and early nephropathy	Parallel groups double-blind
<b>Telmisartan vs Ramipril</b>			
<b>ONTARGET/Tel , 2008</b> n=8542/8576 follow-up: 4.7 y	Telmisartan 80 mg daily versus Ramipril 10 mg daily	patients with vascular disease or high-risk diabetes	Parallel groups double-blind

## References

### ELITE-II, 2000:

Pitt B, Poole-Wilson PA, Segal R, Martinez FA, Dickstein K, Camm AJ, Konstam MA, Riegger G, Klingner GH, Neaton J, Sharma D, Thiyagarajan B Effect of losartan compared with captopril on mortality in patients with symptomatic heart failure: randomised trial—the Losartan Heart Failure Survival Study ELITE II. *Lancet* 2000;355:1582-7 [[10821361](#)]

### OPTIMAAL, 2001:

Dickstein K, Kjekshus J Effects of losartan and captopril on mortality and morbidity in high-risk patients after acute myocardial infarction: the OPTIMAAL randomised trial. *Optimal Trial in Myocardial Infarction with Angiotensin II Antagonist Losartan*. *Lancet* 2002;360:752-60 [[12241832](#)]

### ELITE, 1997:

Pitt B, Segal R, Martinez FA, Meurers G, Cowley AJ, Thomas I, Deedwania PC, Ney DE, Snively DB, Chang PI Randomised trial of losartan versus captopril in patients over 65 with heart failure (Evaluation of Losartan in the Elderly Study, ELITE) *Lancet* 1997;349:747-52 [[9074572](#)]

### VALIANT/Val, 2003:

Pfeffer MA, McMurray JJ, Velazquez EJ, Rouleau JL, Kber L, Maggioni AP, Solomon SD, Swedberg K, Van de Werf F, White H, Leimberger JD, Henis M, Edwards S, Zelenkofske S, Sellers MA, Califf RM Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *N Engl J Med* 2003;349:1893-906 [[14610160](#)]

McMurray J, Solomon S, Pieper K, Reed S, Rouleau J, Velazquez E, White H, Howlett J, Swedberg K, Maggioni A, Kber L, Van de Werf F, Califf R, Pfeffer M The effect of valsartan, captopril, or both on atherosclerotic events after acute myocardial infarction: an analysis of the Valsartan in Acute Myocardial Infarction Trial (VALIANT). J Am Coll Cardiol 2006;47:726-33 [16487836]

**DETAIL, 2004:**

Barnett AH, Bain SC, Bouter P, Karlberg B, Madsbad S, Jervell J, Mustonen J Angiotensin-receptor blockade versus converting-enzyme inhibition in type 2 diabetes and nephropathy. N Engl J Med 2004;351:1952-61 [15516696] 10.1056/NEJMoa042274

**DETAIL, 2004:**

Barnett AH, Bain SC, Bouter P, Karlberg B, Madsbad S, Jervell J, Mustonen J Angiotensin-receptor blockade versus converting-enzyme inhibition in type 2 diabetes and nephropathy. N Engl J Med 2004;351:1952-61 [15516696]

**ONTARGET/Tel, 2008:**

Yusuf S, Teo KK, Pogue J, Dyal L, Copland I, Schumacher H, Dagenais G, Sleight P, Anderson C Telmisartan, ramipril, or both in patients at high risk for vascular events. N Engl J Med 2008;358:1547-59 [18378520]

Verdecchia P, Sleight P, Mancia G, Fagard R, Trimarco B, Schmieder RE, Kim JH, Jennings G, Jansky P, Chen JH, Liu L, Gao P, Probstfeld J, Teo K, Yusuf S Effects of telmisartan, ramipril, and their combination on left ventricular hypertrophy in individuals at high vascular risk in the Ongoing Telmisartan Alone and in Combination With Ramipril Global End Point Trial and the Telmisartan Randomized Assessment Study in ACE Intolerant Subjects With Cardiovascular Disease. Circulation 2009;120:1380-9 [19770395]

### 3 ARBs+ACEIs vs ACEIs

Trial	Treatments	Patients	Trials design and methods
<b>Valsartan + captopril vs Captopril</b>			
VALIANT/Val+Cap , 2003 n=4885/4909 follow-up: 2.1 y	Valsartan + captopril versus Captopril	patients with myocardial infarction complicated by left ventricular systolic dysfunction, heart failure, or both	Parallel groups double-blind
<b>Telmisartan + ramipril vs Ramipril</b>			
ONTARGET/Tel+Ram , 2008 n=8502/8576 follow-up: 4.7 y	Telmisartan + ramipril versus Ramipril	patients with vascular disease or high-risk diabetes	Parallel groups double-blind

### References

**VALIANT/Val+Cap, 2003:**

Pfeffer MA, McMurray JJ, Velazquez EJ, Rouleau JL, Kber L, Maggioni AP, Solomon SD, Swedberg K, Van de Werf F, White H, Leimberger JD, Henis M, Edwards S, Zelenkofske S, Sellers MA, Califf RM Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. N Engl J Med 2003;349:1893-906 [14610160]

McMurray J, Solomon S, Pieper K, Reed S, Rouleau J, Velazquez E, White H, Howlett J, Swedberg K, Maggioni A, Kber L, Van de Werf F, Califf R, Pfeffer M The effect of valsartan, captopril, or both on atherosclerotic events after acute myocardial infarction: an analysis of the Valsartan in Acute Myocardial Infarction Trial (VALIANT). J Am Coll Cardiol 2006;47:726-33 [16487836]

**ONTARGET/Tel+Ram, 2008:**

Yusuf S, Teo KK, Pogue J, Dyal L, Copland I, Schumacher H, Dagenais G, Sleight P, Anderson C Telmisartan, ramipril, or both in patients at high risk for vascular events. N Engl J Med 2008;358:1547-59 [18378520]

Verdecchia P, Sleight P, Mancia G, Fagard R, Trimarco B, Schmieder RE, Kim JH, Jennings G, Jansky P, Chen JH, Liu L, Gao P, Probstfield J, Teo K, Yusuf S Effects of telmisartan, ramipril, and their combination on left ventricular hypertrophy in individuals at high vascular risk in the Ongoing Telmisartan Alone and in Combination With Ramipril Global End Point Trial and the Telmisartan Randomized Assessment Study in ACE Intolerant Subjects With Cardiovascular Disease. *Circulation* 2009;120:1380-9 [[19770395](#)]

## 4 endopeptidase inhibitors

Trial	Treatments	Patients	Trials design and methods
<b>LCZ696 vs placebo</b>			
Ruilope , 2010 n=NA follow-up: 8 weeks	LCZ696 for 8 weeks versus placebo	patients with mild to moderate hypertension	Parallel groups double blind 18 countries

## References

### Ruilope, 2010:

Ruilope LM, Dukat A, Bhm M, Lacourcire Y, Gong J, Lefkowitz MP Blood-pressure reduction with LCZ696, a novel dual-acting inhibitor of the angiotensin II receptor and neprilysin: a randomised, double-blind, placebo-controlled, active comparator study. *Lancet* 2010 Mar 15;: [[20236700](#)] [10.1016/S0140-6736\(09\)61966-8](#)

## 5 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.