

Clinical trials of calcium channel blockers for acute myocardial infarction in acute short term and post MI studies

TrialResults-center www.trialresultscenter.org

1 acute and long term calcium blockers

Trial	Treatments	Patients	Trials design and methods
nifedipine vs placebo			
SPRINT II , 1993 n=682/676 follow-up: 6 months	Nifedipine 20mgx3 started within 3hr of hospital admission for 6 months versus placebo	men and women with suspected acute MI	Parallel groups Double blind Israel
Sirnes , 1984 n=112/115 follow-up: 42 days	Nifedipine 5x10mg versus placebo	patients with suspected acute myocardial infarction	Parallel groups double blind
Gottlieb , 1988 n=64/68 follow-up: 42 days	Nifedipine 10 to 30mgx4 versus placebo	-	Parallel groups double blind
Eisenberg , 1985 n=25/25 follow-up: 12 days	Nifedipine 20mgx4 versus placebo	patients with nontransmural myocardial infarction	Parallel groups double blind
Bradagan , 1986 n=60/61 follow-up: 2 days	Nifedipine 10mgx4 versus placebo	patients with suspected acute myocardial infarction within 6hrs	Parallel groups double blind
Loogna , 1985 n=23/34 follow-up:	Nifedipine 4x10mg versus placebo	patients with acute myocardial infarction within 6 h from onset of symptoms	Parallel groups double blind
TRENT (Wilcox) , 1986 n=2240/2251 follow-up: 28 days	Nifedipine 4x10mg versus placebo	patients with suspected myocardial infarction	Parallel groups double blind
Walker , 1988 n=217/217 follow-up:	Nifedipine 6x10mg for 48hrs versus placebo	patients with suspected myocardial infarction within six hours from the onset of chest pain	Parallel groups double blind
Jaffe , 1987 n=13/9 follow-up: 7 days	Nifedipine 6x20mg versus placebo	patients with acute myocardial infarction	Parallel groups single blind
Erbel , 1988 n=74/75 follow-up: hospital stay	Nifedipine 20mgx3 versus placebo	patients with acute myocardial infarction	Parallel groups double blind
SPRINT I , 1988 n=1130/1146 follow-up: 1 year	Nifedipine 10mgx3 versus placebo	patient surviving MI 7 and 21 days after admission	Parallel groups Double blind Israel

continued...

Trial	Treatments	Patients	Trials design and methods
verapamil vs placebo			
DAVIT I , 1984 n=1751/1747 follow-up: 6 months	verapamil 120mgx3 versus placebo	-	Parallel groups Double blind Danish
Crea , 1985 n=8/9 follow-up: 10 days	verapamil 80mgx3 versus placebo	patients admitted to the coronary care unit with transmural acute myocardial infarction	Parallel groups single blind
CRIS , 1996 n=531/542 follow-up: 23.5 months	verapamil retard 360 mg daily versus placebo	patients admitted for acute myocardial infarction	Parallel groups Double blind Italy
DAVIT II , 1990 n=878/897 follow-up: 16 months	verapamil 120mgx3 for 18 months versus placebo	patients <76years with diagnosis of acute MI	Parallel groups Double blind Danish
Danish study , 1984 n=1729/1718 follow-up: 12 months	verapamil 0.1mg/kg IV plus 3x120mg orally versus placebo	patients under 75 years of age admitted to the CCU with a suspicion of acute myocardial infarction	Parallel groups double blind Danish

References

SPRINT II, 1993:

Goldbourt U, Behar S, Reicher-Reiss H, Zion M, Mandelzweig L, Kaplinsky E Early administration of nifedipine in suspected acute myocardial infarction. The Secondary Prevention Reinfarction Israel Nifedipine Trial 2 Study Arch Intern Med 1993;153:345-53 [[8285927](#)]

Sirnes, 1984:

Sirnes PA, Overskeid K, Pedersen TR, Bathen J, Drivenes A, Frland GS, Kjekshus JK, Landmark K, Rokseth R, Sirnes KE Evolution of infarct size during the early use of nifedipine in patients with acute myocardial infarction: the Norwegian Nifedipine Multicenter Trial. Circulation 1984;70:638-44 [[6383655](#)]

Gottlieb, 1988:

Gottlieb SO, Becker LC, Weiss JL, Shapiro EP, Chandra NC, Flaherty JT, Gottlieb SH, Ouyang P, Mellits ED, Townsend SN Nifedipine in acute myocardial infarction: an assessment of left ventricular function, infarct size, and infarct expansion. A double blind, randomised, placebo controlled trial Br Heart J 1988;59:411-8 [[3285878](#)]

Eisenberg, 1985:

Eisenberg PR, Lee RG, Biello DR, Geltman EM, Jaffe AS Chest pain after nontransmural infarction: the absence of remediable coronary vasospasm. Am Heart J 1985;110:515-21 [[3898794](#)]

Bradagan, 1986:

Branagan JP, Walsh K, Kelly P, Collins WC, McCafferty D, Walsh MJ Effect of early treatment with nifedipine in suspected acute myocardial infarction. Eur Heart J 1986;7:859-65 [[3792347](#)]

Loogna, 1985:

Loogna E, Sylvn C, Groth T, Mogensen L Complexity of enzyme release during acute myocardial infarction in a controlled study with early nifedipine treatment. Eur Heart J 1985;6:114-9 [[3891346](#)]

TRENT (Wilcox), 1986:

Wilcox RG, Hampton JR, Banks DC, Birkhead JS, Brooksby IA, Burns-Cox CJ, Hayes MJ, Joy MD, Malcolm AD, Mather HG Trial of early nifedipine in acute myocardial infarction: the Trent study. Br Med J (Clin Res Ed) 1986;293:1204-8 [[3096431](#)]

Walker, 1988:

Walker LJ, MacKenzie G, Adgey AA Effect of nifedipine on enzymatically estimated infarct size in the early phase of acute myocardial infarction. Br Heart J 1988;59:403-10 [3285877]

Jaffe, 1987:

Jaffe AS, Biello DR, Sobel BE, Geltman EM Enhancement of metabolism of jeopardized myocardium by nifedipine Int J Cardiol 1987;15:77-89 [3553039]

Erbel, 1988:

Erbel R, Pop T, Meinertz T, Olshausen KV, Treese N, Henrichs KJ, Schuster CJ, Rupprecht HJ, Schlurmann W, Meyer J Combination of calcium channel blocker and thrombolytic therapy in acute myocardial infarction Am Heart J 1988;115:529-38 [3278574]

SPRINT I, 1988:

Secondary prevention reinfarction Israeli nifedipine trial (SPRINT). A randomized intervention trial of nifedipine in patients with acute myocardial infarction. The Israeli Sprint Study Group Eur Heart J 1988;9:354-64 [2898363]

Disegni E, Goldbourt U, Reicher-Reiss H, Kaplinsky E, Zion M, Boyko V, Behar S The predictive value of admission heart rate on mortality in patients with acute myocardial infarction. SPRINT Study Group. Secondary Prevention Reinfarction Israeli Nifedipine Trial J Clin Epidemiol 1995;48:1197-205 [7561981]

DAVIT I, 1984:

Verapamil in acute myocardial infarction. The Danish Study Group on Verapamil in Myocardial Infarction Eur Heart J 1984;5:516-28 [6383832]

Hansen JF Treatment with verapamil after an acute myocardial infarction. Review of the Danish studies on verapamil in myocardial infarction (DAVIT I and II) Drugs 1991;42 Suppl 2:43-53 [1718701]

Crea, 1985:

Crea F, Deanfield J, Crean P, Sharom M, Davies G, Maseri A Effects of verapamil in preventing early postinfarction angina and reinfarction Am J Cardiol 1985;55:900-4 [3920891]

CRIS, 1996:

Rengo F, Carbonin P, Pahor M, DeCaprio L, Bernabei R, Ferrara N, Carosella L, Acanfora D, Parlati S, Vitale D A controlled trial of verapamil in patients after acute myocardial infarction: results of the calcium antagonist reinfarction Italian study (CRIS) Am J Cardiol 1996;77:365-9 [8602564]

Ferrara N Giorn geront 1996;44:577-583

DAVIT II, 1990:

Effect of verapamil on mortality and major events after acute myocardial infarction (the Danish Verapamil Infarction Trial II-DAVIT II) Am J Cardiol 1990;66:779-85 [2220572]

Vaage-Nilsen M, Rasmussen V Effect of verapamil on heart rate variability after an acute myocardial infarction. Danish Verapamil Infarction Trial II Cardiovasc Drugs Ther 1998;12:285-90 [9784908]

Vaage-Nilsen M, Rasmussen V, Hansen JF, Hagerup L, Sorensen MB, Pedersen-Bjergaard O, Mellempgaard K, Hollander NH, Nielsen I, Sigurd BM Effect of verapamil on arrhythmias and heart rate during 16 months following an acute myocardial infarction. The Danish Study Group on Verapamil in Myocardial Infarction Cardiovasc Drugs Ther 1994;8:147-51 [8086325]

Danish study, 1984:

Verapamil in acute myocardial infarction. The Danish Study Group on Verapamil in Myocardial Infarction. Eur Heart J 1984;5:516-28 [6383832]

2 acute short term calcium blockers

Trial	Treatments	Patients	Trials design and methods
diltiazem vs			
Machecourt , 1986 n=38/37 follow-up: 21 days	-	-	Parallel groups single blind

continued...

Trial	Treatments	Patients	Trials design and methods
nifedipine vs			
Muller , 1984 n=93/88 follow-up: 14 days	-	-	Parallel groups double blind
Gordon , 1984 n=13/13 follow-up: 1 day	-	-	Parallel groups single blind
MI study group , 1979 n=904/888	-	-	Parallel groups open
Tiapamil vs control			
Eichler , 1985 n=16/16 follow-up:	Tiapamil 0.5-1 mg/kg plus 25mg/kg/min IV versus no treatment	-	Parallel groups double blind
verapamil vs control			
Bussman , 1984 n=29/25 follow-up: 2 days	Verapamil 5 to 10 mg/h IV versus no treatment	-	Parallel groups open
diltiazem vs placebo			
Gibson , 1986 n=287/289 follow-up: 14 days	Diltiazem 90mgx4 versus placebo	-	Parallel groups double blind
Zannad , 1988 n=17/17 follow-up:	Diltiazem 10-20mg/h IV, 4x60mg orraly versus placebo	-	Parallel groups double blind
MDPIT , 1988 n=1234/1232 follow-up: 25 months (at least 12 months)	Diltiazem 60mgx4 versus placebo	Patient aged 25 to 75 years, admitted to coronary care units with a documented acute myocardial infarction	Parallel groups Double blind US, Canada
nifedipine vs placebo			
SPRINT II , 1993 n=682/676 follow-up: 6 months	Nifedipine 20mgx3 started within 3hr of hopsital admission for 6 months versus placebo	men and women with suspected acute MI	Parallel groups Double blind Israel
Sirnes , 1984 n=112/115 follow-up: 42 days	Nifedipine 5x10mg versus placebo	patients with suspected acute myocardial infarction	Parallel groups double blind
Gottlieb , 1988 n=64/68 follow-up: 42 days	Nifedipine 10 to 30mgx4 versus placebo	-	Parallel groups double blind
Eisenberg , 1985 n=25/25 follow-up: 12 days	Nifedipine 20mgx4 versus placebo	patients with nontransmural myocardial infarction	Parallel groups double blind

continued...

Trial	Treatments	Patients	Trials design and methods
Bradagan , 1986 n=60/61 follow-up: 2 days	Nifedipine 10mgx4 versus placebo	patients with suspected acute myocardial infarction within 6hrs	Parallel groups double blind
Loogna , 1985 n=23/34 follow-up:	Nifedipine 4x10mg versus placebo	patients with acute myocardial infarction within 6 h from onset of symptoms	Parallel groups double blind
TRENT (Wilcox) , 1986 n=2240/2251 follow-up: 28 days	Nifedipine 4x10mg versus placebo	patients with suspected myocardial infarction	Parallel groups double blind
Walker , 1988 n=217/217 follow-up:	Nifedipine 6x10mg for 48hrs versus placebo	patients with suspected myocardial infarction within six hours from the onset of chest pain	Parallel groups double blind
Jaffe , 1987 n=13/9 follow-up: 7 days	Nifedipine 6x20mg versus placebo	patients with acute myocardial infarction	Parallel groups single blind
Erbel , 1988 n=74/75 follow-up: hospital stay	Nifedipine 20mgx3 versus placebo	patients with acute myocardial infarction	Parallel groups double blind
SPRINT I , 1988 n=1130/1146 follow-up: 1 year	Nifedipine 10mgx3 versus placebo	patient surviving MI 7 and 21 days after admission	Parallel groups Double blind Israel
verapamil vs placebo			
DAVIT I , 1984 n=1751/1747 follow-up: 6 months	verapamil 120mgx3 versus placebo	-	Parallel groups Double blind Danish
Crea , 1985 n=8/9 follow-up: 10 days	verapamil 80mgx3 versus placebo	patients admitted to the coronary care unit with transmural acute myocardial infarction	Parallel groups single blind
CRIS , 1996 n=531/542 follow-up: 23.5 months	verapamil retard 360 mg daily versus placebo	patients admitted for acute myocardial infarction	Parallel groups Double blind Italy
DAVIT II , 1990 n=878/897 follow-up: 16 months	verapamil 120mgx3 for 18 months versus placebo	patients <76years with diagnosis of acute MI	Parallel groups Double blind Danish
Danish study , 1984 n=1729/1718 follow-up: 12 months	verapamil 0.1mg/kg IV plus 3x120mg orally versus placebo	patients under 75 years of age admitted to the CCU with a suspicion of acute myocardial infarction	Parallel groups double blind Danish

CT

References

Machecourt, 1986:

Muller, 1984:

Muller JE, Morrison J, Stone PH, Rude RE, Rosner B, Roberts R, Pearle DL, Turi ZG, Schneider JF, Serfas DH Nifedipine therapy for patients with threatened and acute myocardial infarction: a randomized, double-blind, placebo-controlled comparison *Circulation* 1984;69:740-7 [6365350]

Gordon, 1984:

Held PH, Yusuf S, Furberg CD Calcium channel blockers in acute myocardial infarction and unstable angina: an overview *BMJ* 1989;299:1187-92 [2513047]

MI study group, 1979:

Acta cardiol 1979;34(suppl24):7-46

Eichler, 1985:

Eichler HG, Mabin TA, Commerford PJ, Lloyd EA, Beck W, Opie LH Tiapamil, a new calcium antagonist: hemodynamic effects in patients with acute myocardial infarction. *Circulation* 1985;71:779-86 [3882269]

Bussman, 1984:

Bussmann WD, Seher W, Gruengras M Reduction of creatine kinase and creatine kinase-MB indexes of infarct size by intravenous verapamil *Am J Cardiol* 1984;54:1224-30 [6391131]

Gibson, 1986:

Gibson RS, Boden WE, Theroux P, Strauss HD, Pratt CM, Gheorghiadu M, Capone RJ, Crawford MH, Schlant RC, Kleiger RE Diltiazem and reinfarction in patients with non-Q-wave myocardial infarction. Results of a double-blind, randomized, multicenter trial. *N Engl J Med* 1986;315:423-9 [3526151]

Zannad, 1988:

Zannad F, Amor M, Karcher G, Maurin P, Ethevenot G, Sebag C, Bertrand A, Pernot C, Gilgenkrantz JM Effect of diltiazem on myocardial infarct size estimated by enzyme release, serial thallium-201 single-photon emission computed tomography and radionuclide angiography. *Am J Cardiol* 1988;61:1172-7 [3287881]

MDPIT, 1988:

The effect of diltiazem on mortality and reinfarction after myocardial infarction. The Multicenter Diltiazem Postinfarction Trial Research Group *N Engl J Med* 1988;319:385-92 [2899840]

Bigger JT Jr, Coromilas J, Rolnitzky LM, Fleiss JL, Kleiger RE Effect of diltiazem on cardiac rate and rhythm after myocardial infarction. Multicenter Diltiazem Postinfarction Trial Investigators *Am J Cardiol* 1990;65:539-46 [2178379]

Moss AJ, Oakes D, Rubison M, McDermott M, Carleen E, Eberly S, Brown M Effects of diltiazem on long-term outcome after acute myocardial infarction in patients with and without a history of systemic hypertension. The Multicenter Diltiazem Postinfarction Trial Research Group *Am J Cardiol* 1991;68:429-33 [1872266]

Gibson RS, Hansen JF, Messerli F, Schechtman KB, Boden WE Long-term effects of diltiazem and verapamil on mortality and cardiac events in non-Q-wave acute myocardial infarction without pulmonary congestion: post hoc subset analysis of the multicenter diltiazem postinfarction trial and the second danish verapamil infarction trial studies *Am J Cardiol* 2000;86:275-9 [10922432]

SPRINT II, 1993:

Goldbourt U, Behar S, Reicher-Reiss H, Zion M, Mandelzweig L, Kaplinsky E Early administration of nifedipine in suspected acute myocardial infarction. The Secondary Prevention Reinfarction Israel Nifedipine Trial 2 Study *Arch Intern Med* 1993;153:345-53 [8285927]

Sirnes, 1984:

Sirnes PA, Overskeid K, Pedersen TR, Bathen J, Drivenes A, Frland GS, Kjekshus JK, Landmark K, Rokseth R, Sirnes KE Evolution of infarct size during the early use of nifedipine in patients with acute myocardial infarction: the Norwegian Nifedipine Multicenter Trial. *Circulation* 1984;70:638-44 [6383655]

Gottlieb, 1988:

Gottlieb SO, Becker LC, Weiss JL, Shapiro EP, Chandra NC, Flaherty JT, Gottlieb SH, Ouyang P, Mellits ED, Townsend SN Nifedipine in acute myocardial infarction: an assessment of left ventricular function, infarct size, and infarct expansion. A double blind, randomised, placebo controlled trial *Br Heart J* 1988;59:411-8 [3285878]

Eisenberg, 1985:

Eisenberg PR, Lee RG, Biello DR, Geltman EM, Jaffe AS Chest pain after nontransmural infarction: the absence of remediable coronary vasospasm. *Am Heart J* 1985;110:515-21 [3898794]

Bradagan, 1986:

Branagan JP, Walsh K, Kelly P, Collins WC, McCafferty D, Walsh MJ Effect of early treatment with nifedipine in suspected acute myocardial infarction. *Eur Heart J* 1986;7:859-65 [3792347]

Loogna, 1985:

Loogna E, Sylvn C, Groth T, Mogensen L Complexity of enzyme release during acute myocardial infarction in a controlled study with early nifedipine treatment. *Eur Heart J* 1985;6:114-9 [3891346]

TRENT (Wilcox), 1986:

Wilcox RG, Hampton JR, Banks DC, Birkhead JS, Brooksby IA, Burns-Cox CJ, Hayes MJ, Joy MD, Malcolm AD, Mather HG Trial of early nifedipine in acute myocardial infarction: the Trent study. *Br Med J (Clin Res Ed)* 1986;293:1204-8 [3096431]

Walker, 1988:

Walker LJ, MacKenzie G, Adgey AA Effect of nifedipine on enzymatically estimated infarct size in the early phase of acute myocardial infarction. *Br Heart J* 1988;59:403-10 [3285877]

Jaffe, 1987:

Jaffe AS, Biello DR, Sobel BE, Geltman EM Enhancement of metabolism of jeopardized myocardium by nifedipine *Int J Cardiol* 1987;15:77-89 [3553039]

Erbel, 1988:

Erbel R, Pop T, Meinertz T, Olshausen KV, Treese N, Henrichs KJ, Schuster CJ, Rupprecht HJ, Schlurmann W, Meyer J Combination of calcium channel blocker and thrombolytic therapy in acute myocardial infarction *Am Heart J* 1988;115:529-38 [3278574]

SPRINT I, 1988:

Secondary prevention reinfarction Israeli nifedipine trial (SPRINT). A randomized intervention trial of nifedipine in patients with acute myocardial infarction. The Israeli Sprint Study Group *Eur Heart J* 1988;9:354-64 [2898363]

Disegni E, Goldbourt U, Reicher-Reiss H, Kaplinsky E, Zion M, Boyko V, Behar S The predictive value of admission heart rate on mortality in patients with acute myocardial infarction. SPRINT Study Group. Secondary Prevention Reinfarction Israeli Nifedipine Trial *J Clin Epidemiol* 1995;48:1197-205 [7561981]

DAVIT I, 1984:

Verapamil in acute myocardial infarction. The Danish Study Group on Verapamil in Myocardial Infarction *Eur Heart J* 1984;5:516-28 [6383832]

Hansen JF Treatment with verapamil after an acute myocardial infarction. Review of the Danish studies on verapamil in myocardial infarction (DAVIT I and II) *Drugs* 1991;42 Suppl 2:43-53 [1718701]

Crea, 1985:

Crea F, Deanfield J, Crean P, Sharom M, Davies G, Maseri A Effects of verapamil in preventing early postinfarction angina and reinfarction *Am J Cardiol* 1985;55:900-4 [3920891]

CRIS, 1996:

Rengo F, Carbonin P, Pahor M, DeCaprio L, Bernabei R, Ferrara N, Carosella L, Acanfora D, Parlati S, Vitale D A controlled trial of verapamil in patients after acute myocardial infarction: results of the calcium antagonist reinfarction Italian study (CRIS) *Am J Cardiol* 1996;77:365-9 [8602564]

Ferrara N *Giorn geront* 1996;44:577-583

DAVIT II, 1990:

Effect of verapamil on mortality and major events after acute myocardial infarction (the Danish Verapamil Infarction Trial II-DAVIT II) *Am J Cardiol* 1990;66:779-85 [2220572]

Vaage-Nilsen M, Rasmussen V Effect of verapamil on heart rate variability after an acute myocardial infarction. Danish Verapamil Infarction Trial II *Cardiovasc Drugs Ther* 1998;12:285-90 [9784908]

Vaage-Nilsen M, Rasmussen V, Hansen JF, Hagerup L, Sorensen MB, Pedersen-Bjergaard O, Mellempgaard K, Hollander NH, Nielsen I, Sigurd BM Effect of verapamil on arrhythmias and heart rate during 16 months following an acute myocardial infarction. The Danish Study Group on Verapamil in Myocardial Infarction *Cardiovasc Drugs Ther* 1994;8:147-51 [8086325]

Danish study, 1984:

Verapamil in acute myocardial infarction. The Danish Study Group on Verapamil in Myocardial Infarction. *Eur Heart J* 1984;5:516-28 [6383832]

3 long term calcium blockers

Trial	Treatments	Patients	Trials design and methods
nifedipine vs			
Muller , 1984 n=93/88 follow-up: 14 days	-	-	Parallel groups double blind
Gordon , 1984 n=13/13 follow-up: 1 day	-	-	Parallel groups single blind
MI study group , 1979 n=904/888	-	-	Parallel groups open
diltiazem vs placebo			
Gibson , 1986 n=287/289 follow-up: 14 days	Diltiazem 90mgx4 versus placebo	-	Parallel groups double blind
Zannad , 1988 n=17/17 follow-up:	Diltiazem 10-20mg/h IV, 4x60mg orrally versus placebo	-	Parallel groups double blind
MDPIT , 1988 n=1234/1232 follow-up: 25 months (at least 12 months)	Diltiazem 60mgx4 versus placebo	Patient aged 25 to 75 years, admitted to coronary care units with a documented acute myocardial infarction	Parallel groups Double blind US, Canada
nifedipine vs placebo			
SPRINT II , 1993 n=682/676 follow-up: 6 months	Nifedipine 20mgx3 started within 3hr of hopsital admission for 6 months versus placebo	men and women with suspected acute MI	Parallel groups Double blind Israel
Sirnes , 1984 n=112/115 follow-up: 42 days	Nifedipine 5x10mg versus placebo	patients with suspected acute myocardial infarction	Parallel groups double blind
Gottlieb , 1988 n=64/68 follow-up: 42 days	Nifedipine 10 to 30mgx4 versus placebo	-	Parallel groups double blind
Eisenberg , 1985 n=25/25 follow-up: 12 days	Nifedipine 20mgx4 versus placebo	patients with nontransmural myocardial infarction	Parallel groups double blind
Bradagan , 1986 n=60/61 follow-up: 2 days	Nifedipine 10mgx4 versus placebo	patients with suspected acute myocardial infarction within 6hrs	Parallel groups double blind
Loogna , 1985 n=23/34 follow-up:	Nifedipine 4x10mg versus placebo	patients with acute myocardial infarction within 6 h from onset of symptoms	Parallel groups double blind

continued...

Trial	Treatments	Patients	Trials design and methods
TRENT (Wilcox) , 1986 n=2240/2251 follow-up: 28 days	Nifedipine 4x10mg versus placebo	patients with suspected myocardial infarction	Parallel groups double blind
Walker , 1988 n=217/217 follow-up:	Nifedipine 6x10mg for 48hrs versus placebo	patients with suspected myocardial infarction within six hours from the onset of chest pain	Parallel groups double blind
Jaffe , 1987 n=13/9 follow-up: 7 days	Nifedipine 6x20mg versus placebo	patients with acute myocardial infarction	Parallel groups single blind
Erbel , 1988 n=74/75 follow-up: hospital stay	Nifedipine 20mgx3 versus placebo	patients with acute myocardial infarction	Parallel groups double blind
SPRINT I , 1988 n=1130/1146 follow-up: 1 year	Nifedipine 10mgx3 versus placebo	patient surviving MI 7 and 21 days after admission	Parallel groups Double blind Israel
verapamil vs placebo			
DAVIT I , 1984 n=1751/1747 follow-up: 6 months	verapamil 120mgx3 versus placebo	-	Parallel groups Double blind Danish
Crea , 1985 n=8/9 follow-up: 10 days	verapamil 80mgx3 versus placebo	patients admitted to the coronary care unit with transmural acute myocardial infarction	Parallel groups single blind
CRIS , 1996 n=531/542 follow-up: 23.5 months	verapamil retard 360 mg daily versus placebo	patients admitted for acute myocardial infarction	Parallel groups Double blind Italy
DAVIT II , 1990 n=878/897 follow-up: 16 months	verapamil 120mgx3 for 18 months versus placebo	patients <76years with diagnosis of acute MI	Parallel groups Double blind Danish
Danish study , 1984 n=1729/1718 follow-up: 12 months	verapamil 0.1mg/kg IV plus 3x120mg orally versus placebo	patients under 75 years of age admitted to the CCU with a suspicion of acute myocardial infarction	Parallel groups double blind Danish

References

Muller, 1984:

Muller JE, Morrison J, Stone PH, Rude RE, Rosner B, Roberts R, Pearle DL, Turi ZG, Schneider JF, Serfas DH Nifedipine therapy for patients with threatened and acute myocardial infarction: a randomized, double-blind, placebo-controlled comparison *Circulation* 1984;69:740-7 [6365350]

Gordon, 1984:

Held PH, Yusuf S, Furberg CD Calcium channel blockers in acute myocardial infarction and unstable angina: an overview *BMJ* 1989;299:1187-92 [2513047]

MI study group, 1979:

Acta cardiologica 1979;34(suppl24):7-46

Gibson, 1986:

Gibson RS, Boden WE, Theroux P, Strauss HD, Pratt CM, Gheorghide M, Capone RJ, Crawford MH, Schlant RC, Kleiger RE Diltiazem and reinfarction in patients with non-Q-wave myocardial infarction. Results of a double-blind, randomized, multicenter trial. *N Engl J Med* 1986;315:423-9 [3526151]

Zannad, 1988:

Zannad F, Amor M, Karcher G, Maurin P, Ethevenot G, Sebag C, Bertrand A, Pernot C, Gilgenkrantz JM Effect of diltiazem on myocardial infarct size estimated by enzyme release, serial thallium-201 single-photon emission computed tomography and radionuclide angiography. *Am J Cardiol* 1988;61:1172-7 [3287881]

MDPIT, 1988:

The effect of diltiazem on mortality and reinfarction after myocardial infarction. The Multicenter Diltiazem Postinfarction Trial Research Group *N Engl J Med* 1988;319:385-92 [2899840]

Bigger JT Jr, Coromilas J, Rolnitzky LM, Fleiss JL, Kleiger RE Effect of diltiazem on cardiac rate and rhythm after myocardial infarction. Multicenter Diltiazem Postinfarction Trial Investigators *Am J Cardiol* 1990;65:539-46 [2178379]

Moss AJ, Oakes D, Rubison M, McDermott M, Carleen E, Eberly S, Brown M Effects of diltiazem on long-term outcome after acute myocardial infarction in patients with and without a history of systemic hypertension. The Multicenter Diltiazem Postinfarction Trial Research Group *Am J Cardiol* 1991;68:429-33 [1872266]

Gibson RS, Hansen JF, Messerli F, Schechtman KB, Boden WE Long-term effects of diltiazem and verapamil on mortality and cardiac events in non-Q-wave acute myocardial infarction without pulmonary congestion: post hoc subset analysis of the multicenter diltiazem postinfarction trial and the second danish verapamil infarction trial studies *Am J Cardiol* 2000;86:275-9 [10922432]

SPRINT II, 1993:

Goldbourt U, Behar S, Reicher-Reiss H, Zion M, Mandelzweig L, Kaplinsky E Early administration of nifedipine in suspected acute myocardial infarction. The Secondary Prevention Reinfarction Israel Nifedipine Trial 2 Study *Arch Intern Med* 1993;153:345-53 [8285927]

Sirnes, 1984:

Sirnes PA, Overskeid K, Pedersen TR, Bathen J, Drivenes A, Frland GS, Kjekshus JK, Landmark K, Rokseth R, Sirnes KE Evolution of infarct size during the early use of nifedipine in patients with acute myocardial infarction: the Norwegian Nifedipine Multicenter Trial. *Circulation* 1984;70:638-44 [6383655]

Gottlieb, 1988:

Gottlieb SO, Becker LC, Weiss JL, Shapiro EP, Chandra NC, Flaherty JT, Gottlieb SH, Ouyang P, Mellits ED, Townsend SN Nifedipine in acute myocardial infarction: an assessment of left ventricular function, infarct size, and infarct expansion. A double blind, randomised, placebo controlled trial *Br Heart J* 1988;59:411-8 [3285878]

Eisenberg, 1985:

Eisenberg PR, Lee RG, Biello DR, Geltman EM, Jaffe AS Chest pain after nontransmural infarction: the absence of remediable coronary vasospasm. *Am Heart J* 1985;110:515-21 [3898794]

Bradagan, 1986:

Branagan JP, Walsh K, Kelly P, Collins WC, McCafferty D, Walsh MJ Effect of early treatment with nifedipine in suspected acute myocardial infarction. *Eur Heart J* 1986;7:859-65 [3792347]

Loogna, 1985:

Loogna E, Sylvn C, Groth T, Mogensen L Complexity of enzyme release during acute myocardial infarction in a controlled study with early nifedipine treatment. *Eur Heart J* 1985;6:114-9 [3891346]

TRENT (Wilcox), 1986:

Wilcox RG, Hampton JR, Banks DC, Birkhead JS, Brooksby IA, Burns-Cox CJ, Hayes MJ, Joy MD, Malcolm AD, Mather HG Trial of early nifedipine in acute myocardial infarction: the Trent study. *Br Med J (Clin Res Ed)* 1986;293:1204-8 [3096431]

Walker, 1988:

Walker LJ, MacKenzie G, Adgey AA Effect of nifedipine on enzymatically estimated infarct size in the early phase of acute myocardial infarction. *Br Heart J* 1988;59:403-10 [3285877]

Jaffe, 1987:

Jaffe AS, Biello DR, Sobel BE, Geltman EM Enhancement of metabolism of jeopardized myocardium by nifedipine *Int J Cardiol* 1987;15:77-89 [3553039]

Erbel, 1988:

Erbel R, Pop T, Meinertz T, Olshausen KV, Treese N, Henrichs KJ, Schuster CJ, Rupprecht HJ, Schlurmann W, Meyer J Combination of calcium channel blocker and thrombolytic therapy in acute myocardial infarction *Am Heart J* 1988;115:529-38 [3278574]

SPRINT I, 1988:

Secondary prevention reinfarction Israeli nifedipine trial (SPRINT). A randomized intervention trial of nifedipine in patients with acute myocardial infarction. The Israeli Sprint Study Group *Eur Heart J* 1988;9:354-64 [2898363]

Disegni E, Goldbourt U, Reicher-Reiss H, Kaplinsky E, Zion M, Boyko V, Behar S The predictive value of admission heart rate on mortality in patients with acute myocardial infarction. SPRINT Study Group. Secondary Prevention Reinfarction Israeli Nifedipine Trial *J Clin Epidemiol* 1995;48:1197-205 [7561981]

DAVIT I, 1984:

Verapamil in acute myocardial infarction. The Danish Study Group on Verapamil in Myocardial Infarction *Eur Heart J* 1984;5:516-28 [6383832]

Hansen JF Treatment with verapamil after an acute myocardial infarction. Review of the Danish studies on verapamil in myocardial infarction (DAVIT I and II) *Drugs* 1991;42 Suppl 2:43-53 [1718701]

Crea, 1985:

Crea F, Deanfield J, Crean P, Sharom M, Davies G, Maseri A Effects of verapamil in preventing early postinfarction angina and reinfarction *Am J Cardiol* 1985;55:900-4 [3920891]

CRIS, 1996:

Rengo F, Carbonin P, Pahor M, DeCaprio L, Bernabei R, Ferrara N, Carosella L, Acanfora D, Parlati S, Vitale D A controlled trial of verapamil in patients after acute myocardial infarction: results of the calcium antagonist reinfarction Italian study (CRIS) *Am J Cardiol* 1996;77:365-9 [8602564]

Ferrara N *Giorn geront* 1996;44:577-583

DAVIT II, 1990:

Effect of verapamil on mortality and major events after acute myocardial infarction (the Danish Verapamil Infarction Trial II-DAVIT II) *Am J Cardiol* 1990;66:779-85 [2220572]

Vaage-Nilsen M, Rasmussen V Effect of verapamil on heart rate variability after an acute myocardial infarction. Danish Verapamil Infarction Trial II *Cardiovasc Drugs Ther* 1998;12:285-90 [9784908]

Vaage-Nilsen M, Rasmussen V, Hansen JF, Hagerup L, Sorensen MB, Pedersen-Bjergaard O, Mellempgaard K, Hollander NH, Nielsen I, Sigurd BM Effect of verapamil on arrhythmias and heart rate during 16 months following an acute myocardial infarction. The Danish Study Group on Verapamil in Myocardial Infarction *Cardiovasc Drugs Ther* 1994;8:147-51 [8086325]

Danish study, 1984:

Verapamil in acute myocardial infarction. The Danish Study Group on Verapamil in Myocardial Infarction. *Eur Heart J* 1984;5:516-28 [6383832]

4 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.