

Clinical trials of antithrombotics for thrombosis prevention in arthroscopy

TrialResults-center www.trialresultscenter.org

1 Low molecular weight heparin

Trial	Treatments	Patients	Trials design and methods
deltaparin vs control			
Michot , 2002 n=66/64 follow-up: 30 days	deltaparin 2500IU 60-120min before procedure, followed 6hrs after the end of the procedure by 2500IU (<70kg) or 5000 IU(>70kg) versus no treatment	patients requiring diagnostic or therapeutic arthroscopic knee surgery as outpatients; aged 18 to 80 years.	Parallel groups open Switzerland.
enoxaparin vs control			
Canata , 2003 n=18/18 follow-up: 6 days	enoxaparin sc daily (dose not specified) versus no treatment	ACL reconstruction for symptomatic ACL-deficient knees	Parallel groups Italy
nadroparin vs control			
KANT (7 days) , 2008 n=657/660 follow-up: 3 months	once-daily subcutaneous injection of LMWH (nadroparin, 3800 anti-Xa IU) for 7 days versus full-length graduated compression stocking for 7 days	patients undergoing knee arthroscopy	Parallel groups open (blinded assessment) Italy
Roth , 1995 n=61/61 follow-up: 4 days	0.3 ml sc fraxiparine 2 hours before the operation and self administered daily (except the first two doses) for 4 days after surgery versus no treatment	patients undergoing ambulatory arthroscopic	Parallel groups Germany
nadroparin 14d vs control			
KANT (14 days) , 2008 n=444/660 follow-up: 3 months	once-daily subcutaneous injection of LMWH (nadroparin, 3800 anti-Xa IU) for 14 days versus full-length graduated compression stocking for 7 days	patients undergoing knee arthroscopy	Parallel groups open (blinded assessment) Italy
reviparin vs control			
Wirth , 2001 n=117/122 follow-up: 7-10 days	reviparin 1,750 anti Xa IU Sc once daily for 7-10 days versus no treatment	elective knee arthroscopy	Parallel groups open (blind assesement) Germany
extended-duration thromboprophylaxis vs standard-duration thromboprophylaxis			

continued...

Trial	Treatments	Patients	Trials design and methods
Marlovits , 2007 n=87/88 follow-up: <=28 days	extended-duration thromboprophylaxis with enoxaparin for 20 days versus standard thromboprophylaxis with enoxaparin (3 to 8 days)	patients who had undergone arthroscopic surgery of the anterior cruciate ligament	Parallel groups double-blind Austria

References

Michot, 2002:

Michot M, Conen D, Holtz D, Erni D, Zumstein MD, Ruffin GB, Renner N Prevention of deep-vein thrombosis in ambulatory arthroscopic knee surgery: A randomized trial of prophylaxis with low-molecular weight heparin. *Arthroscopy* 2002;18:257-63 [[11877611](#)]

Canata, 2003:

Canata GL, Chiey A. Prevention of venous thromboembolism after ACL reconstruction: a prospective, randomized study.(International Society of Arthroscopy, Knee Surgery and OrthopaedicSports Medicine) 2003; Vol. Poster 71-2003.

KANT (7 days), 2008:

Camporese G, NTita K, Rossi F, Bernardi E, Verlato F, Salmistraro G, Cordova R, et al. Different thromboprophylaxis approaches in patients undergoing knee arthroscopy (KANT study): interim report of prospective randomized study-f *Journal of Thrombosis & Haemostasis*.

Camporese G, Bernardi E, Prandoni P, Noventa F, Verlato F, Simioni P, Ntita K, Salmistraro G, Frangos C, Rossi F, Cordova R, Franz F, Zucchetto P, Kontothanassis D, Andreozzi GM Low-molecular-weight heparin versus compression stockings for thromboprophylaxis after knee arthroscopy: a randomized trial. *Ann Intern Med* 2008;149:73-82 [[18626046](#)]

Roth, 1995:

Roth P. Prophylaxis of deep vein thrombosis in outpatients undergoing arthroscopic meniscus operation *Orthopdische Praxis* 1995;5:3458.

KANT (14 days), 2008:

Camporese G, Bernardi E, Prandoni P, Noventa F, Verlato F, Simioni P, Ntita K, Salmistraro G, Frangos C, Rossi F, Cordova R, Franz F, Zucchetto P, Kontothanassis D, Andreozzi GM Low-molecular-weight heparin versus compression stockings for thromboprophylaxis after knee arthroscopy: a randomized trial. *Ann Intern Med* 2008;149:73-82 [[18626046](#)]

Wirth, 2001:

Wirth T, Schneider B, Misselwitz F, Lomb M, Tyl H, Egbring R, Griss P Prevention of venous thromboembolism after knee arthroscopy with low-molecular weight heparin (reviparin): Results of a randomized controlled trial. *Arthroscopy* 2001;17:393-9 [[11288011](#)]

Marlovits, 2007:

Marlovits S, Striessnig G, Schuster R, Stocker R, Luxl M, Trattng S, Vcsei V Extended-duration thromboprophylaxis with enoxaparin after arthroscopic surgery of the anterior cruciate ligament: a prospective, randomized, placebo-controlled study. *Arthroscopy* 2007;23:696-702 [[17637403](#)] [10.1016/j.arthro.2007.02.001](#)

2 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.