

# Clinical trials of antiplatelets drug for DVT prophylaxis in orthopedic surgery

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 platelet aggregation inhibitors

Trial	Treatments	Patients	Trials design and methods
<b>Aspirin vs no treatment</b>			
Pasteyer , 1977 n=20/20 follow-up: 2 weeks	Aspirin 1000mg daily + Hep versus control (Hep alone)	Elective orthopaedic surgery	Parallel groups
Rocha , 1986 n=60/30 follow-up: 1 weeks	Aspirin 250mg or 1000mg daily versus control (combination of heparin plus dihydroergotamine)	total hip replacement	Parallel groups open
<b>aspirin + dipyridamol vs no treatment</b>			
Morris-B , 1977 n=32/32 follow-up:	Aspirin 900 mg daily + dipyridamole versus control	elderly patients with hip fractures	Parallel groups open
Lyon-I , 1975 n=20/20 follow-up: 2 weeks	Aspirin 1500 mg daily + Dipyridamole versus control	Elective orthopaedic surgery	
<b>dipyridamol vs no treatment</b>			
Morris-A , 1977 n=24/24 follow-up:	dipyridamole versus control	elderly patients with hip fractures	Parallel groups open
<b>Flurbiprofen vs no treatment</b>			
Mocris-C , 1977 n=20/20 follow-up:	Flurbiprofen versus control	elderly patients with hip fractures	Parallel groups open
<b>Hydroxychloroquine vs no treatment</b>			
Massachusetts-I , 1981 n=50/25 follow-up:	Aspirin 1200mg daily, Aspirin 1200mg daily + hydroxychloroquine 300 mg b.i.d. versus control	patients with upper femoral fractures	Parallel groups open
<b>Aspirin vs placebo</b>			
Zekert-I , 1974 n=138/140 follow-up:	Aspirin 1500mg daily versus placebo	patients undergoing surgery of hip-joint proximal fractures	Parallel groups double-blind
Powers , 1976 n=66/63	A1300 versus placebo	traumatic orthopaedic surgery	

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
Erfurt-B , 1979 n=44/44 follow-up:	A1500 versus placebo	traumatic orthopaedic surgery	double-blind
Stockholm-I , 1975 n=26/25 follow-up: 2 weeks	Aspirin 2000mg daily versus placebo	elective surgery of the hip	double blind
Harris-I , 1977 n=58/59 follow-up: 1 weeks	Aspirin 1200mg daily versus placebo	patients over 40 years of age, who had undergone total hip replacement	Parallel groups double-blind
McKenna-I , 1980 n=24/12 follow-up: 2 weeks	Aspirin 975mg or 3900mg daily versus placebo	total knee replacement	Parallel groups double-blind
Sautter , 1983 n=68/77 follow-up: 3 weeks	Aspirin 900mg daily + sulfapyrazone versus placebo	patient with total hip replacement	Parallel groups
McBride , 1983 n=21/22 follow-up: 1 weeks	A1800+Dipyridamole versus placebo	Elective orthopaedic surgery	
<b>aspirin + dipyridamol vs placebo</b>			
Encke-II , 1976 n=34/25 follow-up:	Aspirin 1500mg daily, Aspirin 990mg daily + dipyridamol versus placebo	patients with abdominal operations	Parallel groups double-blind
Hamburg , 1976 n=21/11 follow-up: 3 weeks	A+Dipyridamole,A1000 versus placebo	Elective orthopaedic surgery	
<b>Hydroxychloroquine vs placebo</b>			
Danish-A , 1976 n=48/50 follow-up:	Hydroxychloroquine sulphate versus placebo	patients with fractures of the hip, pelvis, or thoracolumbar spine	Parallel groups double-blind
Danish-B , 1976 n=27/28 follow-up:	Hydroxychloroquine sulphate versus placebo	patients with fractures of the hip, pelvis, or thoracolumbar spine	Parallel groups
Massachusetts-II , 1976 n=51/51 follow-up: 3 weeks	Hydroxychloroquine 600mg daily versus placebo	fractures or orthopaedic operations involving the skeleton between the knee and the pelvis	
Cooke , 1977 n=25/25 follow-up: 2 weeks	Hydroxychloroquine versus placebo	elective surgery on the hip	Parallel groups double-blind
Hume-A , 1977 n=20/20 follow-up: 2 weeks	Hydroxychloroquine versus placebo	total hip replacement	
Stockholm-II , 1981 n=18/17 follow-up: 2 weeks	Hydroxychloroquine versus placebo	total hip replacement	

continued...

Trial	Treatments	Patients	Trials design and methods
<b>RA233 vs placebo</b>			
65279;Wood , 1973 n=21/9 follow-up:	RA233; Aspirin 600mg daily +RA233 versus placebo	traumatic orthopaedic surgery	
<b>Ticlopidine vs placebo</b>			
McKenna-II , 1983 n=29/29 follow-up: 2 weeks	Ticlopidine versus placebo	Elective orthopaedic surgery	
Lyon-II , 3000 n=20/20 follow-up: 3 weeks	Ticlopidine versus placebo	Elective orthopaedic surgery	
Gardecki , 3000 n=48/46 follow-up: 2 weeks	Ticlopidine versus placebo	Elective orthopaedic surgery	

## References

### Pasteyer, 1977:

Flicoteaux H, Kher A, Jean N, Blery M, Judet T, Honnart F, et al. Comparison of low dose heparin and low dose herparin combined with aspirin in prevention of deep vein thrombosis after total hip replacement. *Pathol Biol (Paris)* 1977;25(suppl):55-8.

### Rocha, 1986:

Alfaro MJ, Pramo JA, Rocha E Prophylaxis of thromboembolic disease and platelet-related changes following total hip replacement: a comparative study of aspirin and heparin-dihydroergotamine. *Thromb Haemost* 1986;56:53-6 [3535158]

### Morris-B , 1977:

Morris GK, Mitchell JR Preventing venous thromboembolism in elderly patients with hip fractures: studies of low-dose heparin, dipyridamole, aspirin, and flurbiprofen. *Br Med J* 1977;1:535-7 [843794]

### Lyon-I, 1975:

Dechavanne M, Ville D, Viala JJ, Kher A, Faivre J, Pousset MB, Dejour H Controlled trial of platelet anti-aggregating agents and subcutaneous heparin in prevention of postoperative deep vein thrombosis in high risk patients. *Haemostasis* 1975;4:94-100 [1205340]

### Morris-A , 1977:

Morris GK, Mitchell JR Preventing venous thromboembolism in elderly patients with hip fractures: studies of low-dose heparin, dipyridamole, aspirin, and flurbiprofen. *Br Med J* 1977;1:535-7 [843794]

### Mocris-C , 1977:

Morris GK, Mitchell JR Preventing venous thromboembolism in elderly patients with hip fractures: studies of low-dose heparin, dipyridamole, aspirin, and flurbiprofen. *Br Med J* 1977;1:535-7 [843794]

### Massachusetts-I , 1981:

Snook GA, Chrisman OD, Wilson TC Thromboembolism after surgical treatment of hip fractures. *Clin Orthop Relat Res* 1981;:21-4 [7014058]

### Zekert-I , 1974:

Zekert F, Kohn P, Vormittag E, Poigenfrst J, Thien M [Prevention of thromboembolism using acetylsalicylic acid in the surgery of hip-joint proximal fractures] *Monatsschr Unfallheilkd Versich Versorg Verkehrsmed* 1974;77:97-110 [4277091]

### Powers , 1976:

Hansen EH, Jessing P, Lindewald H, Ostergaard P, Olesen T, Malver EI Hydroxychloroquine sulphate in prevention of deep venous thrombosis following fracture of the hip, pelvis, or thoracolumbar spine. *J Bone Joint Surg Am* 1976;58:1089-93 [[1002750](#)]

Powers PJ, Gent M, Jay RM, Julian DH, Turpie AG, Levine M, Hirsh J A randomized trial of less intense postoperative warfarin or aspirin therapy in the prevention of venous thromboembolism after surgery for fractured hip. *Arch Intern Med* 1989;149:771-4 [[2650646](#)]

#### **Erfurt-B , 1979:**

Hartung B, Schreiber U, Rdiger H [Study of the platelet aggregation inhibitor MICRISTIN as to its efficacy in the prevention of thromboembolism in the postoperative phase following surgical interventions] *Folia Haematol Int Mag Klin Morphol Blutforsch* 1979;106:810-27 [[94873](#)]

#### **Stockholm-I, 1975:**

Soreff J, Johnsson H, Diener L, Gransson L Acetylsalicylic acid in a trial to diminish thromboembolic complications after elective hip surgery. *Acta Orthop Scand* 1975;46:246-55 [[1096521](#)]

#### **Harris-I, 1977:**

Harris WH, Salzman EW, Athanasoulis CA, Waltman AC, DeSanctis RW Aspirin prophylaxis of venous thromboembolism after total hip replacement. *N Engl J Med* 1977;297:1246-9 [[335247](#)]

#### **McKenna-I, 1980:**

McKenna R, Galante J, Bachmann F, Wallace DL, Kaushal PS, Meredith P Prevention of venous thromboembolism after total knee replacement by high-dose aspirin or intermittent calf and thigh compression. *Br Med J* 1980;280:514-7 [[6989432](#)]

#### **Sautter, 1983:**

Sautter RD, Koch EL, Myers WO, Ray JR 3rd, Mazza JJ, Larson DE, Chen HM, Milbauer JP, Treuhaft PS, Plotka ED Aspirin-sulfinpyrazone in prophylaxis of deep venous thrombosis in total hip replacement. *JAMA* 1983;250:2649-54 [[6355542](#)]

#### **McBride, 1983:**

McBride JA, Turpie AG, Kraus V, Hiltz C. Failure of aspirin and dipyridamole to influence the incidence of leg scan detected venous thrombosis after elective hip surgery *Thrombosis et Diathesis Haemorrhagica* 1975;34:abstract 204.

#### **Encke-II , 1976:**

Encke A, Stock C, Dumke HO [Double-blind study for the prevention of postoperative thrombosis] *Chirurg* 1976;47:670-3 [[1001131](#)]

#### **Hamburg, 1976:**

Boehringer Ingelheim DVT nach Hirntumoroperationen Boehringer Ingelheim, 1976. (Internal report.)

#### **Danish-A , 1976:**

Hansen EH, Jessing P, Lindewald H, Ostergaard P, Olesen T, Malver EI Hydroxychloroquine sulphate in prevention of deep venous thrombosis following fracture of the hip, pelvis, or thoracolumbar spine. *J Bone Joint Surg Am* 1976;58:1089-93 [[1002750](#)]

#### **Danish-B , 1976:**

Hansen EH, Jessing P, Lindewald H, Ostergaard P, Olesen T, Malver EI Hydroxychloroquine sulphate in prevention of deep venous thrombosis following fracture of the hip, pelvis, or thoracolumbar spine. *J Bone Joint Surg Am* 1976;58:1089-93 [[1002750](#)]

#### **Massachusetts-II, 1976:**

Chrisman OD, Snook GA, Wilson TC, Short JY Prevention of venous thromboembolism by administration of hydroxychloroquine. A preliminary report. *J Bone Joint Surg Am* 1976;58:918-20 [[789380](#)]

#### **Cooke, 1977:**

Cooke ED, Dawson MH, Ibbotson RM, Bowcock SA, Ainsworth ME, Pilcher MF Failure of orally administered hydroxychloroquine sulphate to prevent venous thromboembolism following elective hip operations. *J Bone Joint Surg Am* 1977;59:496-500 [[325009](#)]

#### **Hume-A, 1977:**

Hume M, Bierbaum B, Kuriakose TX, Surprenant, J Prevention of postoperative thrombosis by aspirin. *Am J Surg* 1977;133:420-2 [[322520](#)]

Hume M, Donaldson WR, Surprenant J Sex, aspirin, and venous thrombosis. *Orthop Clin North Am* 1978;9:761-7 [[358040](#)]

**Stockholm-II, 1981:**

Johansson E, Forsberg K, Johnsson H Clinical and experimental evaluation of the thromboprophylactic effect of hydroxychloroquine sulfate after total hip replacement. *Haemostasis* 1981;10:89-96 [7007179]

**65279;Wood , 1973:**

Wood EH, Prentice CR, McGrouther DA, Sinclair J, McNicol GP Trial of aspirin and RA 233 in prevention of post-operative deep vein thrombosis. *Thromb Diath Haemorrh* 1973;30:18-24 [4788747]

**McKenna-II, 1983:**

McKenna R, Galante J, Molony B, Kamm B. Failure of ticlopidine hydrochloride to prevent DVT in orthopedic patients *Blood* 1983;62(suppl 1):304A

**Lyon-II, 3000:****Gardecki, 3000:**

## 2 unfractionated heparin

Trial	Treatments	Patients	Trials design and methods
<b>UFH vs no treatment</b>			
<b>Bergqvist , 1979</b> n=84/77	UFH twice daily for 5 days versus no treatment	elective orthopedic surgery	
<b>Bergqvist , 1979</b> n=32/23	UFH twice daily for 5 days versus no treatment	traumatic orthopedic surgery	
<b>Dechavanne , 1974</b> n=29/29	UFH 3 times daily for 10 days versus no treatment	elective orthopedic surgery	
<b>Dechavanne , 1975</b> n=20/21	UFH 3 times daily for 10 days versus no treatment	elective orthopedic surgery	
<b>Galasko , 1976</b> n=50/50	UFH twice daily (duration unknown) versus no treatment	traumatic orthopedic surgery	
<b>Gallus , 1973</b> n=23/23	UFH 3 times daily for A versus no treatment	traumatic orthopedic surgery	
<b>Mannucci , 1976</b> n=23/24	UFH 3 times daily for 7 days versus no treatment	elective orthopedic surgery	
<b>Morris , 1974</b> n=36/36	UFH twice daily for 10 days versus no treatment	elective orthopedic surgery	
<b>Morris , 1977</b> n=24/24	UFH 3 times daily for 10 days versus no treatment	traumatic orthopedic surgery	

continued...

Trial	Treatments	Patients	Trials design and methods
VTCSG , 1975 n=34/30	UFH twice daily for 10 days versus no treatment	elective orthopedic surgery	
Welin-Berger , 1982 n=20/20	UFH twice daily for 7 days versus no treatment	elective orthopedic surgery	
<b>UFH vs placebo</b>			
Abraham-Inpijn , 1975 n=12/13	UFH twice daily for 8 days versus placebo	elective orthopedic surgery	
Hampson , 1974 n=48/52	UFH 3 times daily for 7- 10 days versus placebo	elective orthopedic surgery	
Lahnborg , 1980 n=70/69	UFH twice daily for 10 days versus placebo	traumatic orthopedic surgery	
Moskovitz , 1978 n=29/23	UFH 3 times daily for 7 days versus placebo	traumatic orthopedic surgery	
Moskovitz , 1978 n=35/32	UFH 3 times daily for 7 days versus placebo	elective orthopedic surgery	
Svend-Hansen , 1981 n=65/65	UFH 3 times daily for 14 days versus placebo	traumatic orthopedic surgery	
Williams , 1978 n=106/106	UFH twice daily for 14 days versus placebo	elective orthopedic surgery	
Xabregas , 1977 n=27/26	UFH twice daily for 14 days versus placebo	traumatic orthopedic surgery	

## References

### Bergqvist , 1979:

Bergqvist D, Efsing HO, Hallbk T, Hedlund T Thromboembolism after elective and post-traumatic hip surgery—a controlled prophylactic trial with dextran 70 and low-dose heparin. Acta Chir Scand 1979;145:213-8 [386676]

### Bergqvist, 1979:

Bergqvist D, Efsing HO, Hallbk T, Hedlund T Thromboembolism after elective and post-traumatic hip surgery—a controlled prophylactic trial with dextran 70 and low-dose heparin. Acta Chir Scand 1979;145:213-8 [386676]

### Dechavanne, 1974:

Dechavanne M, Saudin F, Viala JJ, Kher A, Bertrix L, de Mourgues G [Prevention of venous thrombosis. Success of high doses of heparin during total hip replacement for osteoarthritis] Nouv Presse Med 1974;3:1317-9 [4843085]

### Dechavanne, 1975:

Dechavanne M, Ville D, Viala JJ, Kher A, Faivre J, Pousset MB, Dejour H Controlled trial of platelet anti-aggregating agents and subcutaneous heparin in prevention of postoperative deep vein thrombosis in high risk patients. *Haemostasis* 1975;4:94-100 [[1205340](#)]

**Galasko , 1976:**

Galasko CS, Edwards DH, Fearn CB, Barber HM The value of low dosage heparin for the prophylaxis of thromboembolism in patients with transcervical and intertrochanteric femoral fractures. *Acta Orthop Scand* 1976;47:276-82 [[782146](#)]

**Gallus , 1973:**

Gallus AS, Hirsh J, Tuttle RJ, Trebilcock R, O'Brien SE, Carroll JJ, Minden JH, Hudecki SM Small subcutaneous doses of heparin in prevention of venous thrombosis. *N Engl J Med* 1973;288:545-51 [[4568221](#)]

**Mannucci , 1976:**

Mannucci PM, Citterio LE, Panajotopoulos N Low-dose heparin and deep-vein thrombosis after total hip replacement. *Thromb Haemost* 1976;36:157-64 [[1036806](#)]

**Morris, 1974:**

Morris GK, Henry AP, Preston BJ Prevention of deep-vein thrombosis by low-dose heparin in patients undergoing total hip replacement. *Lancet* 1974;2:797-800 [[4138248](#)]

**Morris , 1977:**

Morris GK, Mitchell JR Preventing venous thromboembolism in elderly patients with hip fractures: studies of low-dose heparin, dipyridamole, aspirin, and flurbiprofen. *Br Med J* 1977;1:535-7 [[843794](#)]

**VTCSG, 1975:**

Small doses of subcutaneous sodium heparin in the prevention of deep vein thrombosis after elective hip operations. *Br J Surg* 1975;62:348-50 [[1139128](#)]

**Welin-Berger, 1982:**

Welin-Berger T, Bygdeman S, Mebius C Deep vein thrombosis following hip surgery. Relation to activated factor X inhibitor activity: effect of heparin and dextran. *Acta Orthop Scand* 1982;53:937-45 [[6184938](#)]

**Abraham-Inpijn , 1975:**

Abraham-Inpijn L, Vreeken J Effect of low-dose heparin on incidence of postoperative thrombosis in orthopaedic patients. *Arch Chir Neerl* 1975;27:63-8 [[1098580](#)]

**Hampson, 1974:**

Hampson WG, Harris FC, Lucas HK, Roberts PH, McCall IW, Jackson PC, Powell NL, Staddon GE Failure of low-dose heparin to prevent deep-vein thrombosis after hip-replacement arthroplasty. *Lancet* 1974;2:795-7 [[4138905](#)]

**Lahnborg, 1980:**

Moreno-Gonzalez E, Sanmartin JH, Pascual MH, Moreno-Azcoita M, Selas PR Total esophagectomy by right anterior thoracotomy and immediate esophageal reconstruction for carcinoma of the esophagus. *Acta Chir Scand* 1980;146:19-23 [[7376779](#)]

Lahnborg G Effect of low-dose heparin and dihydroergotamine on frequency of postoperative deep-vein thrombosis in patients undergoing post-traumatic hip surgery. *Acta Chir Scand* 1980;146:319-22 [[7468062](#)]

**Moskovitz, 1978:**

Moskovitz PA, Ellenberg SS, Feffer HL, Kenmore PI, Neviasser RJ, Rubin BE, Varma VM Low-dose heparin for prevention of venous thromboembolism in total hip arthroplasty and surgical repair of hip fractures. *J Bone Joint Surg Am* 1978;60:1065-70 [[363722](#)]

**Moskovitz, 1978:**

Moskovitz PA, Ellenberg SS, Feffer HL, Kenmore PI, Neviasser RJ, Rubin BE, Varma VM Low-dose heparin for prevention of venous thromboembolism in total hip arthroplasty and surgical repair of hip fractures. *J Bone Joint Surg Am* 1978;60:1065-70 [[363722](#)]

**Svend-Hansen, 1981:**

Svend-Hansen H, Bremerskov V, Gtrik J, Ostri P Low-dose heparin in proximal femoral fractures. Failure to prevent deep-vein thrombosis. *Acta Orthop Scand* 1981;52:77-80 [[7010893](#)]

**Williams , 1978:**

Williams JW, Eikman EA, Greenberg SH, Hewitt JC, Lopez-Cuenca E, Jones GP, Madden JA Failure of low dose heparin to prevent pulmonary embolism after hip surgery or above the knee amputation. *Ann Surg* 1978;188:468-74 [697431]

**Xabregas , 1977:**

Xabregas A, Gray L, Ham JM Heparin prophylaxis of deep vein thrombosis in patients with a fractured neck of the femur. *Med J Aust* 1978;1:620-2 [355810]

Lawrence JC, Xabregas A, Gray L, Ham JM Seasonal variation in the incidence of deep vein thrombosis. *Br J Surg* 1977;64:777-80 [338089]

### 3 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.