

# Clinical trials of LMWH for thrombosis prevention in abdominal surgery

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 low molecular weight heparin

Trial	Treatments	Patients	Trials design and methods
<b>certoparin vs unfractionated heparin</b>			
Schmitz-Huebner , 1984 n=84/42 follow-up: 1 month	Certoparin (dose 1 and dose 2) b.i.d. versus UFH 10 000 units	Abdominal surgery	Blind
Sasahara , 1986 n=137/132 follow-up: 7 days	Certoparin 3000 + DHE versus UFH 10 000 units +DHE	Abdominal surgery	Blind
Voigt , 1986 n=103/97 follow-up: 10 days	Certoparin 3000 + DHE versus UFH 10 000 units	Abdominal surgery	Blind
Welzel , 1988 n=98/103 follow-up: 7 days	Certoparin 2500 + DHE versus UFH 10 000 units+DHE	Abdominal surgery	Open
Kakkar , 1989 n=88/91 follow-up:	Certoparin 3000 + DHE versus UFH 10 000 units+DHE	Abdominal surgery	Blind
Adolf , 1989 n=205/205 follow-up: 1 month	Certoparin 3000 versus UFH 15 000 units	Abdominal surgery	Blind
Baumgartner , 1989 n=99/102 follow-up: 10 days	Certoparin 3000 + DHE versus UFH 5 000 units+DHE	Abdominal surgery	Blind
Hoffmann and Largiade , 1990 n=464/452 follow-up:	Certoparin 3000 + DHE versus UFH 10 000 units	Abdominal surgery	NA
Kopenhagen , 1990 n=51/53	Certoparin 3000 anti Xa units versus UFH 15 000 units	Abdominal surgery	Blind
Schielke , 1991 n=47/51	Certoparin 3000 anti Xa units + DHE versus UFH 10 000 units + DHE	Abdominal surgery	Open
Kopenhagen , 1992 n=336/337	Certoparin 3000 anti Xa units versus UFH 15 000 units	Abdominal surgery	Blind
<b>dalteparin vs unfractionated heparin</b>			

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
Bergqvist , 1986 n=215/217 follow-up: 1 month	Dalteparin 5000 versus UFH 10 000 units	Abdominal surgery	Blind
Onarheim , 1986 n=25/27 follow-up: 1 month	Dalteparin 5000 versus UFH 10 000 units	Abdominal surgery	Blind
Koller , 1986 n=23/20 follow-up: 30 days	Dalteparin 7500 versus UFH 10 000 units	Abdominal surgery	Blind
Koller , 1986 n=75/75 follow-up: 30 days	Dalteparin 2500 versus UFH 10 000 units	Abdominal surgery	Blind
Fricker , 1988 n=40/40 follow-up: 1-2 months	Dalteparin 5000 versus UFH 15 000 units	Abdominopelvic surgery	Open
Bergqvist , 1988 n=505/497 follow-up: 1 month	Dalteparin 5000 versus UFH 10 000 units	Abdominal surgery	Blind
Caen , 1988 n=195/190 follow-up: 1 month	Dalteparin 2500 versus UFH 10 000 units	Abdominal surgery	Blind
Hartl , 1990 n=126/124 follow-up: >7 days	Dalteparin 2500 versus UFH 10 000 units	Abdominal surgery	Blind
Kakkar , 1993 n=1894/1915	Dalteparin 2500 anti Xa units versus UFH 10 000 units	Abdominal surgery	Blind
<b>enoxaparin vs unfractionated heparin</b>			
McLeod (Canadian) , 1995 n=674/675	Enoxaparin 4000 anti Xa units versus UFH 15 000 units	Colorectal surgery	Blind
Gonzalez , 1996 n=84/82	Bemiparin 2500 anti Xa units versus UFH 10 000 units	Abdominal surgery	Blind
<b>nadroparin vs unfractionated heparin</b>			
EFS , 1988 n=968/941 follow-up: 1 month	Nadroparin 2850 versus UFH 15 000 units	Abdominal surgery	Open
<b>parnaparin vs unfractionated heparin</b>			
Catania , 1988 n=88/85 follow-up:	Parnaparin 3200 versus UFH 15 000 units	Abdominal surgery	Open
Salcuni , 1988 n=73/68 follow-up:	Parnaparin 3200 versus UFH 15 000 units	Abdominal surgery	Open

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>Verardi , 1989</b> n=44/44 follow-up:	Parnaparin 6400 versus UFH 10 000 units	Abdominal/urological surgery	NA
<b>Garcea , 1992</b> n=45/45	Parnaparin 3200 anti Xa units versus UFH 15 000 units	Abdominal surgery	Open

## References

### Schmitz-Huebner, 1984:

Schmitz-Huebner U, Bnte H, Freise G, Reers B, Rskemeyer C, Scherer R, Schulte H, van de Loo J Clinical efficacy of low molecular weight heparin in postoperative thrombosis prophylaxis. *Klin Wochenschr* 1984;62:349-53 [[6374278](#)]

### Sasahara, 1986:

Sasahara AA, Koppenhagen K, Hring R, Welzel D, Wolf H Low molecular weight heparin plus dihydroergotamine for prophylaxis of postoperative deep vein thrombosis. *Br J Surg* 1986;73:697-700 [[3530367](#)]

### Voigt, 1986:

Voigt J, Hamelmann H, Hedderich J, Seifert J, Buchhammer T, Khler A [Effectiveness and side effects of low-molecular weight heparin-dihydroergotamine in preventing thromboembolism in abdominal surgery] *Zentralbl Chir* 1986;111:1269-305 [[3544605](#)]

### Welzel, 1988:

Welzel D, Wolf H, Koppenhagen K Antithrombotic defense during the postoperative period. Clinical documentation of low molecular weight heparin. *Arzneimittelforschung* 1988;38:120-3 [[2835055](#)]

### Kakkar, 1989:

Kakkar VV, Stringer MD, Hedges AR, Parker CJ, Welzel D, Ward VP, Sanderson RM, Cooper D, Kakkar S Fixed combinations of low-molecular weight or unfractionated heparin plus dihydroergotamine in the prevention of postoperative deep vein thrombosis. *Am J Surg* 1989;157:413-8 [[2539025](#)]

### Adolf, 1989:

Adolf J, Knee H, Roder JD, van de Fliedrt E, Siewert JR [Thromboembolism prophylaxis with low molecular weight heparin in abdominal surgery] *Dtsch Med Wochenschr* 1989;114:48-53 [[2535983](#)]

### Baumgartner, 1989:

Baumgartner A, Jacot N, Moser G, Krhenbhl B Prevention of postoperative deep vein thrombosis by one daily injection of low molecular weight heparin and dihydroergotamine. *Vasa* 1989;18:152-6 [[2545054](#)]

### Hoffmann and Largiade, 1990:

### Koppenhagen, 1990:

### Schielke, 1991:

### Koppenhagen, 1992:

### Bergqvist, 1986:

Bergqvist D, Burmark US, Frisell J, Hallbk T, Lindblad B, Risberg B, Trngren S, Wallin G Low molecular weight heparin once daily compared with conventional low-dose heparin twice daily. A prospective double-blind multicentre trial on prevention of postoperative thrombosis. *Br J Surg* 1986;73:204-8 [[3512031](#)]

### Onarheim, 1986:

Onarheim H, Lund T, Heimdal A, Arnesj B A low molecular weight heparin (KABI 2165) for prophylaxis of postoperative deep venous thrombosis. *Acta Chir Scand* 1986;152:593-6 [[3544625](#)]

### Koller, 1986:

Koller M, Schoch U, Buchmann P, Largiadri F, von Felten A, Frick PG Low molecular weight heparin (KABI 2165) as thromboprophylaxis in elective visceral surgery. A randomized, double-blind study versus unfractionated heparin. *Thromb Haemost* 1986;56:243-6 [[3551180](#)]

**Koller, 1986:**

Koller M, Schoch U, Buchmann P, Largiadri F, von Felten A, Frick PG Low molecular weight heparin (KABI 2165) as thromboprophylaxis in elective visceral surgery. A randomized, double-blind study versus unfractionated heparin. *Thromb Haemost* 1986;56:243-6 [[3551180](#)]

**Fricker, 1988:**

Fricker JP, Vergnes Y, Schach R, Heitz A, Eber M, Grunebaum L, Wiesel ML, Kher A, Barbier P, Cazenave JP Low dose heparin versus low molecular weight heparin (Kabi 2165, Fragmin) in the prophylaxis of thromboembolic complications of abdominal oncological surgery. *Eur J Clin Invest* 1988;18:561-7 [[2852111](#)]

**Bergqvist, 1988:**

Bergqvist D, Mtzsch T, Burmark US, Frisell J, Guilbaud O, Hallbk T, Horn A, Lindhagen A, Ljungnr H, Ljungstrm KG Low molecular weight heparin given the evening before surgery compared with conventional low-dose heparin in prevention of thrombosis. *Br J Surg* 1988;75:888-91 [[2846113](#)]

**Caen, 1988:**

Caen JP A randomized double-blind study between a low molecular weight heparin Kabi 2165 and standard heparin in the prevention of deep vein thrombosis in general surgery. A French multicenter trial. *Thromb Haemost* 1988;59:216-20 [[2838923](#)]

**Hartl, 1990:**

Hartl P, Brcke P, Dienstl E, Vinazzer H Prophylaxis of thromboembolism in general surgery: comparison between standard heparin and Fragmin. *Thromb Res* 1990;57:577-84 [[2158151](#)]

**Kakkar, 1993:**

**McLeod (Canadian), 1995:**

**Gonzalez, 1996:**

**EFS, 1988:**

Comparison of a low molecular weight heparin and unfractionated heparin for the prevention of deep vein thrombosis in patients undergoing abdominal surgery. The European Fraxiparin Study (EFS) Group. *Br J Surg* 1988;75:1058-63 [[2905187](#)]

**Catania, 1988:**

Catania G, Salantri G Prevention of postoperative deep vein thrombosis by two different heparin types. *Int J Clin Pharmacol Ther Toxicol* 1988;26:304-9 [[2842266](#)]

**Salcuni, 1988:**

**Verardi, 1989:**

Verardi S, Cortese F, Baroni B, Boffo V, Casciani CU [Role of low molecular weight heparin in the prevention of postoperative deep venous thrombosis. Our experience in 88 cases] *G Chir* 1989;10:674-8 [[2562010](#)]

**Garcea, 1992:**

## 2 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.