

Clinical trials of Drug eluting stent for coronary artery disease in diabetic patients

TrialResults-center www.trialresultscenter.org

1 drug-eluting stents

| Trial | Treatments | Patients | Trials design and methods |
|---|---|---|--|
| paclitaxel eluting stent vs bare-metal stent | | | |
| TAXUS II (diabetics) , 2003 <i>unpublished</i> n=37/41 follow-up: 12 months | TAXUS versus NIR stent | Diabetic patients with stable or unstable AP, silent ischaemia; single de novo target lesion with estimated stenosis >50% and <99% , | Parallel groups double-blind Europe |
| TAXUS IV (diabetics) , 2005 [NCT00292474] n=155/163 follow-up: 9 months | TAXUS versus EXPRESS | Diabetic patients with stable or unstable AP, provokable ischaemia with a single, previously untreated coronary-artery stenosis (vessel diameter, 2.5 to 3.75 mm; lesion length, 10 to 28 mm) | Parallel groups double-blind United States |
| TAXUS V (diabetics) , 2005 n=178/171 follow-up: 9 months | TAXUS versus BMS | Diabetic patients with stable or unstable AP, silent ischaemia with complex or previously unstudied lesions (requiring 2.25-mm, 4.0-mm, and/or multiple stents) | Parallel groups double-blind United States |
| TAXUS VI (diabetics) , 2005 [NCT00297804] n=39/50 follow-up: 9 months | TAXUS versus Express2 stent | Diabetic patients with stable or unstable AP, silent ischaemia with long, complex coronary artery lesions | Parallel groups double-blind Europe |
| sirolimus eluting stent vs bare-metal stent | | | |
| DECODE , 2005 <i>unpublished</i> [NCT00489164] n=54/29 follow-up: 1 year | CYPHER (Up to 3 stents per patient were allowed) versus Bx VELOCITY (Up to 3 stents per patient were allowed) | Stable or unstable angina in diabetic patients with with up to 2 de novo lesions in up to 2 native coronary vessels | Parallel groups open US, Asia/Pacific |
| DESSERT , 2008 n=75/75 follow-up: 12 months | Cypher and Cypher Select versus Sonic (Cordis) | de novo lesions of diabetic patients treated with insulin and/or oral antidiabetics for >3 months | Parallel groups single-blind Italy |
| DIABETES , 2005 n=80/80 follow-up: 9 months | Cypher versus Bx Velocity/Sonic | de novo lesions in native coronary arteries in 1, 2, or 3 native vessels with symptoms or objective evidence of ischemia; vessel size smaller than 4.0 mm | Parallel groups open Spanish |
| Ravel (diabetics) , 2004 n=19/25 follow-up: 6 months | coated Bx velocity versus Bx VELOCITY | sub groups of diabetic patients with de novo native coronary artery lesions 2.5 to 3.5 mm in diameter by visual assessment that could be covered by an 18-mm stent | Parallel groups NA Europe |

continued...

| Trial | Treatments | Patients | Trials design and methods |
|--|--|--|--|
| SES-SMART (diabetics) , 2005 n=29/45 follow-up: 8 months | Cypher versus Bx Sonic | Diabetic patients with de novo target lesion <=2.75 mm in diameter in a native coronary artery that could be completely covered by a single stent (maximum length 33 mm) | Parallel groups single-blind Italy |
| SIRIUS (diabetics) , 2003 n=131/148 follow-up: 12 months | SES versus BMS | sub group of diabetics patients of SIRIUS study | Parallel groups double-blind US |
| CoStar stent vs paclitaxel eluting stent | | | |
| COSTAR II diabetic (sub group) , 2008 n=271/271 follow-up: 8 months | CoStar stent (PES) versus Taxus stent (PES) | patients with de novo single- or multivessel coronary disease | Parallel groups open |
| paclitaxel eluting balloon vs paclitaxel eluting stent | | | |
| PEPCAD IV <i>ongoing</i> [NCT00462631] n=NA follow-up: | Paclitaxel-eluting PTCA-balloon dilation (SeQuent™ Please) followed by cobalt-chromium stent (Coroflex™ Blue) deployment versus Taxus Libert | patients with diabetes mellitus | open |
| sirolimus eluting stent vs paclitaxel eluting stent | | | |
| DES-DIABETES , 2008 n=200/200 follow-up: 9 months (1 year) | sirolimus-eluting stent versus paclitaxel-elutingstent | diabetic patients with angina pectoris and/or a positive stress test and a native coronary lesion | Factorial plan open Korea |
| ISAR-DIABETES , 2005 n=125/125 follow-up: 9 months | Taxus versus Cypher | Diabetic patients. AP or positive stress, no AMI with clinically significant angiographic stenosis in a native coronary vessel | Parallel groups open Germany |
| Kim , 2008 n=85/84 follow-up: 6 months | Cypher versus Taxus | Korean diabetic patients with high-grade de novo coronary lesions (stenosis of >70 percent of the luminal diameter) requiring <3 stents | Parallel groups open Korea |
| REALITY (diabetics) , 2006 <i>unpublished</i> n=187/192 follow-up: 12 months | SES versus PES | - | Parallel groups open worldwide |
| SIRTAX diabetics , 2005 [NCT00297661] n=108/93 follow-up: 12 months | Cypher versus Taxus | Sub groups of diabetics patients with either stable angina or an acute coronary syndrome | Parallel groups single-blind Switzerland |
| TAXi (diabetics) , 3000 <i>unpublished</i> n=33/36 follow-up: 12 months | SES versus PES | - | Parallel groups open Switzerland |

continued...

| Trial | Treatments | Patients | Trials design and methods |
|---|--|---|------------------------------------|
| Tomai , 2008 n=60/60 follow-up: 8 months | sirolimus-eluting stent versus paclitaxel-eluting stent | diabetic patient with multiple de novo coronary artery lesions | Cross over NA Italy |
| Lipsia-Yukon-DM <i>ongoing</i> [NCT00368953] n=NA follow-up: 9 months | Yukon Choice stent system versus Taxus Libert stent system | Patients With Diabetes Mellitus | |
| paclitaxel eluting stent vs sirolimus eluting stent | | | |
| ISAR-test (diabetics) , 2006 n=73/58 follow-up: 9 months | Taxus versus rapamycin stent | diabetics patients with de novo lesions in native coronary vessels, excluding the left main trunk | Parallel groups open germany |
| zotarolimus eluting stent vs sirolimus eluting stent | | | |
| DIABEDES IV <i>ongoing</i> [NCT00552994] n=NA follow-up: | Cypher select plus versus Xience V | diabetic patients | |

References

TAXUS II (diabetics), 2003:

Hermiller J. Diabetic results: Taxus II, IV and VI TCT [0]

TAXUS IV (diabetics), 2005:

Hermiller JB, Raizner A, Cannon L, Gurbel PA, Kutcher MA, Wong SC, Russell ME, Ellis SG, Mehran R, Stone GW Outcomes with the polymer-based paclitaxel-eluting TAXUS stent in patients with diabetes mellitus: the TAXUS-IV trial. J Am Coll Cardiol 2005;45:1172-9 [15837245]

TAXUS V (diabetics), 2005:

Ellis SG TAXUS V trial global results: expanding the randomizeddata 2005 American College of CardiologyAnnual Scientific Session

TAXUS VI (diabetics), 2005:

Dawkins KD, Grube E, Guagliumi G, Banning AP, Zmudka K, Colombo A, Thuesen L, Hauptman K, Marco J, Wijns W, Popma JJ, Koglin J, Russell ME Clinical efficacy of polymer-based paclitaxel-eluting stents in the treatment of complex, long coronary artery lesions from a multicenter, randomized trial: support for the use of drug-eluting stents in contemporary clinical practice. Circulation 2005;112:3306-13 [16286586]

DECODE, 2005:

Chan C, Zambahari R, Kaul U, Cohen SA, Buchbinder M. Outcomes in diabetic patients with multivessel disease and long lesions: results from the DECODE study Am J Cardiol 2005; 96 (suppl 7A): 31H

DESSERT, 2008:

Maresta A, Varani E, Balducelli M, Varbella F, Lettieri C, Uguccioni L, Sangiorgio P, Zoccai GB Comparison of effectiveness and safety of sirolimus-eluting stents versus bare-metal stents in patients with diabetes mellitus (from the Italian Multicenter Randomized DESSERT Study). Am J Cardiol 2008;101:1560-6 [18489933]

DIABETES, 2005:

Sabat M, Jimnez-Quevedo P, Angiolillo DJ, Gmez-Hospital JA, Alfonso F, Hernandez-Antoln R, Goicolea J, Bauelos C, Escaned J, Moreno R, Fernandez C, Fernandez-Aviles F, Macaya C Randomized comparison of sirolimus-eluting stent versus standard stent for percutaneous coronary revascularization in diabetic patients: the diabetes and sirolimus-eluting stent (DIABETES) trial. Circulation 2005;112:2175-83 [16203930]

Jimnez-Quevedo P, Sabat M, Angiolillo DJ, Alfonso F, Hernndez-Antoln R, SanMartn M, Gmez-Hospital JA, Bauelos C, Escaned J, Moreno R, Fernndez C, Fernndez-Avils F, Macaya C Long-term clinical benefit of sirolimus-eluting stent implantation in diabetic patients with de novo coronary stenoses: long-term results of the DIABETES trial. Eur Heart J 2007;28:1946-52 [17562666]

Ravel (diabetics), 2004:

SES-SMART (diabetics), 2005:

SIRIUS (diabetics), 2003:

COSTAR II diabetic (sub group), 2008:

PEPCAD IV, 0:

DES-DIABETES, 2008:

ISAR-DIABETES, 2005:

Kim, 2008:

REALITY (diabetics), 2006:

SIRTAX diabetics, 2005:

TAXi (diabetics), 3000:

Tomai, 2008:

Lipsia-Yukon-DM, 0:

ISAR-test (diabetics), 2006:

DIABEDES IV, 0:

2 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.