

# Clinical trials of Drug eluting stent for coronary artery disease in long or complex lesion

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 drug-eluting stents

Trial	Treatments	Patients	Trials design and methods
<b>paclitaxel eluting stent vs bare-metal stent</b>			
<b>TAXUS V (all patients) , 2005</b> [NCT00301522] n=577/579 follow-up: 9 months	TAXUS versus bare metal EXPRESS-2	Stable or unstable AP, silent ischaemia with single coronary artery stenosis including complex or previously unstudied lesions (requiring 2.25-mm, 4.0-mm, and/or multiple stents)	Parallel groups double-blind United States
<b>sirolimus eluting stent vs bare-metal stent</b>			
<b>PRISON II , 2006</b> [NCT00258596] n=100/100 follow-up: 6 months	Cypher versus BxVelocity	Chronic total occlusion, positive exercise stress test	Parallel groups single-blind Belgium
<b>SCANDSTENT , 2006</b> [NCT00151658] n=163/159 follow-up: 7 months	Cypher versus Sonic	Stable or unstable AP, recent AMI (non ST-elevation); with one or more de novo complex lesions in native coronary vessels (occluded, bifurcational, ostial or angulated)	Parallel groups open Denmark
<b>paclitaxel eluting stent vs CABG</b>			
<b>SYNTAX , 2009</b> [NCT00114972] n=903/897 follow-up: 1 year	paclitaxel (taxus Express SR) versus Coronary Artery Bypass Surgery (on- or off-pump bypass)	patients with previously untreated three-vessel or left main coronary artery disease (or both) (complex lesions)	Parallel groups open
<b>Genous stent vs paclitaxel eluting stent</b>			
<b>TRIAS-HR , 2008</b> <i>unpublished</i> [ISRCTN74297220] n=98/95 follow-up: 12 months	Genous stent (antibody-coated bare-metal stent) followed by one month of dual antiplatelet therapy versus Taxus or Cypher followed by at least six months of dual antiplatelet therapy	high-risk patients (long lesions, small vessels, chronic total occlusions, or any lesion in a diabetic patient)	Parallel groups single-blind
<b>sirolimus eluting stent vs paclitaxel eluting stent</b>			
<b>CORPAL , 2005</b> <i>unpublished</i> n=331/321 follow-up:	sirolimus versus paclitaxel	Documented myocardial ischaemia, no AMI	Parallel groups open Spain
<b>LONG DES II , 2006</b> n=250/250 follow-up: 9 months	SES versus PES	Long lesions. AP or positive stress, no AMI	Parallel groups single-blind Korea

## References

### **TAXUS V (all patients), 2005:**

Stone GW, Ellis SG, Cannon L, Mann JT, Greenberg JD, Spriggs D, O'Shaughnessy CD, DeMaio S, Hall P, Popma JJ, Koglin J, Russell ME Comparison of a polymer-based paclitaxel-eluting stent with a bare metal stent in patients with complex coronary artery disease: a randomized controlled trial. *JAMA* 2005;294:1215-23 [[16160130](#)]

### **PRISON II, 2006:**

Suttorp MJ, Laarman GJ, Rahel BM, Kelder JC, Bosschaert MA, Kiemeneij F, Ten Berg JM, Bal ET, Rensing BJ, Eefting FD, Mast EG Primary Stenting of Totally Occluded Native Coronary Arteries II (PRISON II): a randomized comparison of bare metal stent implantation with sirolimus-eluting stent implantation for the treatment of total coronary occlusions. *Circulation* 2006;114:921-8 [[16908768](#)]

### **SCANDSTENT, 2006:**

Kelbaek H, Thuesen L, Helqvist S, Klwgaard L, Jrgensen E, Aljabbari S, Saunamki K, Krusell LR, Jensen GV, Btker HE, Lassen JF, Andersen HR, Thayssen P, Galle A, van Weert A The Stenting Coronary Arteries in Non-stress/benestent Disease (SCANDSTENT) trial. *J Am Coll Cardiol* 2006;47:449-55 [[16412876](#)]

Kelbaek H, Klwgaard L, Helqvist S, Lassen JF, Krusell LR, Engstrm T, Btker HE, Jrgensen E, Saunamki K, Aljabbari S, Thayssen P, Galle A, Jensen GV, Thuesen L Long-term outcome in patients treated with sirolimus-eluting stents in complex coronary artery lesions: 3-year results of the SCANDSTENT (Stenting Coronary Arteries in Non-Stress/Benestent Disease) trial. *J Am Coll Cardiol* 2008 May 27;51:2011-6 [[18498953](#)]

### **SYNTAX, 2009:**

Lee TH, Hillis LD, Nabel EG CABG vs. stenting—clinical implications of the SYNTAX trial. *N Engl J Med* 2009 Feb 19;360:e10 [[19228613](#)] [10.1056/NEJMp0900462](#)

Serruys PW, Morice MC, Kappetein AP, Colombo A, Holmes DR, Mack MJ, Sthle E, Feldman TE, van den Brand M, Bass EJ, Van Dyck N, Leadley K, Dawkins KD, Mohr FW Percutaneous coronary intervention versus coronary-artery bypass grafting for severe coronary artery disease. *N Engl J Med* 2009 Mar 5;360:961-72 [[19228612](#)] [10.1056/NEJMoa0804626](#)

Banning AP, Westaby S, Morice MC, Kappetein AP, Mohr FW, Berti S, Glauber M, Kellett MA, Kramer RS, Leadley K, Dawkins KD, Serruys PW Diabetic and nondiabetic patients with left main and/or 3-vessel coronary artery disease: comparison of outcomes with cardiac surgery and paclitaxel-eluting stents. *J Am Coll Cardiol* 2010;55:1067-75 [[20079596](#)] [10.1016/j.jacc.2009.09.057](#)

### **TRIAS-HR, 2008:**

### **CORPAL, 2005:**

de Lezo J, Medina A, Pan M, et al. de Lezo J, Medina A, Pan M, et al. Drug-eluting stent for complex lesions: latest angiographic data from randomized rapamycin versus paclitaxel CORPAL study *J Am Coll Cardiol* 2005; 45: 75A.

### **LONG DES II, 2006:**

Kim YH, Park SW, Lee SW, Park DW, Yun SC, Lee CW, Hong MK, Kim HS, Ko JK, Park JH, Lee JH, Choi SW, Seong IW, Cho YH, Lee NH, Kim JH, Chun KJ, Park SJ Sirolimus-eluting stent versus paclitaxel-eluting stent for patients with long coronary artery disease. *Circulation* 2006;114:2148-53 [[17060388](#)]

Entry terms: liraglutide, liraglutide, victoza, nouveau, exenatide, exendin 4, exendin-4, Ex4 peptide, Byetta, AC 2993, AC 2993 LAR,

## 2 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.