

Clinical trials of fibrinolysis for acute myocardial infarction in all type of patients

TrialResults-center www.trialresultscenter.org

1 fibrinolysis

Trial	Treatments	Patients	Trials design and methods
APSAC vs control			
APSIM , 1989 n=112/119 follow-up: 3 weeks	APSAC 30 U over 5 min versus control (conventional heparin therapy, 5,000 IU in a bolus injection)	patients with a first acute myocardial infarction within 5 h after the onset of symptoms	Parallel groups open France
urokinase vs control			
USIM , 1991 n=1128/1073 follow-up: in hospital	urokinase bolus dose of 1 million U repeated after 60 minutes plus heparin versus control (heparin alone)	patients with acute myocardial infarction within 4 hours of the onset of pain	Parallel groups open Italy
APSAC vs placebo			
AIMS , 1988 n=624/634 follow-up: 1 y	APSAC 30U IV in 5 min versus Placebo	Hommes et femmes, <70 ans	Parallel groups double blind
German Multicenter Trial , 1988 n=162/151 follow-up: 28 jours	APSAC 30 units en IV en 5 min, puis hparine en IV (17 U/kg/h) 4 h aprs l'injection d'APSAC versus Hparine 5000 U en bolus en IV, puis 17 U/kg/h	Hommes et femmes, <70 ans	Parallel groups
streptokinase vs placebo			
EMERAS (7-12h) , 1993 n=2257/2277 follow-up:	intravenous streptokinase 1.5 MU versus placebo	patients presenting 7-12 h from symptom onset	Parallel groups double blind
EMERAS (all delay) , 1993 n=2257/2277 follow-up:	streptokinase 1.5 MU versus placebo	patients entering hospital up to 24 h after the onset of suspected acute myocardial infarction	Parallel groups double blind south america
GISSI I , 1986 n=5860/5852 follow-up: 1 y	Streptokinase 1.5 MU en perfusion IV en 1 heure versus usual care	patients within 12 h after the onset of symptoms and with no contraindications to SK	Parallel groups open
ISAM , 1986 n=859/882 follow-up: 21 days	1.5 million IU of streptokinase over 1h versus Placebo	patients within six hours after the onset of symptoms of myocardial infarction	Parallel groups double blind

continued...

Trial	Treatments	Patients	Trials design and methods
ISIS 2 pilot , 1987 n=NA follow-up:	streptokinase 1.5 MU versus placebo	patients with suspected acute myocardial infarction	Parallel groups double blind
ISIS-2 (SK) , 1988 n=8592/8595 follow-up: 15 mo	1-hour intravenous infusion of 1.5 MU of streptokinase versus Placebo	patients within 24h of the onset of suspected acute myocardial infarction	plan factoriel 2*2 double blind
Western Washington Intravenous Trial , 1988 [NCT00000507] n=191/177 follow-up: 1.4 y	Streptokinase en IV, 1.5 M UI en 60 min aprs injection de benadryl 50 mg en IV et hydrocortisone 100 mg en IV; hparine en IV 1000 UI/h 2h aprs la streptokinase puis warfarine pendant au moins 3 mois versus Traitement standard, avec ou sans anticoagulant (dcid par le mdecin)	Hommes et femmes, <ou = 75 ans	Parallel groups
t-PA vs placebo			
ASSET , 1988 n=2516/2495 follow-up: 6 months	rt-PA 100 mg versus Placebo	patient with suspected acute myocardial infarction	Parallel groups double blind
LATE , 1993 n=2836/2875 follow-up: 6 mo	intravenous alteplase (100 mg over 3 h) versus placebo	patients with symptoms and electrocardiographic criteria consistent with AMI between 6 and 24 h from symptom onset	Parallel groups double blind
TAMI 6 , 1992 n=96/101 follow-up: 6 months	tissue-type plasminogen activator 100 mg over 2 hours versus placebo	patients with 6 to 24 hours of symptoms and ECG ST elevation	Parallel groups double blind USA
bolus t-PA vs accelerated t-PA			
COBALT , 1997 n=3585/3584 follow-up: 30 days	of 50 mg of alteplase over a period of 1 to 3 minutes followed 30 minutes later by a second bolus of 50 mg (or 40 mg for patients who weighed less than 60 kg). versus weight-adjusted, accelerated infusion of 100 mg of alteplase	patients with acute myocardial infarction	Parallel groups double blind
lanoteplase vs accelerated t-PA			
InTIME-II , 2000 n=10038/5022 follow-up: 30 days	lanoteplase 120 KU. kg(-1) as a single intravenous bolus versus up to 100 mg accelerated alteplase given over 90 min	patients presenting within 6 h of onset of ST elevation acute myocardial infarction	Parallel groups double blind worldwide
reteplase vs accelerated t-PA			
GUSTO III , 1997 n=10138/4921 follow-up: 30 days	reteplase, in two bolus doses or 10 MU each given 30 minutes apart versus alteplase, up to 100 mg infused over a period of 90 minutes	patients within 6 hours after the onset of symptoms with ST-segment elevation or bundle-branch block	Parallel groups open 20 countries

continued...

Trial	Treatments	Patients	Trials design and methods
RAPID-2 , 1996 n=169/155 follow-up: 35 days	10 plus 10 megaunits double bolus of reteplase versus front-loaded alteplase	patients with acute myocardial infarction within 12h from onset of ischemic chest pain	Parallel groups open USA, Germany
tenecteplase vs accelerated t-PA			
ASSENT-2 , 1999 n=8461/8488 follow-up: 30d	Tenecteplase en IV bolus (dose en fonction du poids: 30 mg si <60 kg; 35 mg si poids entre 60 et 69.9 kg; 40 mg pour les 80-89.9 kg; 50 mg si >ou = 90 kg) versus Alteplase en IV, bolus de 15 mg, puis 0.75 mg/kg (sans dpasser 50 mg) en 30 min puis 0.50 mg/kg (sans dpasser 35 mg) en 60 min	patients with acute myocardial infarction of less than 6 h duration	Parallel groups double blind 29 countries
accelerated t-PA vs APSAC			
TAPS , 1992 n=199/202 follow-up:	front-loaded administration of rt-PA versus APSAC	patients with acute myocardial infarction.	Parallel groups open
TIMI 4 , 1994 n=NA follow-up: hospital stay	front-loaded rt-PA versus APSAC	patients with acute myocardial infarction	double blind
accelerated t-PA vs streptokinase			
GUSTO tPA Hiv , 1993 n=10396/20251 follow-up: 30 d	tPA acclr (15 mg en bolus, puis 0.75 mg/kg en 30 min sans dpasser 50 mg puis 0.5 mg/kg en 60 min sans dpasser 35 mg) + hparine en IV (5000 U en bolus, 1000 U/h (de prference 1200 U/h si >80 kg), poursuivi au moins 48 h) versus Streptokinase 1.5 MU en 60 min + hparine SC (12500 U 2 fois/j commence 4h aprs thrombolytique) combin streptokinase (1.5 MU en 60 min) + hparine en IV (5000 U en bolus, puis 1000 U/h (1200 U/h si >80 kg) poursuivi au moins 48 h)	Hommes et femmes	Parallel groups International 15 countries
anistreplase vs streptokinase			
TEAM 2 , 1991 n=183/176 follow-up:	anistreplase (30 units/2-5 min) versus streptokinase (1.5 million units/60 min	less than 76 years of age with electrocardiographic ST segment elevation who could be treated within 4 hours of symptom onset	double blind
APSAC vs streptokinase			
ISIS III (SK/APSAC) , 1992 n=13780/13773 follow-up: 6 mo	Streptokinase 1.5 MU infused over about 1 h versus anisoylated plasminogen-streptokinase activator complex (APSAC), anistreplase: 30 U over about 3 min	patients within 24 h of the onset of suspected acute myocardial infarction	Plan factoriel 3 (ou 4) *2 double blind International 17 countries
reteplase vs streptokinase			

continued...

Trial	Treatments	Patients	Trials design and methods
INJECT , 1995 n=3004/3006 follow-up: 6 mo	Retepase 2 bolus de 10 MU 30 min d'intervalle versus Streptokinase 1.5 MU en IV en 60 min	patients with symptoms and electrocardiographic criteria consistent with acute myocardial infarction within 12 h from onset of symptoms	Parallel groups double blind Europe
saruplase vs streptokinase			
COMPASS , 1998 n=1542/1547 follow-up: 1 y	saruplase 20-mg bolus and 60-mg infusion over 60 min versus streptokinase 1.5-MU infusion over 60 min	patients with symptoms compatible with those of acute myocardial infarction for <6 h	Parallel groups double blind
PRIMI (vs SK) , 1989 n=198/203 follow-up: ND	saruplase 20 mg bolus followed by 60 mg infusion for 60 min versus 1.5 million IU streptokinase infused over 60 min	patients with acute myocardial infarction were within 4 h of onset of symptoms	Parallel groups double blind
t-PA vs streptokinase			
International Study Group , 1990 n=10372/10396 follow-up: 6 mo	tPA 100 mg en IV en 3 h (10 mg en bolus, puis 50 mg en 1 h, puis 20 mg/h pendant 2 h) versus Streptokinase 1.5 MU en IV de 30 60 min	patients with suspected acute myocardial infarction of less than 6 h duration	Plan factoriel 2*2 double blind
Centre Illinois , 1993 n=123/130 follow-up:	t-PA 10 mg bolus, followed by 50 mg in the first hour, and 20 mg/hour for the next 2 hours versus SK 375 000 IU bolus, followed by 1 125 000 IU/1 hage/pj	patients with AMI within 3h from onset of chest pain	Parallel groups single blind USA
Cherng , 1992 n=59/63 follow-up: hospital stay	100 mg of rTPA over 3 hours (with early heparinization) versus 1,500,000 units of streptokinase over 1 hour	patients with acute myocardial infarction	Parallel groups open Taiwan
ECSG , 1985 n=64/65 follow-up:	0.75 mg rt-PA/kg over 90 min versus 1 500 000 IU streptokinase over 60 min	patients with acute myocardial infarction of less than 6 h duration	Parallel groups single-blind Europe
GISSI II , 1990 n=6182/6199 follow-up: 6 mo	alteplase 100 mg infused intravenously over 3 h versus streptokinase 1.5 MU infused intravenously over 30-60 min	patients with acute myocardial infarction within 6 h from onset of symptoms	Plan factoriel 2*2 open International 14 countries
ISIS III (SK/tPA) , 1992 n=13780/13746 follow-up: 6 mo	Streptokinase 1.5 MU en IV d'une heure versus tPA 0.04 MU/kg en IV en bolus d'1 min, puis 0.36 MU/kg en 1 h, puis 0.067 MU/kg/h pendant 3 h	Hommes et femmes	Plan factoriel 3 (ou 4) *2 double blind International 17 countries

continued...

Trial	Treatments	Patients	Trials design and methods
PAIMS , 1989 n=86/85 follow-up:	intravenous cumulative dose of 100 mg rt-PA versus .5 million units streptokinase	patients with acute myocardial infarction less than 3 h old	Parallel groups open Italy
TIMI-1 , 1987 [NCT00000505] n=157/159 follow-up:	rt-PA, 40, 20, and 20 mg in successive hours versus SK 1.5 million units over 1 hr	patients with evolving acute myocardial infarction within 7 hr of the onset of symptoms	Parallel groups double blind USA
White , 1989 n=135/135 follow-up:	rt-PA 100 mg over three hours versus streptokinase 1.5 million units over 30 minutes	patients with AMI	Parallel groups double blind New Zealand
t-PA + streptokinase vs streptokinase			
GUSTO tPA-SK Hiv , 1993 n=10374/20251 follow-up: 30 d	tPA en IV 1 mg/kg, sans dpasser 90 mg, dont 10 % en bolus + streptokinase 1 MU en 60 min + hparine en IV (5000 U en bolus, 1000 U/h (de prference 1200 U/h si >80 kg), poursuivi au moins 48 h) versus Streptokinase 1.5 MU en 60 min + hparine SC (12500 U 2 fois/j commence 4h aprs thrombolytique) combin streptokinase (1.5 MU en 60 min) + hparine en IV (5000 U en bolus, puis 1000 U/h (1200 U/h si >80 kg) poursuivi au moins 48 h)	Hommes et femmes	Parallel groups International 15 countries
accelerated t-PA vs t-PA			
RAAMI , 1992 n=143/138 follow-up: hospital stay	100 mg of rt-PA accelerated 90-min regimen (15-mg bolus followed by 50 mg over 30 min, then 35 mg over 60 min) versus 100 mg of rt-PA standard 3-h infusion regimen (an initial 10-mg bolus followed by 50 mg for the 1st h, then 20 mg/h for 2 h)	patients with acute myocardial infarction within 6h from onset of chest pain	Parallel groups open US
APSAC vs t-PA			
TEAM 3 , 1992 n=325 follow-up: 1 months	APSAC, 30 U/2 to 5 min versus rt-PA, 100 mg/3 h,	patient with ST elevation within 4h of the onset of symptoms	double blind
recombinant staphylokinase vs t-PA			
STAR , 1995 n=48/52 follow-up: 90 min	recombinant staphylokinase (10 or 20 mg given intravenously over 30 minutes) versus weight-adjusted rt-PA over 90 minutes	patients with evolving myocardial infarction of <6 hours' duration and with ST-segment elevation	Parallel groups open Belgium
saruplase vs t-PA			

continued...

Trial	Treatments	Patients	Trials design and methods
SESAM , 1997 n=236/237 follow-up: hospital stay	saruplase 80 mg/hour versus alteplase 100 mg every 3 hours	patients with acute myocardial infarction	Parallel groups open Europe
t-PA + urokinase vs t-PA			
TAMI 5 (t-PA+uroK vs tPA) , 1991 n=194/191 follow-up:	t-PA + urokinase versus t-PA	patient with acute myocardial infarction	open
t-PA half dose vs t-PA			
KAMIT , 1991 n=109/107 follow-up: hospital stay	half-dose (50 mg) t-PA with streptokinase (1.5 MU) during 1 hour versus t-PA (100 mg) during 3 hours	patients within 6 hours of myocardial infarction	Parallel groups open USA
saruplase vs urokinase			
PRIMI (vs UK) , 1989 n=198 follow-up:	20 mg bolus followed by 60 mg infusion for 60 min versus 80 mg recombinant pro-urokinase	with a first acute myocardial infarction within 4 h of onset of symptoms	Parallel groups double blind
t-PA vs urokinase			
TAMI 5 (t-PA vs uroK) , 1991 n=191/190 follow-up:	accelerated t-PA 100mg over 3h versus urokinase IV bolus 1.5 MU followed by 1.5 MU over 90min	patient with acute myocardial infarction	open

9

References

APSIM, 1989:

Bassand JP, Machecourt J, Cassagnes J, Anguenot T, Lusson R, Borel E, Peycelon P, Wolf E, Ducellier D Multicenter trial of intravenous anisoylated plasminogen streptokinase activator complex (APSAC) in acute myocardial infarction: effects on infarct size and left ventricular function. J Am Coll Cardiol 1989 Apr;13:988-97 [2647817]

USIM, 1991:

Rossi P, Bolognese L Comparison of intravenous urokinase plus heparin versus heparin alone in acute myocardial infarction. Urochinasi per via Sistemica nell'Infarto Miocardico (USIM) Collaborative Group. Am J Cardiol 1991;68:585-92 [1877476]

AIMS, 1988:

Effect of intravenous APSAC on mortality after acute myocardial infarction: preliminary report of a placebo-controlled clinical trial. AIMS Trial Study Group. Lancet 1988 Mar 12;1:545-9 [2894490]

Long-term effects of intravenous anistreplase in acute myocardial infarction: final report of the AIMS study. AIMS Trial Study Group. Lancet 1990 Feb 24;335:427-31 [1968167]

German Multicenter Trial, 1988:

EMERAS (7-12h), 1993:

Randomised trial of late thrombolysis in patients with suspected acute myocardial infarction. EMERAS (Estudio Multicentrico Estreptoquinasa Replicas de Amrica del Sur) Collaborative Group. Lancet 1993 Sep 25;342:767-72 [8103875]

EMERAS (all delay), 1993:

Randomised trial of late thrombolysis in patients with suspected acute myocardial infarction. EMERAS (Estudio Multicentrico Estreptoquinasa Replicas de Amrica del Sur) Collaborative Group. Lancet 1993;342:767-72 [8103875]

GISSI I, 1986:

Effectiveness of intravenous thrombolytic treatment in acute myocardial infarction. Gruppo Italiano per lo Studio della Streptochinasi nell'Infarto Miocardico (GISSI). Lancet 1986 Feb 22;1:397-402 [2868337]

Long-term effects of intravenous thrombolysis in acute myocardial infarction: final report of the GISSI study. Gruppo Italiano per lo Studio della Streptochinasi nell'Infarto Miocardico (GISSI). Lancet 1987 Oct 17;2:871-4 [2889079]

ISAM, 1986:

A prospective trial of intravenous streptokinase in acute myocardial infarction (I.S.A.M.). Mortality, morbidity, and infarct size at 21 days. The I.S.A.M. Study Group. N Engl J Med 1986 Jun 5;314:1465-71 [2871492] 10.1056/NEJM198606053142301

ISIS 2 pilot, 1987:

Randomized factorial trial of high-dose intravenous streptokinase, of oral aspirin and of intravenous heparin in acute myocardial infarction. ISIS (International Studies of Infarct Survival) pilot study. Eur Heart J 1987 Jun;8:634-42 [2887430]

ISIS-2 (SK), 1988:

Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. Lancet 1988 Aug 13;2:349-60 [2899772]

Western Washington Intravenous Trial, 1988:**ASSET, 1988:**

Wilcox RG, von der Lippe G, Olsson CG, Jensen G, Skene AM, Hampton JR Trial of tissue plasminogen activator for mortality reduction in acute myocardial infarction. Anglo-Scandinavian Study of Early Thrombolysis (ASSET). Lancet 1988 Sep 3;2:525-30 [2900919]

LATE, 1993:

Late Assessment of Thrombolytic Efficacy (LATE) study with alteplase 6-24 hours after onset of acute myocardial infarction. Lancet 1993 Sep 25;342:759-66 [8103874]

TAMI 6, 1992:

Topol EJ, Califf RM, Vandormael M, Grines CL, George BS, Sanz ML, Wall T, O'Brien M, Schwaiger M, Aguirre FV A randomized trial of late reperfusion therapy for acute myocardial infarction. Thrombolysis and Angioplasty in Myocardial Infarction-6 Study Group. Circulation 1992 Jun;85:2090-9 [1591828]

COBALT, 1997:

A comparison of continuous infusion of alteplase with double-bolus administration for acute myocardial infarction. The Continuous Infusion versus Double-Bolus Administration of Alteplase (COBALT) Investigators. N Engl J Med 1997 Oct 16;337:1124-30 [9340504] 10.1056/NEJM199710163371604

InTIME-II, 2000:

Intravenous NPA for the treatment of infarcting myocardium early; InTIME-II, a double-blind comparison of single-bolus alteplase vs accelerated alteplase for the treatment of patients with acute myocardial infarction Eur Heart J 2000;21:2005-13 [11102251] 10.1053/euhj.2000.2498

GUSTO III, 1997:

A comparison of reteplase with alteplase for acute myocardial infarction. The Global Use of Strategies to Open Occluded Coronary Arteries (GUSTO III) Investigators. N Engl J Med 1997 Oct 16;337:1118-23 [9340503] 10.1056/NEJM199710163371603

A comparison of reteplase with alteplase for acute myocardial infarction. The Global Use of Strategies to Open Occluded Coronary Arteries (GUSTO III) Investigators. N Engl J Med 1997;337:1118-23 [9340503]

RAPID-2, 1996:

Bode C, Smalling RW, Berg G, Burnett C, Lorch G, Kalbfleisch JM, Chernoff R, Christie LG, Feldman RL, Seals AA, Weaver WD Randomized comparison of coronary thrombolysis achieved with double-bolus reteplase (recombinant plasminogen activator) and front-loaded, accelerated alteplase (recombinant tissue plasminogen activator) in patients with acute myocardial infarction. The RAPID II Investigators. Circulation 1996;94:891-8 [8790022]

ASSENT-2, 1999:

Van De Werf F, Adgey J, Ardissino D, Armstrong PW, Aylward P, Barbash G, Betriu A, Binbrek AS, Califf R, Diaz R, Fanestus R, Fox K, Granger C, Heikkilä J, Husted S, Jansky P, Langer A, Lupi E, Maseri A, Meyer J, Mlczoch J, Moccetti D, Myburgh D, Oto A, Pao Single-bolus tenecteplase compared with front-loaded alteplase in acute myocardial infarction:

the ASSENT-2 double-blind randomised trial. *Lancet* 1999 Aug 28;354:716-22 [[10475182](#)]

TAPS, 1992:

Neuhaus KL, von Essen R, Tebbe U, Vogt A, Roth M, Riess M, Niederer W, Forycki F, Wirtzfeld A, Maeurer W Improved thrombolysis in acute myocardial infarction with front-loaded administration of alteplase: results of the rt-PA-APSAC patency study (TAPS) *J Am Coll Cardiol* 1992 Apr;19:885-91 [[1552106](#)]

TIMI 4, 1994:

Cannon CP, McCabe CH, Diver DJ, Herson S, Greene RM, Shah PK, Sequeira RF, Leya F, Kirshenbaum JM, Magorien RD Comparison of front-loaded recombinant tissue-type plasminogen activator, anistreplase and combination thrombolytic therapy for acute myocardial infarction: results of the Thrombolysis in Myocardial Infarction (TIMI) 4 trial. *J Am Coll Cardiol* 1994 Dec;24:1602-10 [[7963104](#)]

GUSTO tPA Hiv, 1993:

An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. The GUSTO investigators. *N Engl J Med* 1993 Sep 2;329:673-82 [[8204123](#)]

TEAM 2, 1991:

Anderson JL, Sorensen SG, Moreno FL, Hackworthy RA, Browne KF, Dale HT, Leya F, Dangoisse V, Eckerson HW, Marder VJ Multicenter patency trial of intravenous anistreplase compared with streptokinase in acute myocardial infarction. The TEAM-2 Study Investigators. *Circulation* 1991 Jan;83:126-40 [[1984877](#)]

ISIS III (SK/APSAC), 1992:

ISIS-3: a randomised comparison of streptokinase vs tissue plasminogen activator vs anistreplase and of aspirin plus heparin vs aspirin alone among 41,299 cases of suspected acute myocardial infarction. ISIS-3 (Third International Study of Infarct Survival) Collaborative Group. *Lancet* 1992;339:753-70 [[1347801](#)]

INJECT, 1995:

Randomised, double-blind comparison of reteplase double-bolus administration with streptokinase in acute myocardial infarction (INJECT): trial to investigate equivalence. International Joint Efficacy Comparison of Thrombolytics *Lancet* 1995;346:329-36 [[7623530](#)]

COMPASS, 1998:

Tebbe U, Michels R, Adgey J, Boland J, Caspi A, Charbonnier B, Windeler J, Barth H, Groves R, Hopkins GR, Fennell W, Betriu A, Ruda M, Mlczoch J Randomized, double-blind study comparing saruplase with streptokinase therapy in acute myocardial infarction: the COMPASS Equivalence Trial. Comparison Trial of Saruplase and Streptokinase (COMASS) Investigators *J Am Coll Cardiol* 1998;31:487-93 [[9502624](#)]

PRIMI (vs SK), 1989:

Randomised double-blind trial of recombinant pro-urokinase against streptokinase in acute myocardial infarction. PRIMI Trial Study Group *Lancet* 1989;1:863-8 [[2564949](#)]

International Study Group, 1990:

In-hospital mortality and clinical course of 20,891 patients with suspected acute myocardial infarction randomised between alteplase and streptokinase with or without heparin. The International Study Group. *Lancet* 1990;336:71-5 [[1975322](#)]

Centre Illinois, 1993:

Taylor GJ, Moses HW, Koester D, Colliver JA, Katholi RE, Dove JT, Woodruff RC, Mikell FL, Becker LC, Sheehan FH A difference between front-loaded streptokinase and standard-dose recombinant tissue-type plasminogen activator in preserving left ventricular function after acute myocardial infarction (the Central Illinois Thrombolytic Therapy Study). *Am J Cardiol* 1993;72:1010-4 [[8213579](#)]

Cherng, 1992:

Cherng WJ, Chiang CW, Kuo CT, Lee CP, Lee YS A comparison between intravenous streptokinase and tissue plasminogen activator with early intravenous heparin in acute myocardial infarction. *Am Heart J* 1992;123:841-6 [[1549990](#)]

ECSG, 1985:

Verstraete M, Bernard R, Bory M, Brower RW, Collen D, de Bono DP, Erbel R, Huhmann W, Lennane RJ, Lubsen J Randomised trial of intravenous recombinant tissue-type plasminogen activator versus intravenous streptokinase in acute myocardial infarction. Report from the European Cooperative Study Group for Recombinant Tissue-type Plasminogen Activator. *Lancet* 1985;1:842-7 [[2858711](#)]

GISSI II, 1990:

GISSI-2: a factorial randomised trial of alteplase versus streptokinase and heparin versus no heparin among 12,490 patients with acute myocardial infarction. Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico. *Lancet* 1990;336:65-71 [[1975321](#)]

ISIS III (SK/tPA), 1992:

ISIS-3: a randomised comparison of streptokinase vs tissue plasminogen activator vs anistreplase and of aspirin plus heparin vs aspirin alone among 41,299 cases of suspected acute myocardial infarction. ISIS-3 (Third International Study of Infarct Survival) Collaborative Group. *Lancet* 1992 Mar 28;339:753-70 [[1347801](#)]

PAIMS, 1989:

Magnani B Plasminogen Activator Italian Multicenter Study (PAIMS): comparison of intravenous recombinant single-chain human tissue-type plasminogen activator (rt-PA) with intravenous streptokinase in acute myocardial infarction. *J Am Coll Cardiol* 1989;13:19-26 [[2491867](#)]

TIMI-1, 1987:

Chesebro JH, Knatterud G, Roberts R, Borer J, Cohen LS, Dalen J, Dodge HT, Francis CK, Hillis D, Ludbrook P Thrombolysis in Myocardial Infarction (TIMI) Trial, Phase I: A comparison between intravenous tissue plasminogen activator and intravenous streptokinase. Clinical findings through hospital discharge. *Circulation* 1987;76:142-54 [[3109764](#)]

White, 1989:

White HD, Rivers JT, Maslowski AH, Ormiston JA, Takayama M, Hart HH, Sharpe DN, Whitlock RM, Norris RM Effect of intravenous streptokinase as compared with that of tissue plasminogen activator on left ventricular function after first myocardial infarction. *N Engl J Med* 1989;320:817-21 [[2494454](#)]

GUSTO tPA-SK Hiv, 1993:

An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. The GUSTO investigators. *N Engl J Med* 1993 Sep 2;329:673-82 [[8204123](#)]

RAAMI, 1992:

Carney RJ, Murphy GA, Brandt TR, Daley PJ, Pickering E, White HJ, McDonough TJ, Vermilya SK, Teichman SL Randomized angiographic trial of recombinant tissue-type plasminogen activator (alteplase) in myocardial infarction. RAAMI Study Investigators. *J Am Coll Cardiol* 1992 Jul;20:17-23 [[1607520](#)]

TEAM 3, 1992:

Anderson JL, Becker LC, Sorensen SG, Karagounis LA, Browne KF, Shah PK, Morris DC, Fintel DJ, Mueller HS, Ross AM Anistreplase versus alteplase in acute myocardial infarction: comparative effects on left ventricular function, morbidity and 1-day coronary artery patency. The TEAM-3 Investigators. *J Am Coll Cardiol* 1992 Oct;20:753-66 [[1527286](#)]

STAR, 1995:

Vanderschueren S, Barrios L, Kerdsinchai P, Van den Heuvel P, Hermans L, Vrolix M, De Man F, Benit E, Muyldermans L, Collen D A randomized trial of recombinant staphylokinase versus alteplase for coronary artery patency in acute myocardial infarction. The STAR Trial Group. *Circulation* 1995 Oct 15;92:2044-9 [[7554180](#)]

SESAM, 1997:

Br FW, Meyer J, Vermeer F, Michels R, Charbonnier B, Haerten K, Spiecker M, Macaya C, Hanssen M, Heras M, Boland JP, Morice MC, Dunn FG, Uebis R, Hamm C, Ayzenberg O, Strupp G, Withagen AJ, Klein W, Windeler J, Hopkins G, Barth H, von Fisenne MJ Comparison of saruplase and alteplase in acute myocardial infarction. SESAM Study Group. The Study in Europe with Saruplase and Alteplase in Myocardial Infarction. *Am J Cardiol* 1997 Mar 15;79:727-32 [[9070549](#)]

TAMI 5 (t-PA+uroK vs tPA), 1991:

Califf RM, Topol EJ, Stack RS, Ellis SG, George BS, Kereiakes DJ, Samaha JK, Worley SJ, Anderson JL, Harrelson-Woodlief L Evaluation of combination thrombolytic therapy and timing of cardiac catheterization in acute myocardial infarction. Results of thrombolysis and angioplasty in myocardial infarction—phase 5 randomized trial. TAMI Study Group. *Circulation* 1991 May;83:1543-56 [[1902405](#)]

KAMIT, 1991:

Grines CL, Nissen SE, Booth DC, Gurley JC, Chelliah N, Wolf R, Blankenship J, Branco MC, Bennett K, DeMaria AN A prospective, randomized trial comparing combination half-dose tissue-type plasminogen activator and streptokinase with full-dose tissue-type plasminogen activator. Kentucky Acute Myocardial Infarction Trial (KAMIT) Group. *Circulation* 1991;84:540-9 [[1907228](#)]

PRIMI (vs UK), 1989:

Randomised double-blind trial of recombinant pro-urokinase against streptokinase in acute myocardial infarction. PRIMI Trial Study Group. *Lancet* 1989 Apr 22;1:863-8 [[2564949](#)]

TAMI 5 (t-PA vs uroK), 1991:

Califf RM, Topol EJ, Stack RS, Ellis SG, George BS, Kereiakes DJ, Samaha JK, Worley SJ, Anderson JL, Harrelson-Woodlief L Evaluation of combination thrombolytic therapy and timing of cardiac catheterization in acute myocardial infarction. Results of thrombolysis and angioplasty in myocardial infarction—phase 5 randomized trial. TAMI Study Group. *Circulation* 1991 May;83:1543-56 [[1902405](#)]

2 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.