

Clinical trials of lifestyle intervention for cardiovascular prevention in all type of patients

TrialResults-center www.trialresultscenter.org

1 diet

Trial	Treatments	Patients	Trials design and methods
diet vs control			
NORDIET n=45/43 follow-up:	healthy Nordic diet versus control diet (subjects usual Western diet)	mildly hypercholesterolaemic subjects	Sweden
BARON n=NA	-	-	
HPT n=NA	-	-	
Kumanyika n=NA	-	-	
TAIM n=NA	-	-	
DISH n=NA	-	-	
diet vs usual diet			
Black , 1994 n=56/55 follow-up: 2.0 years	diet with 20 percent of total caloric intake as fat versus usual diet	patients with nonmelanoma skin cancer	Parallel groups open
DART (Burr) , 1989 n=NA follow-up: 2 years	diet advice versus usual diet	men who had recovered from MI	Factorial plan open, blind assessment
Finnish Mental Hospital (Miettinen) , 1985 n=612/610 follow-up: 6.0 years	cholesterol-lowering diet (low in saturated fats and cholesterol and relatively high in polyunsaturated fats) versus usual diet	middle-aged institutionalized women without CHD	Cluster-randomized cross-ove open, blind assessment Finland
Finnish Mental Hospital (Turpeinen) , 1979 n=NA follow-up: 6.0 years	cholesterol-lowering diet (low in saturated fats and cholesterol and relatively high in polyunsaturated fats) versus usual diet	middle-aged institutionalized men without CHD	Cluster-randomized cross-ove open, blind assessment Finland
Goteborg , 1986 n=10004/20028 follow-up: 10 years	multifactorial intervention programme versus no intervention	men, 47-55 years old at entry	Parallel groups open Sweden

continued...

Trial	Treatments	Patients	Trials design and methods
Gteborg (Wilhelmsen) , 1986 n=10004/20028 follow-up: 10.0 years	multifactorial intervention programme versus usual care	men, 47-55 years old at entry	Parallel groups open
Hjermann , 1981 n=604/628 follow-up: 6.5 years	diet versus usual diet	healthy, normotensive men at high risk of coronary heart disease	Parallel groups open Sweden
Kallio , 1979 n=188/187 follow-up: 3.0 years	diet (multifactorial intervention programme) versus usual diet	patients below 65 years who had an acute myocardial infarction	Parallel groups open
Los Angeles VA (Dayton) , 1969 n=424/422 follow-up: 65279;8.0 y	diet versus usual diet	men in domiciliary care, age>55, with or without CHD	Parallel groups double blind USA
Minnesota coronary survey (Frantz) , 1975 n=2197/2196 follow-up: 1.1 y (max 4.5y)	cholesterol lowering diet versus control diet	65279;Adult residents of mental hospitals; no illness restrictions, no cholesterol concentration requirements	Parallel groups double-blind USA
MRC low fat , 1965 n=123/129 follow-up: 3 y	-	-	Parallel groups open
MRC Soya , 1968 n=199/194 follow-up: 3.5 y	Rgime pauvre en graisses satures + 85 g/j d'huile de soja versus usual diet	ambulatory men with recent MI	Parallel groups open, blind assessment
MRFIT , 1982 n=6428/6438 follow-up: 6.5 y	multifactor intervention program versus usual diet	high-risk men aged 35 to 57 years	Parallel groups open
Ornish , 1990 n=28/20 follow-up: 1.0 y	low-fat vegetarian diet, stopping smoking, stress management training, and moderate exercise versus usual-care	Patients with angiographically documented coronary artery disease	Parallel groups open USA
Oslo Diet Heart Study (Leren) , 1966 n=206/206 follow-up: 5 y (11y)	diet versus usual care	middle-aged ambulatory men with prior MI	Parallel groups open, blind assessment
Rose , 1965 n=28/26 follow-up: 1.2 years	Rgime restreint en graisses + 80 g/j huile de mas versus usual diet	men, <70 years	Parallel groups open
Singh , 1992 n=204/202 follow-up: 65279;2.0 years	strict diet versus usual diet	patients with suspected acute myocardial infarction	Parallel groups open

continued...

Trial	Treatments	Patients	Trials design and methods
STARS (St Thomas, diet) , 1992 n=30/30 follow-up: 3.0 years	dietary advice versus usual diet	patients with angina or past myocardial infarction	open, blind assessment
Veterans Ad. (Dayton) , 1969 n=424/422 follow-up: 3.6 and 8 y	cholesterol lowering diet versus usual diet	men in domiciliary care, age>55, with or without CHD	Parallel groups double blind USA
WHI low fat , 2005 [NCT00000611] n=19541/29294 follow-up: 8.1y mean	dietary modification intervention to promote dietary change with the goals of reducing intake of total fat to 20% of energy and increasing consumption of vegetables and fruit to at least 5 servings daily and grains to at least 6 servings daily versus usual diet	postmenopausal women, aged 50 to 79 years, without prior breast cancer	Parallel groups open US
Woodhill , 1966 n=221/237 follow-up: <7 years	diet versus usual diet	men, 30-59 years	Parallel groups open
low fat diet vs mediterranean-style diet			
Tuttle , 2008 n=NA follow-up: 24 months	low-fat versus Mediterranean-style diets	First MI survivors	Parallel groups open

33

References

NORDIET, :

Adamsson V, Reumark A, Fredriksson IB, Hammarström E, Vessby B, Johansson G, Risrus U Effects of a healthy Nordic diet on cardiovascular risk factors in hypercholesterolaemic subjects: a randomized controlled trial (NORDIET). *J Intern Med* 2011;269:150-9 [20964740] [10.1111/j.1365-2796.2010.02290.x](https://doi.org/10.1111/j.1365-2796.2010.02290.x)

BARON, :

Baron JA, Gleason R, Crowe B, Mann JI Preliminary trial of the effect of general practice based nutritional advice. *Br J Gen Pract* 1990;40:137-41 [2115348]

HPT, :

The Hypertension Prevention Trial: three-year effects of dietary changes on blood pressure. Hypertension Prevention Trial Research Group. *Arch Intern Med* 1990;150:153-62 [2404477]

Kumanyika, :

Kumanyika SK, Hebert PR, Cutler JA, Lasser VI, Sugars CP, Steffen-Batey L, Brewer AA, Cameron M, Shepek LD, Cook NR Feasibility and efficacy of sodium reduction in the Trials of Hypertension Prevention, phase I. Trials of Hypertension Prevention Collaborative Research Group. *Hypertension* 1993;22:502-12 [8406655]

TAIM, :

Davis BR, Oberman A, Blafox MD, Wassertheil-Smoller S, Zimbaldi N, Kirchner K, Wylie-Rosett J, Langford HG Lack of effectiveness of a low-sodium/high-potassium diet in reducing antihypertensive medication requirements in overweight persons with mild hypertension. TAIM Research Group. Trial of Antihypertensive Interventions and Management. *Am J Hypertens* 1994;7:926-32 [7826557]

DISH, :

Langford HG, Blafox MD, Oberman A, Hawkins CM, Curb JD, Cutter GR, Wassertheil-Smoller S, Pressel S, Babcock C, Abernethy JD Dietary therapy slows the return of hypertension after stopping prolonged medication. *JAMA* 1985;253:657-64 [3881608]

Black, 1994:

Black HS, Herd JA, Goldberg LH, Wolf JE Jr, Thornby JI, Rosen T, Bruce S, Tschen JA, Foreyt JP, Scott LW Effect of a low-fat diet on the incidence of actinic keratosis. *N Engl J Med* 1994;330:1272-5 [8145782]

DART (Burr), 1989:

Burr ML, Fehily AM, Gilbert JF, Rogers S, Holliday RM, Sweetnam PM, Elwood PC, Deadman NM Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: diet and reinfarction trial (DART). *Lancet* 1989;2:757-61 [2571009]

Finnish Mental Hospital (Miettinen), 1985:

European collaborative trial of multifactorial prevention of coronary heart disease: final report on the 6-year results. World Health Organisation European Collaborative Group. *Lancet* 1986;1:869-72 [2870351]

Miettinen TA, Huttunen JK, Naukkarinen V, Strandberg T, Mattila S, Kumlin T, Sarna S Multifactorial primary prevention of cardiovascular diseases in middle-aged men. Risk factor changes, incidence, and mortality. *JAMA* 1985;254:2097-102 [4046137]

Miettinen M, Turpeinen O, Karvonen MJ, Pekkarinen M, Paavilainen E, Elosuo R Dietary prevention of coronary heart disease in women: the Finnish mental hospital study. *Int J Epidemiol* 1983;12:17-25 [6840954]

Finnish Mental Hospital (Turpeinen), 1979:

Turpeinen O, Karvonen MJ, Pekkarinen M, Miettinen M, Elosuo R, Paavilainen E Dietary prevention of coronary heart disease: the Finnish Mental Hospital Study. *Int J Epidemiol* 1979;8:99-118 [393644]

Goteborg, 1986:

Wilhelmsen L, Berglund G, Elmfeldt D, Tibblin G, Wedel H, Pennert K, Vedin A, Wilhelmsson C, Werk L The multifactor primary prevention trial in Gteborg, Sweden. *Eur Heart J* 1986;7:279-88 [3720755]

Gteborg (Wilhelmsen), 1986:

Wilhelmsen L, Berglund G, Elmfeldt D, Tibblin G, Wedel H, Pennert K, Vedin A, Wilhelmsson C, Werk L The multifactor primary prevention trial in Gteborg, Sweden. *Eur Heart J* 1986;7:279-88 [3720755]

Hjermann, 1981:

Hjermann I, Velve Byre K, Holme I, Leren P Effect of diet and smoking intervention on the incidence of coronary heart disease. Report from the Oslo Study Group of a randomised trial in healthy men. *Lancet* 1981;2:1303-10 [6118715]

Kallio, 1979:

Kallio V, Hmlinen H, Hakkila J, Luurila OJ Reduction in sudden deaths by a multifactorial intervention programme after acute myocardial infarction. *Lancet* 1979;2:1091-4 [91836]

Los Angeles VA (Dayton), 1969:

Rogers MC Sir John Scott Burdon-Sanderson (1828-1905): a pioneer in electrophysiology. *Circulation* 1969;40:1-2 [4893441]

Dayton S, Pearce ML Diet high in unsaturated fat. A controlled clinical trial. *Minn Med* 1969;52:1237-42 [4896402]

Minnesota coronary survey (Frantz), 1975:

Frantz ID Jr, Dawson EA, Ashman PL, Gatewood LC, Bartsch GE, Kuba K, Brewer ER Test of effect of lipid lowering by diet on cardiovascular risk. The Minnesota Coronary Survey. *Arteriosclerosis* 1989;9:129-35 [2643423]

MRC low fat, 1965:

, Low-fat diet in myocardial infarction: A controlled trial. *Lancet* 1965; 2:501-4 [4158171]

MRC Soya, 1968:

, Controlled trial of soya-bean oil in myocardial infarction. *Lancet* 1968; 2:693-9 [4175085]

MRFIT, 1982:

Multiple risk factor intervention trial. Risk factor changes and mortality results. Multiple Risk Factor Intervention Trial Research Group. *JAMA* 1982;248:1465-77 [7050440]

Ornish, 1990:

Ornish D, Brown SE, Scherwitz LW, Billings JH, Armstrong WT, Ports TA, McLanahan SM, Kirkeeide RL, Brand RJ, Gould KL Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial. Lancet 1990;336:129-33 [1973470]

Oslo Diet Heart Study (Leren), 1966:

Leren P, The Oslo diet-heart study. Eleven-year report. Circulation 1970; 42:935-42 [5477261]

Rose, 1965:

ROSE GA, THOMSON WB, WILLIAMS RT CORN OIL IN TREATMENT OF ISCHAEMIC HEART DISEASE. Br Med J 1965 Jun 12;1:1531-3 [14288105]

Singh, 1992:

Singh RB, Rastogi SS, Verma R, Laxmi B, Singh R, Ghosh S, Niaz MA Randomised controlled trial of cardioprotective diet in patients with recent acute myocardial infarction: results of one year follow up. BMJ 1992;304:1015-9 [1586782]

STARS (St Thomas, diet), 1992:

Watts GF, Lewis B, Brunt JN, Lewis ES, Coltart DJ, Smith LD, Mann JI, Swan AV Effects on coronary artery disease of lipid-lowering diet, or diet plus cholestyramine, in the St Thomas' Atherosclerosis Regression Study (STARS) Lancet 1992;339:563-9 [1347091]

Veterans Ad. (Dayton), 1969:

Dayton S, Pearce ML, Hashimoto S, Dixon WJ, Tomiyasu U. A controlled clinical trial of a diet high in unsaturated fat in preventing complications of atherosclerosis. Circulation 1969; 40(supp 2):1-55 [0]

WHI low fat, 2005:

Howard BV, Van Horn L, Hsia J, Manson JE, Stefanick ML, Wassertheil-Smoller S, Kuller LH, LaCroix AZ, Langer RD, Lasser NL, Lewis CE, Limacher MC, Margolis KL, Mysiw WJ, Ockene JK, Parker LM, Perri MG, Phillips L, Prentice RL, Robbins J, Rossouw JE, Sarto Low-fat dietary pattern and risk of cardiovascular disease: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. JAMA 2006 Feb 8;295:655-66 [16467234]

Woodhill, 1966:

Woodhill JM, Palmer AJ, Leelarthae-pin B, McGilchrist C, Blacket RB, Low fat, low cholesterol diet in secondary prevention of coronary heart disease. Adv Exp Med Biol 1978; 109:317-30 [727035]

Tuttle, 2008:

Tuttle KR, Shuler LA, Packard DP, Milton JE, Daratha KB, Bibus DM, Short RA Comparison of low-fat versus Mediterranean-style dietary intervention after first myocardial infarction (from The Heart Institute of Spokane Diet Intervention and Evaluation Trial). Am J Cardiol 2008;101:1523-30 [18489927]

2 Mediterranean diet

Trial	Treatments	Patients	Trials design and methods
Mediterranean diet vs control			
Lyon n=302/303 follow-up:	-	-	
Mediterranean diet with EOVV vs control			
PREDIMED (olive oil) , 2013 [ISRCTN35739639] n=2543/2450 follow-up: 4.8 years	Mediterranean diet supplemented with extra-virgin olive oil versus control diet (advice to reduce dietary fat)	participants who were at high cardiovascular risk, but with no cardiovascular disease	Parallel groups open Spain
Mediterranean diet with nuts vs control			

continued...

Trial	Treatments	Patients	Trials design and methods
PREDIMED (nuts) , 2013 [ISRCTN35739639] n=2454/2450 follow-up: 4.8 years	Mediterranean diet supplemented with mixed nuts versus control diet (advice to reduce dietary fat)	participants who were at high cardiovascular risk, but with no cardiovascular disease	open Spain

References

Lyon, :

de Lorgeril M, Renaud S, Mamelle N, Salen P, Martin JL, Monjaud I, Guidollet J, Touboul P, Delaye J Mediterranean alpha-linolenic acid-rich diet in secondary prevention of coronary heart disease. *Lancet* 1994;343:1454-9 [7911176]

De Lorgeril M, Salen P, Martin JL, Mamelle N, Monjaud I, Touboul P, Delaye J Effect of a mediterranean type of diet on the rate of cardiovascular complications in patients with coronary artery disease. *Insights into the cardioprotective effect of certain nutriments. J Am Coll Cardiol* 1996;28:1103-8 [8890801] [10.1016/S0735-1097\(96\)00280-X](https://doi.org/10.1016/S0735-1097(96)00280-X)

de Lorgeril M, Renaud S, Mamelle N, Salen P, Martin JL, Monjaud I, Guidollet J, Touboul P, Delaye J Mediterranean alpha-linolenic acid-rich diet in secondary prevention of coronary heart disease. *Lancet* 1994;343:1454-9 [7911176]

De Lorgeril M, Salen P, Martin JL, Mamelle N, Monjaud I, Touboul P, Delaye J Effect of a mediterranean type of diet on the rate of cardiovascular complications in patients with coronary artery disease. *Insights into the cardioprotective effect of certain nutriments. J Am Coll Cardiol* 1996;28:1103-8 [8890801] [10.1016/S0735-1097\(96\)00280-X](https://doi.org/10.1016/S0735-1097(96)00280-X)

PREDIMED (olive oil), 2013:

Estruch R, Ros E, Salas-Salvad J, Covas MI, D Pharm, Corella D, Ars F, Gmez-Gracia E, Ruiz-Gutierrez V, Fiol M, Lapetra J, Lamuela-Raventos RM, Serra-Majem L, Pint X, Basora J, Muoz MA, Sorl JV, Martnez JA, Martnez-Gonzlez MA Primary Prevention of Cardiovascular Disease with a Mediterranean Diet. *N Engl J Med* 2013 Feb 25;: [23432189] [10.1056/NEJMoa1200303](https://doi.org/10.1056/NEJMoa1200303)

PREDIMED (nuts), 2013:

Estruch R, Ros E, Salas-Salvad J, Covas MI, D Pharm, Corella D, Ars F, Gmez-Gracia E, Ruiz-Gutierrez V, Fiol M, Lapetra J, Lamuela-Raventos RM, Serra-Majem L, Pint X, Basora J, Muoz MA, Sorl JV, Martnez JA, Martnez-Gonzlez MA Primary Prevention of Cardiovascular Disease with a Mediterranean Diet. *N Engl J Med* 2013 Feb 25;: [23432189] [10.1056/NEJMoa1200303](https://doi.org/10.1056/NEJMoa1200303)

3 Multiple risk factor interventions

Trial	Treatments	Patients	Trials design and methods
Multiple risk factor interventions vs control			
CELL , 1995 n=339/320 follow-up: 18 months	intensive" health care advice through six group sessions versus usual care	subjects aged 30-59 years, with at least two cardiovascular risk factors in addition to moderately high lipid concentrations: total cholesterol >or = 6.5 mmol/l on three occasions, triglycerides <4.0 mmol/l, and ratio of low density lipoprotein cholesterol to high density lipoprotein cholesterol >4.0	Factorial plan open

continued...

Trial	Treatments	Patients	Trials design and methods
Family Heart , 1994 n=3436/5912 follow-up: 1 y	Nurse led programme using a family centred approach with follow up according to degree of risk. Counselling on diet, weight, smoking, exercise, alcohol versus control	men aged 40-59 and their partners	Parallel groups double-blind UK
Gteborg Study , 1986 n=10004/20018 follow-up: 11.8 yr	multifactorial intervention programme on coronary heart disease versus no intervention	random sample of men age 47-55 y	open Sweden
HDFP , 1979 [NCT00000498] n=5485/5455 follow-up: 5 yr	Stepped care: Antihypertensive drugs, diet, smoking advice, weight control, exercise versus usual primary care	persons with high blood pressure	Parallel groups open USA
Helsinki Businessmen Study , 1985 n=612/610 follow-up: 5 yr	Multifactorial prevention of cardiovascular diseases versus no intervention	healthy men 40-58 y at high risk	Parallel groups open Finland
Johns Hopkins , 1983 n=350/50 follow-up: 5 yr	health education interventions versus control	hypertensives men and women	Factorial plan open USA
Meland , 1997 n=69/58 follow-up: 1 y	patient-centred, self-directive intervention of lifestyle changes in general practice versus conventional care	men with high coronary heart disease risk	Parallel groups open
MRFIT , 1982 [NCT00000487?acronym=] n=6428/6438 follow-up: 6 yr	special intervention (SI) program consisting of stepped-care treatment for hypertension, counseling for cigarette smoking, and dietary advice for lowering blood cholesterol levels versus no intervention	high-risk men aged 35 to 57 years	Parallel groups open
Oslo , 1981 n=612/610 follow-up: 5 yr	recommendation to lower their blood lipids by change of diet and to stop smoking versus no intervention	healthy, normotensive men at high risk of coronary heart disease	Parallel groups open Oslo, Norway
OXCHECK , 1994 n=8307/2783 follow-up: 3 yr	health checks by nurses versus no intervention	patients from general practice aged 35-64 years	Parallel groups open UK
WHO Factories , 1982 n=30489/26971 follow-up: 6 years	multifactorial prevention of coronary heart disease versus no intervention	men employed in 80 factories in Belgium, Italy, Poland, and the UK	Parallel groups open Belgium, Italy, Poland, and the UK

References

CELL, 1995:

Lindholm LH, Ekblom T, Dash C, Eriksson M, Tibblin G, Schersten B The impact of health care advice given in primary care on cardiovascular risk. CELL Study Group. *BMJ* 1995;310:1105-9 [[7742677](#)]

Lindholm LH, Ekblom T, Dash C, Isacson A, Schersten B Changes in cardiovascular risk factors by combined pharmacological and nonpharmacological strategies: the main results of the CELL Study. *J Intern Med* 1996;240:13-22 [[8708586](#)]

Family Heart, 1994:

Randomised controlled trial evaluating cardiovascular screening and intervention in general practice: principal results of British family heart study. Family Heart Study Group. *BMJ* 1994;308:313-20 [[8124121](#)]

Gteborg Study, 1986:

Wilhelmsen L, Berglund G, Elmfeldt D, Tibblin G, Wedel H, Pennert K, Vedin A, Wilhelmsson C, Werk L The multifactor primary prevention trial in Gteborg, Sweden. *Eur Heart J* 1986;7:279-88 [[3720755](#)]

HDFP, 1979:

Five-year findings of the hypertension detection and follow-up program. I. Reduction in mortality of persons with high blood pressure, including mild hypertension. Hypertension Detection and Follow-up Program Cooperative Group. *JAMA* 1979;242:2562-71 [[490882](#)]

Helsinki Businessmen Study, 1985:

Miettinen TA, Huttunen JK, Naukkarinen V, Strandberg T, Mattila S, Kumlin T, Sarna S Multifactorial primary prevention of cardiovascular diseases in middle-aged men. Risk factor changes, incidence, and mortality. *JAMA* 1985;254:2097-102 [[4046137](#)]

Strandberg TE, Salomaa VV, Naukkarinen VA, Vanhanen HT, Sarna SJ, Miettinen TA Long-term mortality after 5-year multifactorial primary prevention of cardiovascular diseases in middle-aged men. *JAMA* 1991;266:1225-9 [[1870247](#)]

Strandberg TE, Salomaa VV, Vanhanen HT, Naukkarinen VA, Sarna SJ, Miettinen TA Mortality in participants and non-participants of a multifactorial prevention study of cardiovascular diseases: a 28 year follow up of the Helsinki Businessmen Study. *Br Heart J* 1995;74:449-54 [[7488463](#)]

Johns Hopkins, 1983:

Morisky DE, Levine DM, Green LW, Shapiro S, Russell RP, Smith CR Five-year blood pressure control and mortality following health education for hypertensive patients. *Am J Public Health* 1983;73:153-62 [[6849473](#)]

Meland, 1997:

Meland E, Laerum E, Ulvik RJ Effectiveness of two preventive interventions for coronary heart disease in primary care. *Scand J Prim Health Care* 1997;15:57-64 [[9101627](#)]

Meland E, Maeland JG, Laerum E The importance of self-efficacy in cardiovascular risk factor change. *Scand J Public Health* 1999;27:11-7 [[10847665](#)]

MRFIT, 1982:

Mortality rates after 10.5 years for participants in the Multiple Risk Factor Intervention Trial. Findings related to a priori hypotheses of the trial. The Multiple Risk Factor Intervention Trial Research Group. *JAMA* 1990;263:1795-801 [[2179590](#)]

Multiple risk factor intervention trial. Risk factor changes and mortality results. Multiple Risk Factor Intervention Trial Research Group. *JAMA* 1982;248:1465-77 [[7050440](#)]

Oslo, 1981:

Anderssen S, Holme I, Urdal P, Hjermmann I Diet and exercise intervention have favourable effects on blood pressure in mild hypertensives: the Oslo Diet and Exercise Study (ODES). *Blood Press* 1995;4:343-9 [[8746601](#)]

Hjermmann I, Holme I, Leren P Oslo Study Diet and Antismoking Trial. Results after 102 months. *Am J Med* 1986;80:7-11 [[3511692](#)]

Hjermmann I, Velve Byre K, Holme I, Leren P Effect of diet and smoking intervention on the incidence of coronary heart disease. Report from the Oslo Study Group of a randomised trial in healthy men. *Lancet* 1981;2:1303-10 [[6118715](#)]

Holme I, Hjermmann I, Helgeland A, Leren P The Oslo Study: diet and antismoking advice. Additional results from a 5-year primary preventive trial in middle-aged men. *Prev Med* 1985;14:279-92 [[3903733](#)]

OXCHECK, 1994:

Effectiveness of health checks conducted by nurses in primary care: results of the OXCHECK study after one year. Imperial Cancer Research Fund OXCHECK Study Group. *BMJ* 1994;308:308-12 [[8124120](#)]

WHO Factories, 1982:

European collaborative trial of multifactorial prevention of coronary heart disease: final report on the 6-year results. World Health Organisation European Collaborative Group. Lancet 1986;1:869-72 [2870351]

Multifactorial trial in the prevention of coronary heart disease: 2. Risk factor changes at two and four years. Eur Heart J 1982;3:184-90 [7084265]

4 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.