

Clinical trials of myocardial revascularization for coronary artery disease in multivessel disease

TrialResults-center www.trialresultscenter.org

1 PCI

Trial	Treatments	Patients	Trials design and methods
balloon angioplasty vs CABG			
EAST , 1994 [NCT00000465] n=198/194 follow-up: 3 y	transluminal coronary angioplasty versus coronary-artery bypass grafting	patients with multivessels coronary artery disease	open USA
GABI , 1994 n=182/177 follow-up: 1 y	Percutaneous transluminal coronary angioplasty versus coronary-artery bypass grafting	patients with symptomatic multivessel coronary disease	open Germany
BARI , 1996 [NCT00000462] n=915/914 follow-up: 5.4 y	PTCA versus CABG	Patients with multivessel disease	open USA, Canada
RITA , 1993 n=510/501 follow-up: 2.5 y (6.5y)	percutaneous transluminal coronary angioplasty versus coronary artery bypass surgery	patients with one, two, or three diseased coronary arteries	open UK
ERACI , 1992 n=63/64 follow-up: 3.8 y	Percutaneous transluminal coronary angioplasty versus coronary artery bypass grafting	patients with multivessel disease and lesions suitable for either form of therapy	open Argentina
Toulouse , 1992 n=76/76 follow-up: 2.8 y	PTCA versus CABG	patients with multivessels coronary artery disease	open France
CABRI , 1995 n=541/513 follow-up: 1 y	percutaneous transluminal coronary angioplasty versus coronary artery bypass grafting	patients with symptomatic multivessel coronary disease	open Europe
PCI vs CABG			
AWESOME , 2001 n=222/232 follow-up: 5 years	percutaneous coronary intervention versus coronary artery bypass graft	high-risk patients with medically refractory ischemia	Parallel groups open US (Veterans Affairs Medical Centers)
PCI with drug-eluting stents vs CABG			

continued...

Trial	Treatments	Patients	Trials design and methods
FREEDOM , 2012 [NCT00086450] n=953/947 follow-up: 3.8 yrs (median)	percutaneous coronary stenting versus CABG	patients with diabetes and multivessel coronary artery disease	Parallel groups open international
stent vs CABG			
ARTS , 2001 n=600/605 follow-up: 1 year	Palmaz-Schatz Crown/Cross flex (Cordis) versus Conventional CABG	Multi vessel disease with 2 or more de novo lesion in different major arteries Total occlusion <1month	parallel group open International
CARDia (PCI) , 2008 [ISRCTN19872154] n=256/254 follow-up: 1 y	PCI plus stenting (and routine abciximab) versus CABG	Patients with diabetes and symptomatic multivessel coronary artery disease or complex single-vessel disease.	Parallel groups open UK, Ireland
ERACI II , 2003 n=225/225 follow-up: 30d, 1year	Gianturco Robin II (Cook) Primary device versus Conventional CABG	multi vessel disease Angina CSS III-IV; no angina but large area of heart at risk; unstable =1 vessel to be treated Lesion>3.0mm	parallel group open Argentinad
MASS II , 2007 n=205/203 follow-up: 5y (1y)	PCI (73% stent) versus CABG	patients with multivessel coronary artery disease with stable angina and preserved ventricular function	Parallel groups open South America
Myoprotect , 2004 n=23/21 follow-up: 1 year	percutaneous transluminal coronary angioplasty/stent versus CABG	patients with symptomatic main-stem and main-stem-equivalent lesions with substantially increased risk for bypass surgery	Parallel groups open Europe
SOS , 2002 [NCT00475449] n=488/500 follow-up: 3 years	Stent versus CABG	multiple vessel disease Symptomatic 1 or more vessel suitable for stenting	parallel group open Canada, United Kingdom, Europe
stent vs OPCAB			
OCTOSTENT , 2003 [NCT00975858] n=138/142 follow-up: 1 year	Stent type not reported versus off-pump coronary artery bypass	multi or single vessel disease Moderate LV function CABG or stenting to be considered feasible	Parallel groups open Europe

References

EAST, 1994:

King SB 3rd, Lembo NJ, Weintraub WS, Kosinski AS, Barnhart HX, Kutner MH, Alazraki NP, Guyton RA, Zhao XQ A randomized trial comparing coronary angioplasty with coronary bypass surgery. Emory Angioplasty versus Surgery Trial (EAST) N Engl J Med 1994 Oct 20;331:1044-50 [8090163]

GABI, 1994:

Hamm CW, Reimers J, Ischinger T, Rupprecht HJ, Berger J, Bleifeld W A randomized study of coronary angioplasty compared with bypass surgery in patients with symptomatic multivessel coronary disease. German Angioplasty Bypass Surgery Investigation (GABI) N Engl J Med 1994 Oct 20;331:1037-43 [8090162]

BARI, 1996:

Comparison of coronary bypass surgery with angioplasty in patients with multivessel disease. The Bypass Angioplasty Revascularization Investigation (BARI) Investigators. N

Engl J Med 1996 Jul 25;335:217-25 [8657237]

RITA, 1993:

Coronary angioplasty versus coronary artery bypass surgery: the Randomized Intervention Treatment of Angina (RITA) trial. Lancet 1993 Mar 6;341:573-80 [8094826]

ERACI, 1992:

Rodriguez A, Bouillon F, Perez-Balino N, Paviotti C, Liprandi MI, Palacios IF Argentine randomized trial of percutaneous transluminal coronary angioplasty versus coronary artery bypass surgery in multivessel disease (ERACI): in-hospital results and 1-year follow-up. ERACI Group. J Am Coll Cardiol 1993 Oct;22:1060-7 [8409041]

Toulouse, 1992:

CABRI, 1995:

First-year results of CABRI (Coronary Angioplasty versus Bypass Revascularisation Investigation). CABRI Trial Participants. Lancet 1995 Nov 4;346:1179-84 [7475656]

Martuscelli E, Clementi F, Gallagher MM, D'Eliseo A, Chiricolo G, Nigri A, Marino B, Romeo F Revascularization strategy in patients with multivessel disease and a major vessel chronically occluded; data from the CABRI trial. Eur J Cardiothorac Surg 2008;33:4-8 [17988889]

AWESOME, 2001:

Morrison DA, Sethi G, Sacks J, Henderson W, Grover F, Sedlis S, Esposito R, Ramanathan K, Weiman D, Saucedo J, Antakli T, Paramesh V, Pett S, Vernon S, Birjiniuk V, Welt F, Krucoff M, Wolfe W, Lucke JC, Mediratta S, Booth D, Barbieri C, Lewis D Percutaneous coronary intervention versus coronary artery bypass graft surgery for patients with medically refractory myocardial ischemia and risk factors for adverse outcomes with bypass: a multicenter, randomized trial. Investigators of the Department of Veterans Affairs Cooperative Study 385, the Angina With Extremely Serious Operative Mortality Evaluation (AWESOME). J Am Coll Cardiol 2001;38:143-9 [11451264]

Morrison DA, Sethi G, Sacks J, Grover F, Sedlis S, Esposito R, Ramanathan KB, Weiman D, Krucoff M, Duhaylongsod F, Raya T, Pett S, Vernon S, Birjiniuk V, Booth D, Robinson C, Talley JD, Antakli T, Murphy E, Floten H, Curcovic V, Lucke JC, Lewis D, Barbier A multicenter, randomized trial of percutaneous coronary intervention versus bypass surgery in high-risk unstable angina patients. The AWESOME (Veterans Affairs Cooperative Study 385, angina with extremely serious operative mortality evaluation) investigators from the Cooperative Studies Program of the Department of Veterans Affairs. Control Clin Trials 1999;20:601-19 [10588300]

Morrison DA, Sethi G, Sacks J, Henderson WG, Grover F, Sedlis S, Esposito R Percutaneous coronary intervention versus repeat bypass surgery for patients with medically refractory myocardial ischemia: AWESOME randomized trial and registry experience with post-CABG patients. J Am Coll Cardiol 2002;40:1951-4 [12475454]

Ramanathan KB, Weiman DS, Sacks J, Morrison DA, Sedlis S, Sethi G, Henderson WG Percutaneous intervention versus coronary bypass surgery for patients older than 70 years of age with high-risk unstable angina. Ann Thorac Surg 2005;80:1340-6 [16181866] 10.1016/j.athoracsur.2005.03.057

Rumsfeld JS, Magid DJ, Plomondon ME, Sacks J, Henderson W, Hlatky M, Sethi G, Morrison DA Health-related quality of life after percutaneous coronary intervention versus coronary bypass surgery in high-risk patients with medically refractory ischemia. J Am Coll Cardiol 2003;41:1732-8 [12767656]

Sedlis SP, Morrison DA, Lorin JD, Esposito R, Sethi G, Sacks J, Henderson W, Grover F, Ramanathan KB, Weiman D, Saucedo J, Antakli T, Paramesh V, Pett S, Vernon S, Birjiniuk V, Welt F, Krucoff M, Wolfe W, Lucke JC, Mediratta S, Booth D, Murphy E, Ward H, Percutaneous coronary intervention versus coronary bypass graft surgery for diabetic patients with unstable angina and risk factors for adverse outcomes with bypass: outcome of diabetic patients in the AWESOME randomized trial and registry. J Am Coll Cardiol 2002;40:1555-66 [12427406]

Sedlis SP, Ramanathan KB, Morrison DA, Sethi G, Sacks J, Henderson W Outcome of percutaneous coronary intervention versus coronary bypass grafting for patients with low left ventricular ejection fractions, unstable angina pectoris, and risk factors for adverse outcomes with bypass (the AWESOME Randomized Trial and Registry). Am J Cardiol 2004;94:118-20 [15219521] 10.1016/j.amjcard.2004.03.041

Stroupe KT, Morrison DA, Hlatky MA, Barnett PG, Cao L, Lyttle C, Hynes DM, Henderson WG Cost-effectiveness of coronary artery bypass grafts versus percutaneous coronary intervention for revascularization of high-risk patients. Circulation 2006;114:1251-7 [16966588] 10.1161/CIRCULATIONAHA.105.570838

FREEDOM, 2012:

Farkouh ME, Domanski M, Sleeper LA, Siami FS, Dangas G, Mack M, Yang M, Cohen DJ, Rosenberg Y, Solomon SD, Desai AS, Gersh BJ, Magnuson EA, Lansky A, Boineau R, Weinberger J, Ramanathan K, Sousa JE, Rankin J, Bhargava B, Buse J, Hueb W, Smith CR, Muratov Strategies for Multivessel Revascularization in Patients with Diabetes. N Engl J Med 2012 Nov 4; [23121323] 10.1056/NEJMoa1211585

ARTS, 2001:

Abizaid A, Costa MA, Centemero M, Abizaid AS, Legrand VM, Limet RV, Schuler G, Mohr FW, Lindeboom W, Sousa AG, Sousa JE, van Hout B, Hugenholtz PG, Unger F, Serruys PW Clinical and economic impact of diabetes mellitus on percutaneous and surgical treatment of multivessel coronary disease patients: insights from the Arterial Revascularization Therapy Study (ARTS) trial. *Circulation* 2001;104:533-8 [[11479249](#)]

de Feyter PJ, Serruys PW, Unger F, Beyar R, de Valk V, Milo S, Simon R, Regensburger D, Crean PA, McGovern E, van den Heuvel P, van Cauwelaert C, Penn I, Tyers GF, Lindeboom W Bypass surgery versus stenting for the treatment of multivessel disease in patients with unstable angina compared with stable angina. *Circulation* 2002;105:2367-72 [[12021222](#)]

Legrand VM, Serruys PW, Unger F, van Hout BA, Vrolix MC, Fransen GM, Nielsen TT, Paulsen PK, Gomes RS, de Queiroz e Melo JM, Neves JP, Lindeboom W, Backx B Three-year outcome after coronary stenting versus bypass surgery for the treatment of multivessel disease. *Circulation* 2004;109:1114-20 [[14993134](#)]

Serruys PW, Unger F, Sousa JE, Jatene A, Bonnier HJ, Schnberger JP, Buller N, Bonser R, van den Brand MJ, van Herwerden LA, Morel MA, van Hout BA Comparison of coronary-artery bypass surgery and stenting for the treatment of multivessel disease. *N Engl J Med* 2001;344:1117-24 [[11297702](#)]

Unger F, Serruys PW, Yacoub MH, Ilsley C, Paulsen PK, Nielsen TT, Eysmann L, Kiemeneij F Revascularization in multivessel disease: comparison between two-year outcomes of coronary bypass surgery and stenting. *J Thorac Cardiovasc Surg* 2003;125:809-20 [[12698143](#)]

CARDia (PCI), 2008:

Kapur A, Hall RJ, Malik IS, Qureshi AC, Butts J, de Belder M, Baumbach A, Angelini G, de Belder A, Oldroyd KG, Flather M, Roughton M, Nihoyannopoulos P, Bagger JP, Morgan K, Beatt KJ Randomized comparison of percutaneous coronary intervention with coronary artery bypass grafting in diabetic patients. 1-year results of the CARDia (Coronary Artery Revascularization in Diabetes) trial. *J Am Coll Cardiol* 2010 Feb 2;55:432-40 [[20117456](#)] [10.1016/j.jacc.2009.10.014](#)

ERACI II, 2003:

Rodriguez A, Bernardi V, Navia J, Baldi J, Grinfeld L, Martinez J, Vogel D, Grinfeld R, Delacasa A, Garrido M, Oliveri R, Mele E, Palacios I, O'Neill W Argentine Randomized Study: Coronary Angioplasty with Stenting versus Coronary Bypass Surgery in patients with Multiple-Vessel Disease (ERACI II): 30-day and one-year follow-up results. ERACI II Investigators. *J Am Coll Cardiol* 2001;37:51-8 [[11153772](#)]

Rodriguez A, Rodriguez Alemparte M, Baldi J, Navia J, Delacasa A, Vogel D, Oliveri R, Fernandez Pereira C, Bernardi V, O'Neill W, Palacios IF Coronary stenting versus coronary bypass surgery in patients with multiple vessel disease and significant proximal LAD stenosis: results from the ERACI II study. *Heart* 2003;89:184-8 [[12527674](#)]

MASS II, 2007:

Hueb W, Lopes NH, Gersh BJ, Soares P, Machado LA, Jatene FB, Oliveira SA, Ramires JA Five-year follow-up of the Medicine, Angioplasty, or Surgery Study (MASS II): a randomized controlled clinical trial of 3 therapeutic strategies for multivessel coronary artery disease. *Circulation* 2007 Mar 6;115:1082-9 [[17339566](#)]

Hueb W, Lopes N, Gersh BJ, Soares PR, Ribeiro EE, Pereira AC, Favarato D, Rocha AS, Hueb AC, Ramires JA Ten-year follow-up survival of the Medicine, Angioplasty, or Surgery Study (MASS II): a randomized controlled clinical trial of 3 therapeutic strategies for multivessel coronary artery disease. *Circulation* 2010;122:949-57 [[20733102](#)] [10.1161/CIRCULATIONAHA.109.911669](#)

Myoprotect, 2004:

Pohl T, Giehl W, Reichart B, Kupatt C, Raake P, Paul S, Reichensperner H, Steinbeck G, Boekstegers P Retroinfusion-supported stenting in high-risk patients for percutaneous intervention and bypass surgery: results of the prospective randomized myoprotect I study. *Catheter Cardiovasc Interv* 2004;62:323-30 [[15224298](#)] [10.1002/ccd.20060](#)

SOS, 2002:

Coronary artery bypass surgery versus percutaneous coronary intervention with stent implantation in patients with multivessel coronary artery disease (the Stent or Surgery trial): a randomised controlled trial. *Lancet* 2002;360:965-70 [[12383664](#)]

Stables RH Design of the 'Stent or Surgery' trial (SoS): a randomized controlled trial to compare coronary artery bypass grafting with percutaneous transluminal coronary angioplasty and primary stent implantation in patients with multi-vessel coronary artery disease. *Semin Interv Cardiol* 1999;4:201-7 [[10738353](#)]

Zhang Z, Mahoney EM, Stables RH, Booth J, Nugara F, Spertus JA, Weintraub WS Disease-specific health status after stent-assisted percutaneous coronary intervention and coronary artery bypass surgery: one-year results from the Stent or Surgery trial. *Circulation* 2003;108:1694-700 [[12975252](#)]

Zhang Z, Weintraub WS, Mahoney EM, Spertus JA, Booth J, Nugara F, Stables RH, Vaccarino V Relative benefit of coronary artery bypass grafting versus stent-assisted percutaneous coronary intervention for angina pectoris and multivessel coronary disease in women versus men (one-year results from the Stent or Surgery trial). *Am J Cardiol* 2004;93:404-9 [[14969611](#)]

Booth J, Clayton T, Pepper J, Nugara F, Flather M, Sigwart U, Stables RH Randomized, controlled trial of coronary artery bypass surgery versus percutaneous coronary intervention in patients with multivessel coronary artery disease: six-year follow-up from the Stent or Surgery Trial (SoS). *Circulation* 2008;118:381-8 [[18606919](#)]

Booth J, Clayton T, Pepper J, Nugara F, Flather M, Sigwart U, Stables RH Randomized, controlled trial of coronary artery bypass surgery versus percutaneous coronary intervention in patients with multivessel coronary artery disease: six-year follow-up from the Stent or Surgery Trial (SoS). *Circulation* 2008 Jul 22;118:381-8 [[18606919](#)]

OCTOSTENT, 2003:

Eefting F, Nathoe H, van Dijk D, Jansen E, Lahpor J, Stella P, Suyker W, Diephuis J, Suryapranata H, Ernst S, Borst C, Buskens E, Grobbee D, de Jaegere P Randomized comparison between stenting and off-pump bypass surgery in patients referred for angioplasty. *Circulation* 2003;108:2870-6 [[14656913](#)]

van Dijk D, Nierich AP, Eefting FD, Buskens E, Nathoe HM, Jansen EW, Borst C, Knape JT, Brede JJ, Robles de Medina EO, Grobbee DE, Diephuis JC, de Jaegere PP The Octopus Study: rationale and design of two randomized trials on medical effectiveness, safety, and cost-effectiveness of bypass surgery on the beating heart. *Control Clin Trials* 2000;21:595-609 [[11146152](#)]

2 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.