

Clinical trials of antibiotics for stable angina in all type of patient

TrialResults-center www.trialresultscenter.org

1 antibiotics

Trial	Treatments	Patients	Trials design and methods
Azithromycin vs placebo			
Gupta et al , 1997 n=43/41 follow-up: 65279;18mo	65279;Azithromycin 500 mg/d for 3 d (28 received 1 course, 12 received 2 courses 3 mo apart) versus placebo	Male patients at least 6 mo from documented MI and with titers to Chlamydia pneumoniae $\geq 1:64$	Parallel groups double blind
ACADEMIC , 1999 n=150/152 follow-up: 2y	Azithromycin 500 mg/d for 3 d then 500 mg/wk for 3 mo versus placebo	Patients with CAD and C pneumoniae titers of $\geq 1:16$. Patients were at least 5 d from an MI	Parallel groups double blind
STAMINA (Azithromycin) , 2002 n=111/107 follow-up: 1y	Azithromycin 500 mg/d for 3 d plus omeprazole 20 mg 2/d for 1 wk plus metronidazole 400 mg 2/d for 1 wk versus placebo	Patients with ACS	Parallel groups double blind England
AZACS , 2003 n=2004/2008 follow-up: 6mo	Azithromycin 500 mg on day 1 followed by 250 mg/d for 4d versus placebo	Patients with ACS	Parallel groups double blind
WIZARD , 2003 n=3879/3868 follow-up: 14mo	Azithromycin 600 mg/d for 3 d then 1/wk for 11 wk versus placebo	Patients with a history of MI of more than 6 weeks before and with C pneumoniae titers of $\geq 1:16$	Parallel groups double blind North America, Europe, Argentina, India
ACES , 2005 [NCT00000617] n=2004/2008 follow-up: 4y	Azithromycin 600 mg/wk for 1 y versus placebo	Patients with stable CAD	Parallel groups double blind US
clarithromycin vs placebo			
CLARIFY , 2001 n=74/74 follow-up: 1y	Clarithromycin 500 mg/d for 85 d versus placebo	Patients with ACS	Parallel groups double blind
CLARICOR , 2006 [NCT00121550] n=2172/2201 follow-up: 3 years	clarithromycin 500 mg/day versus placebo	patients with adischarge diagnosis of myocardial infarction or angina pectoris	Parallel groups double blind Denmark
Gatifloxacin vs placebo			

continued...

Trial	Treatments	Patients	Trials design and methods
PROVE-IT , 2005 n=2076/2086 follow-up: 24mo	Gatifloxacin 400 mg/d for 10 d/mo for 2y versus placebo	Patients hospitalized with ACS in the preceding 10 d	Parallel groups double blind
Roxithromycin vs placebo			
ROXIS , 1999 n=102/100 follow-up: 6mo	Roxithromycin 150 mg 2/d for 30 d versus placebo	Patients with documented history of CAD and ACS	Parallel groups double blind
Leowattana et al , 2001 n=40/20 follow-up: 3mo	Roxithromycin 150 mg/d for 30 d versus placebo	Patients with ACS	Parallel groups
ANTIBIO , 2003 n=433/437 follow-up: 1y	Roxithromycin 300 mg/d for 6 wk versus placebo	Patients with unstable angina or MI	Parallel groups double blind

References

Gupta et al, 1997:

Gupta S, Leatham EW, Carrington D, Mendall MA, Kaski JC, Camm AJ Elevated Chlamydia pneumoniae antibodies, cardiovascular events, and azithromycin in male survivors of myocardial infarction. *Circulation* 1997 Jul 15;96:404-7 [[9244203](#)]

ACADEMIC, 1999:

Muhlestein JB, Anderson JL, Carlquist JF, Salunkhe K, Horne BD, Pearson RR, Bunch TJ, Allen A, Trehan S, Nielson C Randomized secondary prevention trial of azithromycin in patients with coronary artery disease: primary clinical results of the ACADEMIC study. *Circulation* 2000 Oct 10;102:1755-60 [[11023928](#)]

STAMINA (Azithromycin), 2002:

Stone AF, Mendall MA, Kaski JC, Edger TM, Risley P, Poloniecki J, Camm AJ, Northfield TC Effect of treatment for Chlamydia pneumoniae and Helicobacter pylori on markers of inflammation and cardiac events in patients with acute coronary syndromes: South Thames Trial of Antibiotics in Myocardial Infarction and Unstable Angina (STAMINA). *Circulation* 2002 Sep 3;106:1219-23 [[12208796](#)]

AZACS, 2003:

Cercek B, Shah PK, Noc M, Zahger D, Zeymer U, Matetzky S, Maurer G, Mahrer P Effect of short-term treatment with azithromycin on recurrent ischaemic events in patients with acute coronary syndrome in the Azithromycin in Acute Coronary Syndrome (AZACS) trial: a randomised controlled trial. *Lancet* 2003 Mar 8;361:809-13 [[12642046](#)]

WIZARD, 2003:

O'Connor CM, Dunne MW, Pfeffer MA, Muhlestein JB, Yao L, Gupta S, Benner RJ, Fisher MR, Cook TD Azithromycin for the secondary prevention of coronary heart disease events: the WIZARD study: a randomized controlled trial. *JAMA* 2003 Sep 17;290:1459-66 [[13129985](#)]

ACES, 2005:

Grayston JT, Kronmal RA, Jackson LA, Parisi AF, Muhlestein JB, Cohen JD, Rogers WJ, Crouse JR, Borrowdale SL, Schron E, Knirsch C Azithromycin for the secondary prevention of coronary events. *N Engl J Med* 2005 Apr 21;352:1637-45 [[15843666](#)]

CLARIFY, 2001:

Sinialo J, Mattila K, Valtonen V, Anttonen O, Juvonen J, Melin J, Vuorinen-Markkola H, Nieminen MS Effect of 3 months of antimicrobial treatment with clarithromycin in acute non-q-wave coronary syndrome. *Circulation* 2002 Apr 2;105:1555-60 [[11927522](#)]

CLARICOR, 2006:

Jespersen CM, Als-Nielsen B, Damgaard M, Hansen JF, Hansen S, Hel OH, Hildebrandt P, Hilden J, Jensen GB, Kastrup J, Kolmos HJ, Kjeller E, Lind I, Nielsen H, Petersen L, Gluud C Randomised placebo controlled multicentre trial to assess short term clarithromycin for patients with stable coronary heart disease: CLARICOR trial. *BMJ* 2006 Jan 7;332:22-7 [[16339220](#)]

PROVE-IT, 2005:

Cannon CP, Braunwald E, McCabe CH, Grayston JT, Muhlestein B, Giugliano RP, Cairns R, Skene AM Antibiotic treatment of Chlamydia pneumoniae after acute coronary syndrome. N Engl J Med 2005 Apr 21;352:1646-54 [[15843667](#)]

ROXIS, 1999:

Gurfinkel E, Bozovich G, Beck E, Testa E, Livellara B, Mautner B Treatment with the antibiotic roxithromycin in patients with acute non-Q-wave coronary syndromes. The final report of the ROXIS Study. Eur Heart J 1999 Jan;20:121-7 [[10099908](#)]

Leowattana et al, 2001:**ANTIBIO, 2003:**

2 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.