

# Clinical trials of intensive blood pressure control for hypertension in diabetic patients

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 Treatment blood pressure target

Trial	Treatments	Patients	Trials design and methods
<b>more intensive blood pressure lowering strategie vs less intensive blood pressure lowering strategie</b>			
<b>ABCD target (H) , 2000</b> n=237/233 follow-up: 5 year	intensive treatment with a diastolic blood pressure goal of 75 mmHg versus moderate treatment with a diastolic blood pressure goal of 80-89 mmHg	diabetes patients with DBP $\geq$ 90 mmHg	Parallel groups open
<b>ABCD target (N) , 2002</b> n=237/243 follow-up:	intensive treatment (diastolic blood pressure decrease of 10 mmHg below baseline DBP) versus moderate treatment (diastolic blood pressure goal of 80-89 mmHg)	diabetes patients with diastolic blood pressure between 80 and 89mmHg	Parallel groups open
<b>ACCORD blood pressure , 2008</b> [NCT00000620] n=2362/2371 follow-up: 4.7y	intensive therapy, targeting a systolic pressure of less than 120 mm Hg versus standard therapy, targeting a systolic pressure of less than 140 mm Hg	patients with a median glycated hemoglobin level of 8.1% at high risk for cardiovascular events	Factorial plan open USA, Canada

## References

### ABCD target (H) , 2000:

Estacio RO, Jeffers BW, Gifford N, Schrier RW Effect of blood pressure control on diabetic microvascular complications in patients with hypertension and type 2 diabetes. Diabetes Care 2000;23 Suppl 2:B54-64 [[10860192](#)]

### ABCD target (N) , 2002:

Schrier RW, Estacio RO, Esler A, Mehler P Effects of aggressive blood pressure control in normotensive type 2 diabetic patients on albuminuria, retinopathy and strokes. Kidney Int 2002;61:1086-97 [[11849464](#)] [10.1046/j.1523-1755.2002.00213.x](#)

### ACCORD blood pressure, 2008:

Gerstein HC, Miller ME, Byington RP, Goff DC Jr, Bigger JT, Buse JB, Cushman WC, Genuth S, Ismail-Beigi F, Grimm RH Jr, Probstfield JL, Simons-Morton DG, Friedewald WT Effects of intensive glucose lowering in type 2 diabetes. N Engl J Med 2008;358:2545-59 [[18539917](#)] [10.1056/NEJMoa0802743](#)

Cushman WC, Grimm RH Jr, Cutler JA, Evans GW, Capes S, Corson MA, Sadler LS, Alderman MH, Peterson K, Bertoni A, Basile JN Rationale and design for the blood pressure intervention of the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial. Am J Cardiol 2007;99:44i-55i [[17599425](#)] [10.1016/j.amjcard.2007.03.005](#)

Effects of Intensive Blood-Pressure Control in Type 2 Diabetes Mellitus. N Engl J Med 2010 Mar 14;: [[20228401](#)] [10.1056/NEJMoa1001286](#)

## 2 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.