

Clinical trials of direct antithrombins for DVT prophylaxis in all type of patients

TrialResults-center www.trialresultscenter.org

1 oral direct thrombin inhibitor

Trial	Treatments	Patients	Trials design and methods
dabigatran 150mg vs enoxaparin			
RE-NOVATE (150mg) , 2007 [NCT00168818] n=1174/1162 follow-up: 28-35 days, median 33d	dabigatran etexilate 150 mg q.d. 28-35 days versus Enoxaparin 40 mg q.d. for 28-25 days	Total hip replacement	double blind Europe, Australia, South Africa
dabigatran 220mg vs enoxaparin			
RE-NOVATE 2 <i>unpublished</i> [NCT00657150] n=1010/1003 follow-up: 28-35 days (mean 32d)	dabigatran 220mg once daily for 28-35 Days versus enoxaparin 40mg subcutaneous once daily for 28-35 Days	patients undergoing total hip-replacement surgery	Parallel groups double-blind
RE-NOVATE (220mg) , 2007 [NCT00168818] n=1157/1162 follow-up: 28-35 days, median 33d	dabigatran etexilate 220 mg q.d. for 28-35 days versus Enoxaparin 40 mg q.d. for 23-35 days	Total hip replacement	Parallel groups double blind Europe, Australia, South Africa
dabigatran 150mg vs enoxaparin (europe regimen)			
RE-MODEL (150mg) , 2007 n=708/699 follow-up: 6-10 days, mean 8 days	dabigatran etexilate 150 mg q.d. for 6-10 days versus Enoxaparin 40 mg q.d. for 6-10 days	Total knee replacement	Parallel groups double blind Europe, Australia, South Africa
dabigatran 220mg vs enoxaparin (europe regimen)			
RE-MODEL (220mg) , 2007 n=694/699 follow-up: 6-10 days, mean 8 days	dabigatran etexilate 220 mg q.d. 6-10 days versus Enoxaparin 40 mg q.d. for 6-10 days	patients undergoing total knee replacement	double blind Europe, Australia, South Africa
dabigatran 150mg vs enoxaparin (US regimen)			
RE-MOBILIZE (150mg) , 2008 n=877/876 follow-up: 12-15 days, median 14d	dabigatran etexilate 150 mg q.d. for 12-15 days versus enoxaparin 30 mg SC BID after surgery for 12-15 days	Total knee replacement	double blind US, Canada, Mexico, UK
dabigatran 220mg vs enoxaparin (US regimen)			

continued...

Trial	Treatments	Patients	Trials design and methods
RE-MOBILIZE (220mg) , 2008 n=862/876 follow-up: 12-15 days, median 14d	dabigatran etexilate 220 mg for 12-15 days versus Enoxaparin 30mg SC BID after surgery for 12-15 days	Total knee replacement	Parallel groups double blind US, Canada, Mexico, UK

References

RE-NOVATE (150mg), 2007:

Eriksson BI, Dahl OE, Rosencher N, Kurth AA, van Dijk CN, Frostick SP, Prins MH, Hettiarachchi R, Hantel S, Schnee J, Bller HR Dabigatran etexilate versus enoxaparin for prevention of venous thromboembolism after total hip replacement: a randomised, double-blind, non-inferiority trial. *Lancet* 2007;370:949-56 [17869635]

RE-NOVATE 2, 0:

RE-NOVATE (220mg), 2007:

Eriksson BI, Dahl OE, Rosencher N, Kurth AA, van Dijk CN, Frostick SP, Prins MH, Hettiarachchi R, Hantel S, Schnee J, Bller HR Dabigatran etexilate versus enoxaparin for prevention of venous thromboembolism after total hip replacement: a randomised, double-blind, non-inferiority trial. *Lancet* 2007;370:949-56 [17869635]

RE-MODEL (150mg), 2007:

Eriksson BI, Dahl OE, Rosencher N, Kurth AA, van Dijk CN, Frostick SP, Klebo P, Christiansen AV, Hantel S, Hettiarachchi R, Schnee J, Bller HR Oral dabigatran etexilate vs. subcutaneous enoxaparin for the prevention of venous thromboembolism after total knee replacement: the RE-MODEL randomized trial. *J Thromb Haemost* 2007;5:2178-85 [17764540]
10.1111/j.1538-7836.2007.02748.x

RE-MODEL (220mg), 2007:

Eriksson BI, Dahl OE, Rosencher N, Kurth AA, van Dijk CN, Frostick SP, Klebo P, Christiansen AV, Hantel S, Hettiarachchi R, Schnee J, Bller HR Oral dabigatran etexilate vs. subcutaneous enoxaparin for the prevention of venous thromboembolism after total knee replacement: the RE-MODEL randomized trial. *J Thromb Haemost* 2007 Nov;5:2178-85 [17764540]

RE-MOBILIZE (150mg), 2008:

Ginsberg JS, Davidson BL, Comp PC, Francis CW, Friedman RJ, Huo MH, Lieberman JR, Muntz JE, Raskob GE, Clements ML, Hantel S, Schnee JM, Caprini JA Oral thrombin inhibitor dabigatran etexilate vs North American enoxaparin regimen for prevention of venous thromboembolism after knee arthroplasty surgery. *J Arthroplasty* 2009;24:1-9 [18534438]
10.1016/j.arth.2008.01.132

RE-MOBILIZE (220mg), 2008:

The Oral Thrombin Inhibitor Dabigatran Etexilate vs the North American Enoxaparin Regimen for the Prevention of Venous Thromboembolism after Knee Arthroplasty Surgery. *J Arthroplasty* 2008;: [18534438]

2 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.