

# Clinical trials of insulin secretagogues peptides (incretins) for diabetes type 2 in all type of patients

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 albiglutide

Trial	Treatments	Patients	Trials design and methods
<b>albiglutide vs placebo</b>			
Seino , 2009 [NCT00530309] n=32/8 follow-up: 4 weeks (+5 wk)	albiglutide 15 mg weekly, 30 mg weekly, 50 mg biweekly, and 100 mg monthly versus placebo	Japanese subjects with type 2 diabetes mellitus	Parallel groups single-blind Japan
<b>albiglutide biweekly vs placebo (add on MET)</b>			
Rosenstock (30 mg every two weeks) , 2009 [NCT00518115] n=32/50 follow-up: 16 weeks	albiglutide 30mg weekly versus placebo	patients with type 2 diabetes inadequately controlled with diet and exercise or metformin monotherapy	Parallel groups double blind US, Mexico, Chile, Dominical republic
<b>albiglutide weekly vs placebo (add on MET)</b>			
Rosenstock (30 mg weekly) , 2009 [NCT00518115] n=31/52 follow-up: 16 weeks	albiglutide 30mg weekly versus placebo	patients with type 2 diabetes inadequately controlled with diet and exercise or metformin monotherapy	Parallel groups double-blind US, Mexico, Chile, Dominical republic
<b>albiglutide vs liraglutide</b>			
HARMONY 7 (114179) <i>ongoing</i> [NCT01128894] n=NA follow-up:	weekly albiglutide subcutaneous injection versus liraglutide daily subcutaneous injection, starting at 0.6mg, then up-titrating to 1.2mg then 1.8mg in accordance with prescribing information	subjects with type 2 diabetes	open USA

## References

### Seino, 2009:

Seino Y, Nakajima H, Miyahara H, Kurita T, Bush MA, Yang F, Stewart MW, Safety, tolerability, pharmacokinetics and pharmacodynamics of albiglutide, a long-acting GLP-1-receptor agonist, in Japanese subjects with type 2 diabetes mellitus. *Curr Med Res Opin* 2009;25:3049-57. [[19863477](#)] [10.1185/03007990903372999](#)

### Rosenstock (30 mg every two weeks), 2009:

Rosenstock J, Reusch J, Bush M, Yang F, Stewart M, , Potential of albiglutide, a long-acting GLP-1 receptor agonist, in type 2 diabetes: a randomized controlled trial exploring weekly, biweekly, and monthly dosing. *Diabetes Care* 2009;32:1880-6. [[19592625](#)] [10.2337/dc09-0366](#)

### Rosenstock (30 mg weekly), 2009:

Rosenstock J, Reusch J, Bush M, Yang F, Stewart M, , Potential of albiglutide, a long-acting GLP-1 receptor agonist, in type 2 diabetes: a randomized controlled trial exploring weekly, biweekly, and monthly dosing. *Diabetes Care* 2009;32:1880-6. [[19592625](#)] [10.2337/dc09-0366](#)

### **HARMONY 7 (114179), 0:**

Pratley RE, Nauck MA, Barnett AH, Feinglos MN, Ovalle F, Harman-Boehm I, Ye J, Scott R, Johnson S, Stewart M, Rosenstock J Once-weekly albiglutide versus once-daily liraglutide in patients with type 2 diabetes inadequately controlled on oral drugs (HARMONY 7): a randomised, open-label, multicentre, non-inferiority phase 3 study. *Lancet Diabetes Endocrinol* 2014;2:289-97 [[24703047](#)] [10.1016/S2213-8587\(13\)70214-6](#)

## 2 glucagon-like peptide 1 receptor agonist

Trial	Treatments	Patients	Trials design and methods
<b>liraglutide vs placebo</b>			
<b>LEADER , 2016</b> [NCT01179048] n=4668/4672 follow-up: 3.8 years (median)	Maximum dose of 1.8 mg liraglutide, injected subcutaneously once daily versus placebo	subjects with type 2 diabetes	double-blind Africa, Asia, Europe, North and South America

## References

### **LEADER, 2016:**

Steinberg WM, Nauck MA, Zinman B, Daniels GH, Bergenstal RM, Mann JF, Steen Ravn L, Moses AC, Stockner M, Baeres FM, Marso SP, Buse JB LEADER 3-lipase and amylase activity in subjects with type 2 diabetes: baseline data from over 9000 subjects in the LEADER Trial. *Pancreas* 2014;43:1223-31 [[25275271](#)]

Petrie JR, Marso SP, Bain SC, Franek E, Jacob S, Masmiquel L, Leiter LA, Haluzik M, Satman I, Omar M, Shestakova M, Van Gaal L, Mann JF, Baeres FM, Zinman B, Poulter NR LEADER-4: blood pressure control in patients with type 2 diabetes and high cardiovascular risk: baseline data from the LEADER randomized trial. *J Hypertens* 2016;: [[26855018](#)]

Masmiquel L, Leiter LA, Vidal J, Bain S, Petrie J, Franek E, Raz I, Comlekci A, Jacob S, van Gaal L, Baeres FM, Marso SP, Eriksson M LEADER 5: prevalence and cardiometabolic impact of obesity in cardiovascular high-risk patients with type 2 diabetes mellitus: baseline global data from the LEADER trial. *Cardiovasc Diabetol* 2016;15:29 [[26864124](#)]

Marso SP, Daniels GH, Brown-Frandsen K, Kristensen P, Mann JF, Nauck MA, Nissen SE, Pocock S, Poulter NR, Ravn LS, Steinberg WM, Stockner M, Zinman B, Bergenstal RM, Buse JB Liraglutide and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med* 2016;: [[27295427](#)]

## 3 glucagon-like peptide analogs

Trial	Treatments	Patients	Trials design and methods
<b>liraglutide other doses vs</b>			
<b>NN2211-1333</b> n=NA follow-up:	liraglutide versus placebo	obese subjects with type 2 diabetes	
<b>exenatide vs glargine</b>			

continued...

Trial	Treatments	Patients	Trials design and methods
<a href="#">NCT00360334</a> [NCT00360334] n=118/116 follow-up:	-	-	
<b>exenatide 20g vs glibenclamide</b>			
<a href="#">Derosa , 2010</a> n=63/65 follow-up:	exenatide 10 microg twice a day versus glibenclamide 5 mg three times a day	patients with uncontrolled type 2 diabetes mellitus receiving therapy with metformin	
<b>exenatide vs placebo</b>			
<a href="#">Liutkus , 2010</a> n=111/54 follow-up:	exenatide twice-daily versus placebo	subjects suboptimally controlled with TZDs with or without metformin	
<b>exenatide 10g/d vs placebo</b>			
<a href="#">H8O-MC-GWBJ, 9698, 10g/d , 2008</a> n=NA follow-up: 24 weeks	exenatide twice daily 5 et 10 g for 24 weeks versus placebo	Drug-Naive Patients with Type 2 Diabetes and inadequate glycemic control through diet and exercise	Parallel groups double-blind 4 countries
<b>exenatide 20g/d vs placebo</b>			
<a href="#">Apovian , 2010</a> n=96/98 follow-up: 24 weeks	10 microg exenatide twice daily injection + lifestyle modification program versus placebo + lifestyle modification program	overweight or obese participants with type 2 diabetes treated with metformin and/or sulfonylurea	Parallel groups double-blind
<a href="#">H8O-MC-GWBJ, 9698, 20g/d , 2008 unpublished</a> n=78/78 follow-up: 24 weeks	exenatide twice daily 10 g for 24 weeks versus placebo	Drug-Naive Patients with Type 2 Diabetes and inadequate glycemic control through diet and exercise	Parallel groups double-blind 4 countries
<b>exenatide other doses vs placebo</b>			
<a href="#">Moretto (DOUBLONS avec druker) , 2008</a> [NCT00381342] n=155/78 follow-up: 24 weeks	Exenatide 1020 g daily versus Placebo	-	Parallel groups double blind United States, Puerto Rico, Romania, Russia, India
<a href="#">NCT00085969 unpublished</a> [NCT00085969] n=99 follow-up: 28 days	exenatide for 28 days versus placebo	subjects with type 2 diabetes mellitus	double-blind USA
<a href="#">Poon , 2005</a> [NCT00044694] n=NA follow-up: 28 days	exenatide at 2.5, 5.0, 7.5, or 10.0 microg administered b.i.d. for 28 days versus placebo	patients with type 2 diabetes	Parallel groups double-blind
<b>liraglutide other doses vs placebo</b>			
<a href="#">Harder , 2004</a> n=21/12 follow-up: 8 weeks	single daily subcutaneous dose of 0.6 mg liraglutide for 8 weeks versus placebo	obese subjects with type 2 diabetes	Parallel groups double-blind Denmark

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>Kaku 0.6mg , 2010</b> n=88/88 follow-up: 24 weeks	liraglutide 0.6 mg/day versus placebo	Japanese patients with type 2 diabetes	Parallel groups double-blind Japan
<b>Madsbad (vs placebo) , 2004</b> n=135/29 follow-up: 12 weeks	Liraglutide 0.045, 0.225, 0.45, 0.60, and 0.75 mg daily versus Placebo	Outpatients with type 2 diabetes	open UK, Scandinavia
<b>NN2211-1571 (Vilsbll) , 2007</b> [NCT00154401] n=123/40 follow-up: 14 weeks	liraglutide 0.65 mg, 1.25 mg or 1.9 mg for 14 weeks versus placebo	subjects with type 2 diabetes	Parallel groups double-blind Denmark, France, Slovakia, Netherlands
<b>Seino , 2008</b> [NCT00154414] n=180/46 follow-up: 14 weeks	Liraglutide 0.1, 0.3, 0.6 or 0.9 mg once daily for 14 weeks versus Placebo	Japanese subjects with type 2 diabetes	Parallel groups double blind Japan
<b>LIBRA ongoing</b> [NCT01270789] n=NA follow-up: 1 year	-	patients with T2DM	double-blind Canada
<b>NCT00978393 ongoing</b> [NCT00978393] n=NA follow-up:	High dose liraglutide treatment (3.0 mg) followed by low dose liraglutide treatment (1.8 mg) s.c. once daily versus placebo	non-diabetic obese volunteers	double-blind Netherlands
<b>NN2211-1799 ongoing</b> [NCT00620282] n=NA follow-up: 3 months	liraglutide Stepwise dose increase, s.c. injection, once daily versus placebo	subjects with type 2 diabetes who are on diet and lifestyle changes or treated with metformin alon	double-blind USA
<b>NN2211-1800 ongoing</b> [NCT00943501] n=NA follow-up:	-	children with type 2 diabetes	double-blind USA, Europe
<b>NN2211-3619 ongoing</b> [NCT01206101] n=NA follow-up:	-	Type 1 Diabetes Undergoing Islet Cell Transplantation	
<b>NN8022-1922 ongoing</b> [NCT01272232] n=NA follow-up: 56 weeks	Liraglutide 3.0 mg for subcutaneous (under the skin) injection once daily for 56 weeks in addition to subject's pre-trial background treatment versus placebo	overweight or obese subjects with type 2 diabetes	double-blind Africa, Asia, Europe and the United States of America

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>NN8022-1923</b> <i>ongoing</i> [NCT00781937] n=NA follow-up: 56 weeks	Liraglutide 3.0 mg per day administered in a 6.0 mg/mL, 3 mL FlexPen for subcutaneous (s.c.) injection, once daily versus placebo	obese non-diabetic subjects or overweight subjects who have medical problems such as hypertension or dyslipidaemia	double-blind North America
<b>taspoglutide vs placebo</b>			
<b>BC20750</b> <i>ongoing</i> [NCT00744926] n=NA follow-up:	taspoglutide 10mg sc once weekly, or taspoglutide 20mg sc once weekly (after 4 weeks of taspoglutide 10mg sc once weekly) versus placebo	patients with type 2 diabetes mellitus inadequately controlled with diet and exercise	double-blind USA
<b>BC20963</b> <i>ongoing</i> [NCT00744367] n=NA follow-up: 24 weeks	taspoglutide 10mg once weekly, taspoglutide 20 mg once weekly (after 4 weeks of taspoglutide 10 mg once weekly) versus placebo in addition to their continued stable metformin plus pioglitazone treatment	patients with type 2 diabetes mellitus inadequately controlled with metformin plus pioglitazone	double-blind USA
<b>BC21713 (vs placebo)</b> <i>ongoing</i> [NCT00754988] n=NA follow-up:	taspoglutide (10mg once weekly or 10mg once weekly for 4 weeks followed by 20mg once weekly), versus placebo or sitagliptin 100mg once daily in addition to their continued prestudy metformin treatment	patients with type 2 diabetes mellitus inadequately controlled with metformin	parallel groups double-blind USA
<b>BP21572</b> <i>ongoing</i> [NCT00809705] n=NA follow-up: 12 weeks	10mg Taspoglutide sc weekly for 12 weeks b)10mg Taspoglutide sc weekly for 4 weeks followed by 20mg Taspoglutide sc weekly for 8 weeks versus placebo	patients with type 2 diabetes	double-blind germany
<b>taspoglutide 10mg once weekly vs placebo</b>			
<b>Nauck 10 once weekly vs PBO , 2009</b> [NCT00423501] n=257/49 follow-up: 12 weeks	taspoglutide, either 5, 10, or 20 mg once weekly or 10 or 20 mg once every 2 weeks for 8 weeks versus placebo	patients with type 2 diabetes inadequately controlled with metformin	Parallel groups double-blind
<b>taspoglutide 20mg once every 2 weeks vs placebo</b>			
<b>Nauck 20 every 2 weeks vs PBO , 2009</b> n=NA	-	-	-
<b>taspoglutide 20mg once weekly vs placebo</b>			
<b>Nauck 20 once weekly vs PBO , 2009</b> n=NA	-	-	-
<b>exenatide 20g/d vs placebo (add on insulin)</b>			

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>Buse , 2011</b> [NCT00765817] n=138/123 follow-up: 30 weeks	twice-daily 10 g exenatide injections versus placebo (on top insulin glargine)	Adults with type 2 diabetes and an HbA1c level of 7.1% to 10.5% who were receiving insulin glargine alone or in combination with metformin or pioglitazone (or both agents)	Parallel groups double-blind Greece, Israel, Mexico, United Kingdom, USA
<b>exenatide other doses vs placebo (add on MER+/-SU)</b>			
<b>Fineman , 2003</b> n=109 follow-up: 28 days	exenatide 3 regimen (0.08 micro g/kg) for 28 days versus placebo	patients with type 2 diabetes treated with diet and a sulfonylurea and/or metformin	Parallel groups double-blind USA
<b>exenatide 10g/d vs placebo (add on MET)</b>			
<b>DeFronzo 10g/d , 2005</b> [NCT00039013] n=110/113 follow-up: 30 weeks	Exenatide 1020 g daily versus Placebo on-top of Metformin	patients with type 2 diabetes failing to achieve glycemic control with maximally effective metformin doses	Parallel groups double blind USA
<b>exenatide 20g/d vs placebo (add on MET)</b>			
<b>DeFronzo 20g/d , 2005</b> [NCT00039013] n=NA follow-up: 30 weeks	Exenatide 1020 g daily versus Placebo on-top of Metformin	patients with type 2 diabetes failing to achieve glycemic control with maximally effective metformin doses	Parallel groups double blind USA
<b>exenatide weekly vs placebo (add on MET)</b>			
<b>Kim , 2007</b> [NCT00103935] n=30/15 follow-up: 15 weeks	exenatide LAR 0.8 or 2 g daily versus Placebo on-top of metformin	subjects with type 2 diabetes suboptimally controlled with metformin and/or diet and exercise	Parallel groups double blind
<b>liraglutide 1.2mg vs placebo (add on MET)</b>			
<b>LEAD-2 (Nauck) (1.2mg vs placebo) , 2009</b> [NCT00318461] n=241/122 follow-up: 26 weeks	Liraglutide 1.2 mg daily versus Placebo on-top of Metformin	subjects previously treated with oral antidiabetes therapy	Parallel groups double blind 21 countries
<b>liraglutide 1.8mg vs placebo (add on MET)</b>			
<b>LEAD-2 (Nauck) (1.8mg vs placebo) , 2009</b> [NCT00318461] n=242/122 follow-up: 26 weeks	Liraglutide 1.8 mg daily versus Placebo on-top of Metformin	subjects previously treated with oral antidiabetes therapy	Parallel groups double blind 21 countries
<b>liraglutide other doses vs placebo (add on MET)</b>			
<b>NN2211-1796 unpublished</b> [NCT00614120] n=NA follow-up:	liraglutide added to metformin versus glimepiride added to metformin	-	China
<b>NCT01234649 ongoing</b> [NCT01234649] n=NA follow-up:	addition of liraglutide to metformin versus metformin alone	at-risk overweight/obese women with prior gestational diabetes mellitus	double-blind USA

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>tasoglutide vs placebo (add on MET)</b>			
Ratner (20mg once weekly) , 2010 [NCT00460941] n=97/32 follow-up: 8 weeks (+4wk)	tasoglutide s.c. 20mg once weekly for 8 weeks versus placebo s.c. once weekly on top metformin	subjects with Type 2 diabetes inadequately controlled on metformin alone	Parallel groups double-blind Australia, France, Germany, Mexico, Peru, USA
BC22092 ongoing [NCT00823992] n=NA follow-up:	tasoglutide (10mg sc once weekly for 4 weeks followed by 20mg once weekly) in addition to their prescribed, pre-existing metformin therapy versus placebo	obese patients with type 2 diabetes mellitus inadequately controlled with metformin monotherapy	parallel groups double-blind USA
<b>exenatide 20g/d vs placebo (add on MET+/-SU)</b>			
Gao , 2009 [NCT00324363] n=234/232 follow-up: 16 weeks	exenatide 5 mg then 10 mg twice-daily for 4 and 12 weeks versus placebo	Asian descent with type 2 diabetes and inadequate glycemic control taking metformin alone or Met and sulfonylureas	Parallel groups double-blind 4 countries
<b>exenatide 10g/d vs placebo (add on SU)</b>			
Buse 10g/d , 2004 [NCT00039026] n=125/123 follow-up: 30 weeks	Exenatide 5g twice daily versus Placebo on-top of SU	patients with type 2 diabetes failing maximally effective doses of a sulfonylurea as monotherapy	Parallel groups double blind (not adequate) US
<b>exenatide 20g/d vs placebo (add on SU)</b>			
Buse 20g/d , 2004 n=129/123 follow-up: 30 weeks	Exenatide 10g twice daily versus Placebo on-top of SU	patients with type 2 diabetes failing maximally effective doses of a sulfonylurea as monotherapy	double blind (not adequate) US
<b>liraglutide 1.2mg vs placebo (add on SU)</b>			
LEAD-1 SU (1.2 mg vs placebo) , 2009 [NCT00318422] n=228/115 follow-up: 26 weeks	Liraglutide 1.2 mg daily versus Placebo on-top of sulphonylureas	subjects with Type 2 diabetes	Parallel groups double-blind 21 countries
<b>liraglutide 1.8mg vs placebo (add on SU)</b>			
LEAD-1 SU (1.8 mg vs placebo) , 2009 [NCT00318422] n=234/114 follow-up: 26 weeks	Liraglutide 1.8 mg daily versus Placebo on-top of sulphonylureas	patients with type 2 diabetes	Parallel groups double-blind 21 countries
<b>liraglutide other doses vs placebo (add on SU)</b>			
NN2211-1701 ongoing [NCT00395746] n=NA follow-up: 24 weeks	liraglutide in combination with sulphonylurea versus placebo (add on to SU monotherapy)	subjects with type 2 diabetes	Parallel groups double-blind Japan

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>exenatide 10g/d vs placebo (add on SU+/-MET/TZD)</b>			
Kadowaki (trial 8683) , 2009 n=111/40 follow-up: 12 weeks	Exenatide 10g daily for 12 weeks versus Placebo on-top of sulphonylureas +/-metformin/thiazolidinediones	Japanese patients with type 2 diabetes suboptimally controlled despite therapeutic dose of sulfonylurea, SU+biguanide or SU+thiazolidinedione	Parallel groups open Japan
<b>exenatide 10g/d vs placebo (add on SU+MET)</b>			
Kendall 10g/d , 2005 [NCT00035984] n=245/247 follow-up: 30 weeks	Exenatide 5 g bid versus Placebo on-top of sulphonylureas+metformin	patients with type 2 diabetes unable to achieve glyceemic control with metformin-sulfonylurea combination therapy	Parallel groups double blind USA
<b>exenatide 20g/d vs placebo (add on SU+MET)</b>			
Kendall 20g/d , 2005 [NCT00035984] n=241/247 follow-up: 30 weeks	Exenatide 10 g bid versus Placebo on-top of sulphonylureas+metformin	patients with type 2 diabetes unable to achieve glyceemic control with metformin-sulfonylurea combination therapy	Parallel groups double blind USA
<b>liraglutide 1.8mg vs placebo (add on SU+MET)</b>			
LEAD-5 (vs placebo) , 2009 [NCT00331851] n=232/115 follow-up: 26 weeks	Liraglutide 1.8 mg daily versus Placebo on-top of sulphonylureas+metformin	adult patients with type 2 diabetes	Parallel groups double-blind 17 countries
<b>exenatide 20g/d vs placebo (add on TZD+/-MET)</b>			
Zinman 20g/j , 2007 [NCT00099320] n=121/112 follow-up: 16 weeks	Exenatide 20 g daily versus Placebo on-top of thiazolidinediones+/-metformin	patients with type 2 diabetes that was suboptimally controlled with TZD treatment (with or without metformin)	double blind Canada, Spain, and the United States
Zinman 20g/j A MODIFIER , 2007 n=121/112 follow-up: 16 weeks	exenatide Subcutaneous abdominal injections of 10 microg twice daily versus placebo	patients with type 2 diabetes that was suboptimally controlled with TZD treatment (with or without metformin)	Parallel groups double-blind Canada, Spain, and the United States
<b>liraglutide 1.2mg vs placebo (add on TZD+MET)</b>			
LEAD-4 (1.2mg) , 2009 [NCT00333151] n=178/177 follow-up: 26 weeks	Liraglutide 1.2 daily versus Placebo on-top of thiazolidinediones + metformin	patients with type 2 diabetes, A1C 711% (previous OAD monotherapy >=3 months) or 710% (previous OAD combination therapy >=3 months), and BMI 45 kg/m2	Parallel groups double-blind USA, Canada
<b>liraglutide 1.8mg vs placebo (add on TZD+MET)</b>			
LEAD-4 (1.8mg) , 2009 [NCT00333151] n=178/177 follow-up: 26 weeks	Liraglutide 1.8 daily versus Placebo on-top of thiazolidinediones + metformin	patients with type 2 diabetes, A1C 711% (previous OAD monotherapy >=3 months) or 710% (previous OAD combination therapy >=3 months), and BMI 45 kg/m2	double-blind USA, Canada
<b>liraglutide other doses vs placebo (on top SU)</b>			

continued...



<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>Kaku 0.9mg , 2010</b> n=88/88 follow-up: 24 weeks	liraglutide 0.9 mg/day versus placebo	Japanese patients with type 2 diabetes	double-blind Japan
<b>exenatide 10g/d vs placebo add on MET+/-TZD</b>			
<b>Gill , 2010</b> n=28/26 follow-up: 12 weeks	exenatide (5 microg for 4 weeks followed by 10 microg) for 12 weeks versus placebo	subjects with type 2 diabetes mellitus on metformin and/or a thiazolidinedione	Parallel groups double-blind
<b>tasoglutide vs placebo add on standard treatment</b>			
<b>NC25113 ongoing</b> [NCT01018173] n=NA follow-up:	tasoglutide subcutaneously (sc) 10mg weekly for 4 weeks followed by 20mg sc weekly in addition to background anti-hyperglycemic medication and standard of care treatment for cardiovascular disease versus placebo	-	parallel groups double-blind USA
<b>liraglutide other doses vs sitagliptin (add on MET)</b>			
<b>MK-0431-403 ongoing</b> [NCT01296412] n=NA follow-up:	Liraglutide + metformin versus Sitagliptin + metformin	patients with Type 2 Diabetes that is not adequately controlled with metformin alone	parallel groups open
<b>exenatide once monthly vs weekly exenatide</b>			
<b>phase 2 exenatide once monthly unpublished</b> n=121 follow-up: 20 weeks	exenatide once monthly at a low, medium or high dose, each administered once every four weeks, for a total of 20 weeks versus exenatide 2mg once weekly	adults with type 2 diabetes who were not achieving adequate glucose control using diet and exercise alone or with a stable regimen of metformin, pioglitazone, or both	Parallel groups open
<b>exenatide 20g/d vs BIAsp 30 daily</b>			
<b>Bergenstal (once daily) , 2009</b> n=NA follow-up: 24 weeks	exenatide(5 microg BID for 4 weeks and 10 microg BID thereafter) versus biphasic insulin aspart 70/30 (BIAsp 30) 30 QD (12 U before supper)	subjects with type 2 diabetes mellitus insulin naive, not achieving glycemic targets with metformin and sulfonylurea	Parallel groups open
<b>exenatide 20g/d vs BIAsp 30 twice daily</b>			
<b>Bergenstal (twice daily) , 2009</b> n=NA follow-up:	exenatide (5 microg BID for 4 weeks and 10 microg BID thereafter) versus biphasic insulin aspart 70/30 (BIAsp 30) 30 BID (12 U divided equally between pre-breakfast and pre-supper)	subjects with type 2 diabetes mellitus insulin naive, not achieving glycemic targets with metformin and sulfonylurea	Parallel groups open
<b>tasoglutide vs exenatide</b>			

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>BC21625</b> <i>ongoing</i> [NCT00717457] n=NA follow-up:	tasoglutide versus exenatide	patients with type 2 diabetes mellitus inadequately controlled with metformin, thiazolidinedione or a combination of both	parallel groups open USA
<b>exenatide before lunch and dinner vs exenatide before breakfast and dinner</b>			
<b>Exenatide Trial 10749</b> n=187/190 follow-up:	exenatide (10 g twice daily) administered subcutaneously before lunch and dinner versus exenatide (10 g twice daily) administered subcutaneously before breakfast and dinner	patients with type 2 Diabetes using oral antidiabetic therapy	Parallel groups open 2 countries
<b>liraglutide 1.8mg vs exenatide on top MET/SU/MET+SU</b>			
<b>LEAD-6 , 2009</b> [NCT00518882] n=233/231 follow-up: 26 weeks	liraglutide 1.8 mg once a day versus exenatide 10 microg twice a day	Adults with inadequately controlled type 2 diabetes on maximally tolerated doses of metformin, sulphonylurea, or both	Parallel groups open 15 countries
<b>exenatide other doses vs glargine</b>			
<b>HEELA (Davies) , 2009</b> n=NA follow-up:	exenatide 5C10 g bid versus insulin glargine o.d. (titrated to target fasting plasma glucose 5.6 mmol/l)	Patients (BMI>27 kg/m <sup>2</sup> ) with elevated cardiovascular risk and type 2 diabetes inadequately controlled on two or three oral antidiabetic drugs	
<b>liraglutide other doses vs glibenclamide</b>			
<b>Seino , 2010</b> [NCT00393718] n=272/139 follow-up: 24-week	liraglutide 0.9 mg once daily versus glibenclamide once or twice daily at a planned maximum dose of 2.5 mg/day, before or after meals	Japanese subjects with type 2 diabetes, inadequately controlled with diet therapy or oral antidiabetic drug monotherapy	Parallel groups double-blind Japan
<b>liraglutide 1.2mg vs glimepiride</b>			
<b>LEAD-3 mono 1.2mg</b> <b>(Garber) , 2009</b> [NCT00294723] n=251/248 follow-up: 52 weeks (104 weeks)	liraglutide 1.2 mg daily versus glimepiride 8 mg once daily	patients with early type 2 diabetes	Parallel groups double blind North America, Mexico
<b>liraglutide 1.8mg vs glimepiride</b>			
<b>LEAD-3 mono 1.8mg</b> <b>(Garber) , 2009</b> [NCT00294723] n=247/248 follow-up: 52 weeks (104 weeks)	liraglutide 1.8 mg daily versus glimepiride 8 mg once daily	subjects with type 2 diabetes	Parallel groups double blind North America, Mexico
<b>liraglutide other doses vs glimepiride</b>			
<b>Madsbad (vs Glimepiride) ,</b> <b>2004</b> n=135/26 follow-up: 12 weeks	Liraglutide 0.045, 0.225, 0.45, 0.60, and 0.75 mg daily versus Glimepiride	Outpatients with type 2 diabetes	Parallel groups open UK, Scandinavia

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>liraglutide 1.2mg vs glimepiride (add on MET)</b>			
LEAD-2 (Nauck) (1.2 mg vs glimepiride) , 2009 [NCT00318461] n=241/244 follow-up: 26 weeks	Liraglutide 1.2mg daily for 26 weeks versus Glimepiride on-top of Metformin	patients with type 3 diabetes previously treated with oral antidiabetes (OAD) therap	Parallel groups double blind 21 countries
<b>liraglutide 1.8mg vs glimepiride (add on MET)</b>			
LEAD-2 (Nauck) (1.8 mg vs glimepiride) , 2009 [NCT00318461] n=242/244 follow-up: 26 weeks	Liraglutide 1.8 mg daily for 26 weeks versus Glimepiride on-top of Metformin	patients with type 3 diabetes previously treated with oral antidiabetes (OAD) therap	Parallel groups double blind 21 countries
<b>exenatide 20g/d vs insulin (add on SU+MET)</b>			
Heine , 2005 n=282/267 follow-up: 26 weeks	Exenatide 20 g daily versus Insulin on-top of sulphonylureas+metformin	-	open
<b>exenatide 20g/d vs insulin (add on SU/MET)</b>			
Barnett , 2007 [NCT00099619] n=136/127 follow-up: 16 weeks	Exenatide 20 g daily versus Insulin	patients with type 2 diabetes	Cross over open Australia, Greece,Hungary, Italy, Mexico, and Poland
Davis , 2007 [NCT00099333] n=33/16 follow-up: 16 weeks	Exenatide 20 g daily versus Insulin on-top of sulphonylureas/metformin	patients with type 2 diabetes using insulin in combination with oral antidiabetes agents	Parallel groups open USA
<b>exenatide 20g/d vs insulin BIAsp twice daily add on SU+MET</b>			
Nauck , 2007 [NCT00082407] n=253/248 follow-up: 52 weeks	Exenatide 20 g daily versus Insulin on-top of sulphonylureas+metformin	patients with type 2 diabetes who were suboptimally controlled with sulfonyleurea and metformin	Parallel groups open 13 countries
<b>exenatide weekly vs insulin glargine</b>			
DURATION-3 (Diamant) , 2010 [NCT00641056] n=233/223 follow-up: 26 weeks	exenatide (2 mg, once-a-week injection) versus insulin glargine once-daily injection	adults with type 2 diabetes who had suboptimum glycaemic control despite use of maximum tolerated doses of blood-glucose-lowering drugs for 3 months or longer	Parallel groups open (blind analysis) USA, Puerto Rico, Europe, Russia, Australia, Korea, Taiwan, Mexico
<b>liraglutide other doses vs insulin glargine</b>			
EAGLE <i>ongoing</i> [NCT01117350] n=NA follow-up:	Liraglutide (6 mg/mL solution for injection in a 3-mL pre-filled pen (18mg)) versus Insulin Glargine (100 Units/mL solution for injection in a pre-filled SoloStar pen)	Type 2 diabetic patients failing lifestyle management and oral agents	open USA
<b>tasoglutide vs insulin glargine</b>			

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>ZC22565</b> <i>ongoing</i> [NCT01051011] n=NA follow-up:	tasoglutide 10mg subcutaneously (sc) weekly, or tasoglutide 10mg sc weekly for 4 weeks followed by 20mg sc weekly versus insulin glargine at an initial dose of 10 international units sc daily	insulin-naive patients with type 2 diabetes mellitus inadequately controlled on metformin and sulfonylurea combination therapy	parallel groups open China
<b>exenatide 20g/d vs insulin glargine (add on MET)</b>			
<b>Bunck , 2009</b> [NCT00097500] n=36/33 follow-up: 52 weeks	exenatide 10g bid versus insulin glargine	metformin-treated patients with type 2 diabetes	Parallel groups
<b>tasoglutide vs insulin glargine (add on MET)</b>			
<b>BC20965</b> <i>ongoing</i> [NCT00755287] n=NA follow-up: 2 years	tasoglutide (10 mg once weekly, or 10mg once weekly for 4 weeks followed by 20mg once weekly) versus insulin glargine (starting dose 10 IU/day) in addition to continued prestudy metformin treatment	patients with insulin-naive type 2 diabetes mellitus inadequately controlled with metformin and sulphonylurea combination therapy	open USA
<b>exenatide other doses vs insulin glargine (add on MET/SU)</b>			
<b>Trial 8078</b> n=NA follow-up:	exenatide versus Insulin Glargine	Patients with Type 2 Diabetes Using Metformin or Sulfonylurea for Whom Insulin Is the Next Appropriate Therapy	
<b>liraglutide 1.8mg vs insulin glargine (add on SU+MET)</b>			
<b>LEAD-5 (vs Glargine) , 2009</b> [NCT00331851] n=232/234 follow-up: 26 weeks	Liraglutide 1.8 mg daily versus Glargine on-top of sulphonylureas+metformin	adult patients with type 2 diabetes	Parallel groups open 17 countries
<b>exenatide weekly vs liraglutide</b>			
<b>H8O-MC-GWDE</b> <i>ongoing</i> [NCT01029886] n=NA follow-up: 36 weeks	exenatide once weekly for 26 weeks versus once-daily liraglutide for 26 weeks	-	open argentina
<b>liraglutide 1.8mg vs liraglutide 1.2mg</b>			
<b>LEAD 1 (1.8 vs 1.2) , 2009</b> n=NA	-	-	
<b>LEAD 2 (1.8 vs 1.2) , 2009</b> n=NA	-	-	
<b>LEAD 4 (1.8 vs 1.2) , 2009</b> n=NA	-	-	
<b>Pratley (1.8 vs 1.2) , 2010</b> n=NA	-	-	
<b>exenatide 10g twice daily vs liraglutide 1.8mg</b>			

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
LEAD 6 (Buse) exe vs lira , 2009 n=NA	-	-	
<b>liraglutide other doses vs metformin</b>			
Feinglos , 2005 n=176/34 follow-up: 12 weeks	Liraglutide 0.045, 0.225, 0.45, 0.6 or 0.75 mg daily for 12 weeks versus metformin 1000mg twice daily	subjects with Type 2 diabetes	Parallel groups double blind (not adequate)
<b>exenatide weekly vs pioglitazone</b>			
DURATION-2 (Bergental) (vs pioglitazone) , 2010 [NCT00637273] n=170/172 follow-up: 26 weeks	2 mg injected exenatide once weekly plus oral placebo once daily versus 45 mg oral pioglitazone once daily plus injected placebo once weekly	-	Parallel groups double blind USA, India, and Mexico
<b>taspoglutide vs pioglitazone</b>			
BC21893 ongoing [NCT00909597] n=NA follow-up: 24 months	taspoglutide 10mg sc weekly, or taspoglutide 20mg sc weekly after 4 weeks of taspoglutide 10mg sc weekly versus pioglitazone 45mg/day po after 4 weeks of pioglitazone 30mg/day po	patients with type 2 diabetes mellitus inadequately controlled with sulfonylurea monotherapy or sulfonylurea plus metformin combination therapy	parallel groups double-blind USA
<b>liraglutide 1.2mg vs rosiglitazone</b>			
LEAD-1 SU (1.2 vs rosiglitazone) , 2009 n=NA	-	-	
<b>liraglutide other doses vs rosiglitazone (add on SU)</b>			
LEAD-1 SU (1.8 vs rosiglitazone) , 2009 [NCT00318422] n=228/232 follow-up: 26 weeks	Liraglutide 0.6, 1.2 or 1.8 mg daily versus rosiglitazone on-top of sulphonylureas		Parallel groups double-blind 21 countries
<b>exenatide 20g/d vs rosiglitazone add on MET</b>			
DeFronzo (EXE vs ROSI) , 2010 n=45/45 follow-up: 20 weeks	EXE 10 microg b.i.d. versus ROSI 4 mg b.i.d.	subjects with type 2 diabetes on metformin	Parallel groups open
<b>exenatide weekly vs sitagliptin</b>			
DURATION-2 (Bergental) (vs sitagliptin) , 2010 [NCT00637273] n=170 follow-up: 26 weeks	2 mg injected exenatide once weekly plus oral placebo once daily versus 100 mg oral sitagliptin once daily plus injected placebo once weekly	patients treated with metformin	double blind USA, India, and Mexico
<b>liraglutide 1.2mg vs sitagliptin</b>			

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>Pratley 1.2mg , 2010</b> [NCT00700817] n=225/219 follow-up: 26 weeks	liraglutide 1.2mg subcutaneously once daily versus oral sitagliptin 100mg once daily	patients with type 2 diabetes who did not have adequate glycemic control with metformin	Parallel groups open Europe, USA, Canada
<b>liraglutide 1.8mg vs sitagliptin</b>			
<b>Pratley 1.8mg , 2010</b> [NCT00700817] n=221/219 follow-up: 26 weeks	liraglutide 1.8mg subcutaneously once daily versus oral sitagliptin 100mg once daily	patients with type 2 diabetes who did not have adequate glycemic control with metformin	Parallel groups open Europe, USA, Canada
<b>exenatide 20g/d vs sitagliptin (add on MET)</b>			
<b>DeFronzo , 2008</b> [NCT00477581] n=NA follow-up: 2 weeks	exenatide subcutaneous injection versus sitagliptin (100 mg QAM) for 2 weeks	metformin-treated T2D patients	Cross over double-blind USA
<b>exenatide 20g/d vs weekly exenatide</b>			
<b>Drucker , 2008</b> [NCT00308139] n=NA follow-up: 30 weeks	10 mug exenatide administered twice a day versus long-acting release formulation of exenatide 2 mg administered once weekly	patients with type 2 diabetes naive to drug therapy, or on one or more oral antidiabetic agents	Parallel groups open

## References

### **NN2211-1333, :**

Harder H, Nielsen L, Tu DT, Astrup A The effect of liraglutide, a long-acting glucagon-like peptide 1 derivative, on glycemic control, body composition, and 24-h energy expenditure in patients with type 2 diabetes. *Diabetes Care* 2004;27:1915-21 [[15277417](#)]

### **NCT00360334, :**

#### **Derosa, 2010:**

Derosa G, Maffioli P, Salvadeo SA, Ferrari I, Ragonesi PD, Querci F, Franzetti IG, Gadaleta G, Ciccarelli L, Piccinni MN, D'Angelo A, Cicero AF Exenatide versus glibenclamide in patients with diabetes. *Diabetes Technol Ther* 2010;12:233-40 [[20151774](#)] [10.1089/dia.2009.0141](#)

#### **Liutkus, 2010:**

Liutkus J, Rosas Guzman J, Norwood P, Pop L, Northrup J, Cao D, Trautmann M A placebo-controlled trial of exenatide twice-daily added to thiazolidinediones alone or in combination with metformin. *Diabetes Obes Metab* 2010;12:1058-65 [[20977576](#)] [10.1111/j.1463-1326.2010.01251.x](#)

#### **H8O-MC-GWBJ, 9698, 10g/d, 2008:**

#### **Apovian, 2010:**

Apovian CM, Bergenstal RM, Cuddihy RM, Qu Y, Lenox S, Lewis MS, Glass LC Effects of exenatide combined with lifestyle modification in patients with type 2 diabetes. *Am J Med* 2010;123:468.e9-17 [[20399326](#)] [10.1016/j.amjmed.2009.11.019](#)

#### **H8O-MC-GWBJ, 9698, 20g/d, 2008:**

#### **Moretto (DOUBLONS avec drucker), 2008:**

Moretto TJ, Milton DR, Ridge TD, Macconell LA, Okerson T, Wolka AM, Brodows RG Efficacy and tolerability of exenatide monotherapy over 24 weeks in antidiabetic drug-naive patients with type 2 diabetes: a randomized, double-blind, placebo-controlled, parallel-group study. *Clin Ther* 2008;30:1448-60 [[18803987](#)] [10.1016/j.clinthera.2008.08.006](#)

### **NCT00085969, :**

#### **Poon, 2005:**

Poon T, Nelson P, Shen L, Mihm M, Taylor K, Fineman M, Kim D Exenatide improves glycemic control and reduces body weight in subjects with type 2 diabetes: a dose-ranging study. *Diabetes Technol Ther* 2005;7:467-77 [15929678] [10.1089/dia.2005.7.467](#)

Nelson P, Poon T, Guan X, Schnabel C, Wintle M, Fineman M The incretin mimetic exenatide as a monotherapy in patients with type 2 diabetes. *Diabetes Technol Ther* 2007;9:317-26 [17705687] [10.1089/dia.2006.0024](#)

**Harder, 2004:**

Harder H, Nielsen L, Tu DT, Astrup A, The effect of liraglutide, a long-acting glucagon-like peptide 1 derivative, on glycemic control, body composition, and 24-h energy expenditure in patients with type 2 diabetes. *Diabetes Care* 2004;27:1915-21. [15277417]

**Kaku 0.6mg, 2010:**

Kaku K, Rasmussen MF, Clauson P, Seino Y Improved glycaemic control with minimal hypoglycaemia and no weight change with the once-daily human glucagon-like peptide-1 analogue liraglutide as add-on to sulphonylurea in Japanese patients with type 2 diabetes. *Diabetes Obes Metab* 2010;12:341-7 [20380655] [10.1111/j.1463-1326.2009.01194.x](#)

**Madsbad (vs placebo), 2004:**

Madsbad S, Schmitz O, Ranstam J, Jakobsen G, Matthews DR Improved glycemic control with no weight increase in patients with type 2 diabetes after once-daily treatment with the long-acting glucagon-like peptide 1 analog liraglutide (NN2211): a 12-week, double-blind, randomized, controlled trial. *Diabetes Care* 2004;27:1335-42 [15161785]

**NN2211-1571 (Vilsbll), 2007:**

Vilsbll T, Zdravkovic M, Le-Thi T, Krarup T, Schmitz O, Courrges JP, Verhoeven R, Bugnov I, Madsbad S Liraglutide, a long-acting human glucagon-like peptide-1 analog, given as monotherapy significantly improves glycemic control and lowers body weight without risk of hypoglycemia in patients with type 2 diabetes. *Diabetes Care* 2007 Jun;30:1608-10 [17372153]

Courrges JP, Vilsbll T, Zdravkovic M, Le-Thi T, Krarup T, Schmitz O, Verhoeven R, Bugov I, Madsbad S, Beneficial effects of once-daily liraglutide, a human glucagon-like peptide-1 analogue, on cardiovascular risk biomarkers in patients with Type 2 diabetes. *Diabet Med* 2008;25:1129-31. [19183322] [10.1111/j.1464-5491.2008.02484.x](#)

Horowitz M, Vilsbll T, Zdravkovic M, Hammer M, Madsbad S, Patient-reported rating of gastrointestinal adverse effects during treatment of type 2 diabetes with the once-daily human GLP-1 analogue, liraglutide. *Diabetes Obes Metab* 2008;10:593-6. [18435773] [10.1111/j.1463-1326.2008.00861.x](#)

**Seino, 2008:**

Seino Y, Rasmussen MF, Zdravkovic M, Kaku K Dose-dependent improvement in glycemia with once-daily liraglutide without hypoglycemia or weight gain: A double-blind, randomized, controlled trial in Japanese patients with type 2 diabetes. *Diabetes Res Clin Pract* 2008;81:161-8 [18495285] [10.1016/j.diabres.2008.03.018](#)

**LIBRA, 0:**

**NCT00978393, 0:**

**NN2211-1799, 0:**

**NN2211-1800, 0:**

**NN2211-3619, 0:**

**NN8022-1922, 0:**

**NN8022-1923, 0:**

**BC20750, 0:**

**BC20963, 0:**

**BC21713 (vs placebo), 0:**

**BP21572, 0:**

**Nauck 10 once weekly vs PBO, 2009:**

Nauck MA, Ratner RE, Kapitza C, Berria R, Boldrin M, Balena R, Treatment with the human once-weekly glucagon-like peptide-1 analog taspoglutide in combination with metformin improves glycemic control and lowers body weight in patients with type 2 diabetes inadequately controlled with metformin alone: a double-blind placebo-controlled study. *Diabetes Care* 2009;32:1237-43. [19366970] [10.2337/dc08-1961](#)

**Nauck 20 every 2 weeks vs PBO, 2009:**

**Nauck 20 once weekly vs PBO, 2009:**

**Buse, 2011:**

Buse JB, Bergenstal RM, Glass LC, Heilmann CR, Lewis MS, Kwan AY, Hoogwerf BJ, Rosenstock J Use of twice-daily exenatide in Basal insulin-treated patients with type 2 diabetes: a randomized, controlled trial. *Ann Intern Med* 2011 Jan 18;154:103-12 [21138825] [10.1059/0003-4819-154-2-201101180-00300](#)

**Fineman, 2003:**

Fineman MS, Bicsak TA, Shen LZ, Taylor K, Gaines E, Varns A, Kim D, Baron AD Effect on glycemic control of exenatide (synthetic exendin-4) additive to existing metformin and/or sulfonylurea treatment in patients with type 2 diabetes. *Diabetes Care* 2003;26:2370-7 [12882864]

**DeFronzo 10g/d, 2005:**

DeFronzo RA, Ratner RE, Han J, Kim DD, Fineman MS, Baron AD Effects of exenatide (exendin-4) on glycemic control and weight over 30 weeks in metformin-treated patients with type 2 diabetes. *Diabetes Care* 2005;28:1092-100 [15855572]

Blonde L, Klein EJ, Han J, Zhang B, Mac SM, Poon TH, Taylor KL, Trautmann ME, Kim DD, Kendall DM Interim analysis of the effects of exenatide treatment on A1C, weight and cardiovascular risk factors over 82 weeks in 314 overweight patients with type 2 diabetes. *Diabetes Obes Metab* 2006;8:436-47 [16776751] [10.1111/j.1463-1326.2006.00602.x](#)

Ratner RE, Maggs D, Nielsen LL, Stonehouse AH, Poon T, Zhang B, Bicsak TA, Brodows RG, Kim DD Long-term effects of exenatide therapy over 82 weeks on glycaemic control and weight in over-weight metformin-treated patients with type 2 diabetes mellitus. *Diabetes Obes Metab* 2006;8:419-28 [16776749] [10.1111/j.1463-1326.2006.00589.x](#)

**DeFronzo 20g/d, 2005:**

DeFronzo RA, Ratner RE, Han J, Kim DD, Fineman MS, Baron AD Effects of exenatide (exendin-4) on glycemic control and weight over 30 weeks in metformin-treated patients with type 2 diabetes. *Diabetes Care* 2005;28:1092-100 [15855572]

**Kim, 2007:**

Kim D, MacConell L, Zhuang D, Kothare PA, Trautmann M, Fineman M, Taylor K Effects of once-weekly dosing of a long-acting release formulation of exenatide on glucose control and body weight in subjects with type 2 diabetes. *Diabetes Care* 2007;30:1487-93 [17353504] [10.2337/dc06-2375](#)

**LEAD-2 (Nauck) (1.2mg vs placebo), 2009:**

Nauck M, Frid A, Hermansen K, Shah NS, Tankova T, Mitha IH, Zdravkovic M, Dring M, Matthews DR Efficacy and safety comparison of liraglutide, glimepiride, and placebo, all in combination with metformin, in type 2 diabetes: the LEAD (liraglutide effect and action in diabetes)-2 study. *Diabetes Care* 2009;32:84-90 [18931095] [10.2337/dc08-1355](#)

Sullivan SD, Alfonso-Cristancho R, Conner C, Hammer M, Blonde L Long-term outcomes in patients with type 2 diabetes receiving glimepiride combined with liraglutide or rosiglitazone. *Cardiovasc Diabetol* 2009;8:12 [19245711] [10.1186/1475-2840-8-12](#)

Nauck M, Marre M Adding liraglutide to oral antidiabetic drug monotherapy: efficacy and weight benefits. *Postgrad Med* 2009;121:5-15 [19491535] [10.3810/pgm.2009.05.1997](#)

**LEAD-2 (Nauck) (1.8mg vs placebo), 2009:**

Nauck M, Frid A, Hermansen K, Shah NS, Tankova T, Mitha IH, Zdravkovic M, Dring M, Matthews DR Efficacy and safety comparison of liraglutide, glimepiride, and placebo, all in combination with metformin, in type 2 diabetes: the LEAD (liraglutide effect and action in diabetes)-2 study. *Diabetes Care* 2009;32:84-90 [18931095] [10.2337/dc08-1355](#)

**NN2211-1796, 0:**

Buse JB, Garber A, Rosenstock J, Schmidt WE, Brett JH, Videbk N, Holst J, Nauck M Liraglutide treatment is associated with a low frequency and magnitude of antibody formation with no apparent impact on glycemic response or increased frequency of adverse events: results from the Liraglutide Effect and Action in Diabetes (LEAD) trials. *J Clin Endocrinol Metab* 2011;96:1695-702 [21450987] [10.1210/jc.2010-2822](#)

**NCT01234649, 0:****Ratner (20mg once weekly), 2010:**

Ratner R, Nauck M, Kapitza C, Asnaghi V, Boldrin M, Balena R, Safety and tolerability of high doses of taspoglutide, a once-weekly human GLP-1 analogue, in diabetic patients treated with metformin: a randomized double-blind placebo-controlled study. *Diabet Med* 2010;27:556-62. [20536952] [10.1111/j.1464-5491.2010.02990.x](#)

**BC22092, 0:****Gao, 2009:**

Gao Y, Yoon KH, Chuang LM, Mohan V, Ning G, Shah S, Jang HC, Wu TJ, Johns D, Northrup J, Brodows R Efficacy and safety of exenatide in patients of Asian descent with type 2 diabetes inadequately controlled with metformin or metformin and a sulphonylurea. *Diabetes Res Clin Pract* 2009;83:69-76 [19019476] [10.1016/j.diabres.2008.09.037](#)

**Buse 10g/d, 2004:**



Buse JB, Henry RR, Han J, Kim DD, Fineman MS, Baron AD Effects of exenatide (exendin-4) on glycemic control over 30 weeks in sulfonylurea-treated patients with type 2 diabetes. *Diabetes Care* 2004;27:2628-35 [[15504997](#)]

Riddle MC, Henry RR, Poon TH, Zhang B, Mac SM, Holcombe JH, Kim DD, Maggs DG Exenatide elicits sustained glycaemic control and progressive reduction of body weight in patients with type 2 diabetes inadequately controlled by sulphonylureas with or without metformin. *Diabetes Metab Res Rev* 2006;22:483-91 [[16634116](#)] [10.1002/dmrr.646](#)

Blonde L, Klein EJ, Han J, Zhang B, Mac SM, Poon TH, Taylor KL, Trautmann ME, Kim DD, Kendall DM Interim analysis of the effects of exenatide treatment on A1C, weight and cardiovascular risk factors over 82 weeks in 314 overweight patients with type 2 diabetes. *Diabetes Obes Metab* 2006;8:436-47 [[16776751](#)] [10.1111/j.1463-1326.2006.00602.x](#)

#### **Buse 20g/d, 2004:**

Buse JB, Henry RR, Han J, Kim DD, Fineman MS, Baron AD Effects of exenatide (exendin-4) on glycemic control over 30 weeks in sulfonylurea-treated patients with type 2 diabetes. *Diabetes Care* 2004;27:2628-35 [[15504997](#)]

#### **LEAD-1 SU (1.2 mg vs placebo), 2009:**

Nauck M, Marre M Adding liraglutide to oral antidiabetic drug monotherapy: efficacy and weight benefits. *Postgrad Med* 2009;121:5-15 [[19491535](#)] [10.3810/pgm.2009.05.1997](#)

Marre M, Shaw J, Brndle M, Bebakar WM, Kamaruddin NA, Strand J, Zdravkovic M, Le Thi TD, Colagiuri S Liraglutide, a once-daily human GLP-1 analogue, added to a sulphonylurea over 26 weeks produces greater improvements in glycaemic and weight control compared with adding rosiglitazone or placebo in subjects with Type 2 diabetes (LEAD-1 SU). *Diabet Med* 2009;26:268-78 [[19317822](#)] [10.1111/j.1464-5491.2009.02666.x](#)

Gallwitz B, Vaag A, Falahati A, Madsbad S Adding liraglutide to oral antidiabetic drug therapy: onset of treatment effects over time. *Int J Clin Pract* 2010;64:267-76 [[19925617](#)] [10.1111/j.1742-1241.2009.02265.x](#)

#### **LEAD-1 SU (1.8 mg vs placebo), 2009:**

Marre M, Shaw J, Brndle M, Bebakar WM, Kamaruddin NA, Strand J, Zdravkovic M, Le Thi TD, Colagiuri S Liraglutide, a once-daily human GLP-1 analogue, added to a sulphonylurea over 26 weeks produces greater improvements in glycaemic and weight control compared with adding rosiglitazone or placebo in subjects with Type 2 diabetes (LEAD-1 SU). *Diabet Med* 2009;26:268-78 [[19317822](#)] [10.1111/j.1464-5491.2009.02666.x](#)

#### **NN2211-1701, 0:**

Hegeds L, Moses AC, Zdravkovic M, Le Thi T, Daniels GH GLP-1 and calcitonin concentration in humans: lack of evidence of calcitonin release from sequential screening in over 5000 subjects with type 2 diabetes or nondiabetic obese subjects treated with the human GLP-1 analog, liraglutide. *J Clin Endocrinol Metab* 2011;96:853-60 [[21209033](#)] [10.1210/jc.2010-2318](#)

#### **Kadowaki (trial 8683), 2009:**

Kadowaki T, Namba M, Yamamura A, Sowa H, Wolka AM, Brodows RG Exenatide exhibits dose-dependent effects on glycemic control over 12 weeks in Japanese patients with suboptimally controlled type 2 diabetes. *Endocr J* 2009;56:415-24 [[19194050](#)]

#### **Kendall 10g/d, 2005:**

Kendall DM, Riddle MC, Rosenstock J, Zhuang D, Kim DD, Fineman MS, Baron AD Effects of exenatide (exendin-4) on glycemic control over 30 weeks in patients with type 2 diabetes treated with metformin and a sulfonylurea. *Diabetes Care* 2005;28:1083-91 [[15855571](#)]

#### **Kendall 20g/d, 2005:**

Kendall DM, Riddle MC, Rosenstock J, Zhuang D, Kim DD, Fineman MS, Baron AD Effects of exenatide (exendin-4) on glycemic control over 30 weeks in patients with type 2 diabetes treated with metformin and a sulfonylurea. *Diabetes Care* 2005;28:1083-91 [[15855571](#)]

#### **LEAD-5 (vs placebo), 2009:**

Russell-Jones D, Vaag A, Schmitz O, Sethi BK, Lalic N, Antic S, Zdravkovic M, Ravn GM, Sim R Liraglutide vs insulin glargine and placebo in combination with metformin and sulfonylurea therapy in type 2 diabetes mellitus (LEAD-5 met+SU): a randomised controlled trial. *Diabetologia* 2009;52:2046-55 [[19688338](#)] [10.1007/s00125-009-1472-y](#)

#### **Zinman 20g/j, 2007:**

Zinman B, Hoogwerf BJ, Durn Garca S, Milton DR, Giaconia JM, Kim DD, Trautmann ME, Brodows RG The effect of adding exenatide to a thiazolidinedione in suboptimally controlled type 2 diabetes: a randomized trial. *Ann Intern Med* 2007;146:477-85 [[17404349](#)]

#### **Zinman 20g/j A MODIFIER, 2007:**

Zinman B, Hoogwerf BJ, Durn Garca S, Milton DR, Giaconia JM, Kim DD, Trautmann ME, Brodows RG The effect of adding exenatide to a thiazolidinedione in suboptimally controlled type 2 diabetes: a randomized trial. *Ann Intern Med* 2007;146:477-85 [[17404349](#)]

**LEAD-4 (1.2mg), 2009:**

Zinman B, Gerich J, Buse JB, Lewin A, Schwartz S, Raskin P, Hale PM, Zdravkovic M, Blonde L, , Efficacy and safety of the human glucagon-like peptide-1 analog liraglutide in combination with metformin and thiazolidinedione in patients with type 2 diabetes (LEAD-4 Met+TZD). *Diabetes Care* 2009;32:1224-30. [19289857] [10.2337/dc08-2124](#)

**LEAD-4 (1.8mg), 2009:**

Zinman B, Gerich J, Buse JB, Lewin A, Schwartz S, Raskin P, Hale PM, Zdravkovic M, Blonde L, , Efficacy and safety of the human glucagon-like peptide-1 analog liraglutide in combination with metformin and thiazolidinedione in patients with type 2 diabetes (LEAD-4 Met+TZD). *Diabetes Care* 2009;32:1224-30. [19289857] [10.2337/dc08-2124](#)

**Kaku 0.9mg, 2010:**

Kaku K, Rasmussen MF, Clauson P, Seino Y Improved glycaemic control with minimal hypoglycaemia and no weight change with the once-daily human glucagon-like peptide-1 analogue liraglutide as add-on to sulphonylurea in Japanese patients with type 2 diabetes. *Diabetes Obes Metab* 2010;12:341-7 [20380655] [10.1111/j.1463-1326.2009.01194.x](#)

**Gill, 2010:**

Gill A, Hoogwerf BJ, Burger J, Bruce S, Macconell L, Yan P, Braun D, Giaconia J, Malone J Effect of exenatide on heart rate and blood pressure in subjects with type 2 diabetes mellitus: a double-blind, placebo-controlled, randomized pilot study. *Cardiovasc Diabetol* 2010;9:6 [20109208] [10.1186/1475-2840-9-6](#)

**NC25113, 0:****MK-0431-403, 0:****phase 2 exenatide once monthly, :****Bergenstal (once daily), 2009:**

Bergenstal R, Lewin A, Bailey T, Chang D, Gylvin T, Roberts V Efficacy and safety of biphasic insulin aspart 70/30 versus exenatide in subjects with type 2 diabetes failing to achieve glycemic control with metformin and a sulphonylurea. *Curr Med Res Opin* 2009;25:65-75 [19210140] [10.1185/03007990802597951](#)

**Bergenstal (twice daily), 2009:**

Bergenstal R, Lewin A, Bailey T, Chang D, Gylvin T, Roberts V Efficacy and safety of biphasic insulin aspart 70/30 versus exenatide in subjects with type 2 diabetes failing to achieve glycemic control with metformin and a sulphonylurea. *Curr Med Res Opin* 2009;25:65-75 [19210140] [10.1185/03007990802597951](#)

**BC21625, 0:****Exenatide Trial 10749, :****LEAD-6, 2009:**

Buse JB, Rosenstock J, Sesti G, Schmidt WE, Montanya E, Brett JH, Zychma M, Blonde L Liraglutide once a day versus exenatide twice a day for type 2 diabetes: a 26-week randomised, parallel-group, multinational, open-label trial (LEAD-6). *Lancet* 2009 Jul 4;374:39-47 [19515413] [10.1016/S0140-6736\(09\)60659-0](#)

Buse JB, Sesti G, Schmidt WE, Montanya E, Chang CT, Xu Y, Blonde L, Rosenstock J Switching to once-daily liraglutide from twice-daily exenatide further improves glycemic control in patients with type 2 diabetes using oral agents. *Diabetes Care* 2010;33:1300-3 [20332351] [10.2337/dc09-2260](#)

Schmidt WE, Christiansen JS, Hammer M, Zychma MJ, Buse JB Patient-reported outcomes are superior in patients with Type?2 diabetes treated with liraglutide as compared with exenatide, when added to metformin, sulphonylurea or both: results from a randomized, open-label study. *Diabet Med* 2011 Mar 9; [21388442] [10.1111/j.1464-5491.2011.03276.x](#)

**HEELA (Davies), 2009:**

Davies MJ, Donnelly R, Barnett AH, Jones S, Nicolay C, Kilcoyne A Exenatide compared with long-acting insulin to achieve glycaemic control with minimal weight gain in patients with type 2 diabetes: results of the Helping Evaluate Exenatide in patients with diabetes compared with Long-Acting insulin (HEELA) study. *Diabetes Obes Metab* 2009 Dec;11:1153-62 [19930005]

**Seino, 2010:**

Seino Y, Rasmussen MF, Nishida T, Kaku K, Efficacy and safety of the once-daily human GLP-1 analogue, liraglutide, vs glibenclamide monotherapy in Japanese patients with type 2 diabetes. *Curr Med Res Opin* 2010;26:1013-22. [20199137] [10.1185/03007991003672551](#)

**LEAD-3 mono 1.2mg (Garber), 2009:**

Garber A, Henry R, Ratner R, Garcia-Hernandez PA, Rodriguez-Pattzi H, Olvera-Alvarez I, Hale PM, Zdravkovic M, Bode B Liraglutide versus glimepiride monotherapy for type 2 diabetes (LEAD-3 Mono): a randomised, 52-week, phase III, double-blind, parallel-treatment trial. *Lancet* 2009;373:473-81 [18819705] [10.1016/S0140-6736\(08\)61246-5](#)

Bode BW, Testa MA, Magwire M, Hale PM, Hammer M, Blonde L, Garber A, , Patient-reported outcomes following treatment with the human GLP-1 analogue liraglutide or glimepiride in monotherapy: results from a randomized controlled trial in patients with type 2 diabetes. *Diabetes Obes Metab* 2010;12:604-12. [20590735] [10.1111/j.1463-1326.2010.01196.x](#)

Garber A, Henry RR, Ratner R, Hale P, Chang CT, Bode B Liraglutide, a once-daily human glucagon-like peptide 1 analogue, provides sustained improvements in glycaemic control and weight for 2 years as monotherapy compared with glimepiride in patients with type 2 diabetes. *Diabetes Obes Metab* 2011;13:348-56 [21205128] [10.1111/j.1463-1326.2010.01356.x](#)

**LEAD-3 mono 1.8mg (Garber), 2009:**

Garber A, Henry R, Ratner R, Garcia-Hernandez PA, Rodriguez-Pattzi H, Olvera-Alvarez I, Hale PM, Zdravkovic M, Bode B Liraglutide versus glimepiride monotherapy for type 2 diabetes (LEAD-3 Mono): a randomised, 52-week, phase III, double-blind, parallel-treatment trial. *Lancet* 2009;373:473-81 [18819705] [10.1016/S0140-6736\(08\)61246-5](#)

**Madsbad (vs Glimepiride), 2004:**

Madsbad S, Schmitz O, Ranstam J, Jakobsen G, Matthews DR Improved glycemic control with no weight increase in patients with type 2 diabetes after once-daily treatment with the long-acting glucagon-like peptide 1 analog liraglutide (NN2211): a 12-week, double-blind, randomized, controlled trial. *Diabetes Care* 2004;27:1335-42 [15161785]

**LEAD-2 (Nauck) (1.2 mg vs glimepiride), 2009:**

Nauck M, Frid A, Hermansen K, Shah NS, Tankova T, Mitha IH, Zdravkovic M, Dring M, Matthews DR Efficacy and safety comparison of liraglutide, glimepiride, and placebo, all in combination with metformin, in type 2 diabetes: the LEAD (liraglutide effect and action in diabetes)-2 study. *Diabetes Care* 2009;32:84-90 [18931095] [10.2337/dc08-1355](#)

Hermansen K, Kolotkin RL, Hammer M, Zdravkovic M, Matthews D, Patient-reported outcomes in patients with type 2 diabetes treated with liraglutide or glimepiride, both as add-on to metformin. *Prim Care Diabetes* 2010;4:113-7. [20444662] [10.1016/j.pcd.2010.04.001](#)

*Diabetologia* 52 (2009) S8.

**LEAD-2 (Nauck) (1.8 mg vs glimepiride), 2009:**

Nauck M, Frid A, Hermansen K, Shah NS, Tankova T, Mitha IH, Zdravkovic M, Dring M, Matthews DR Efficacy and safety comparison of liraglutide, glimepiride, and placebo, all in combination with metformin, in type 2 diabetes: the LEAD (liraglutide effect and action in diabetes)-2 study. *Diabetes Care* 2009;32:84-90 [18931095] [10.2337/dc08-1355](#)

**Heine, 2005:**

Heine RJ, Van Gaal LF, Johns D, Mihm MJ, Widel MH, Brodows RG Exenatide versus insulin glargine in patients with suboptimally controlled type 2 diabetes: a randomized trial. *Ann Intern Med* 2005;143:559-69 [16230722]

**Barnett, 2007:**

Barnett AH, Burger J, Johns D, Brodows R, Kendall DM, Roberts A, Trautmann ME Tolerability and efficacy of exenatide and titrated insulin glargine in adult patients with type 2 diabetes previously uncontrolled with metformin or a sulfonylurea: a multinational, randomized, open-label, two-period, crossover noninferiority trial. *Clin Ther* 2007;29:2333-48 [18158075] [10.1016/j.clinthera.2007.11.006](#)

**Davis, 2007:**

Davis SN, Johns D, Maggs D, Xu H, Northrup JH, Brodows RG Exploring the substitution of exenatide for insulin in patients with type 2 diabetes treated with insulin in combination with oral antidiabetes agents. *Diabetes Care* 2007;30:2767-72 [17595353] [10.2337/dc06-2532](#)

**Nauck, 2007:**

Nauck MA, Duran S, Kim D, Johns D, Northrup J, Festa A, Brodows R, Trautmann M A comparison of twice-daily exenatide and biphasic insulin aspart in patients with type 2 diabetes who were suboptimally controlled with sulfonylurea and metformin: a non-inferiority study. *Diabetologia* 2007;50:259-67 [17160407] [10.1007/s00125-006-0510-2](#)

**DURATION-3 (Diamant), 2010:**

Diamant M, Van Gaal L, Stranks S, Northrup J, Cao D, Taylor K, Trautmann M Once weekly exenatide compared with insulin glargine titrated to target in patients with type 2 diabetes (DURATION-3): an open-label randomised trial. *Lancet* 2010 Jun 26;375:2234-2243 [20609969] [10.1016/S0140-6736\(10\)60406-0](#)

**EAGLE, 0:**

**ZC22565, 0:**

**Bunck, 2009:**

Bunck MC, Diamant M, Cornr A, Eliasson B, Malloy JL, Shaginian RM, Deng W, Kendall DM, Taskinen MR, Smith U, Yki-Jrvinen H, Heine RJ One-year treatment with exenatide improves beta-cell function, compared with insulin glargine, in metformin-treated type 2 diabetic patients: a randomized, controlled trial. *Diabetes Care* 2009;32:762-8

[19196887] [10.2337/dc08-1797](#)

Bunck MC, Cornr A, Eliasson B, Heine RJ, Shaginian RM, Taskinen MR, Smith U, Yki-Jrvinen H, Diamant M Effects of exenatide on measures of  $\beta$ -cell function after 3 years in metformin-treated patients with type 2 diabetes. *Diabetes Care* 2011;34:2041-7 [[21868779](#)] [10.2337/dc11-0291](#)

Bunck MC, Cornr A, Eliasson B, Heine RJ, Shaginian RM, Wu Y, Yan P, Smith U, Yki-Jrvinen H, Diamant M, Taskinen MR One-year treatment with exenatide vs. insulin glargine: effects on postprandial glycemia, lipid profiles, and oxidative stress. *Atherosclerosis* 2010;212:223-9 [[20494360](#)] [10.1016/j.atherosclerosis.2010.04.024](#)

**BC20965, 0:**

**Trial 8078, :**

**LEAD-5 (vs Glargine), 2009:**

Russell-Jones D, Vaag A, Schmitz O, Sethi BK, Lalic N, Antic S, Zdravkovic M, Ravn GM, Sim R Liraglutide vs insulin glargine and placebo in combination with metformin and sulfonylurea therapy in type 2 diabetes mellitus (LEAD-5 met+SU): a randomised controlled trial. *Diabetologia* 2009 Oct;52:2046-55 [[19688338](#)]

Sullivan SD, Alfonso-Cristancho R, Conner C, Hammer M, Blonde L Long-term outcomes in patients with type 2 diabetes receiving glimepiride combined with liraglutide or rosiglitazone. *Cardiovasc Diabetol* 2009 Feb 26;8:12 [[19245711](#)]

**H8O-MC-GWDE, 0:**

**LEAD 1 (1.8 vs 1.2), 2009:**

**LEAD 2 (1.8 vs 1.2), 2009:**

**LEAD 4 (1.8 vs 1.2), 2009:**

**Pratley (1.8 vs 1.2), 2010:**

**LEAD 6 (Buse) exe vs lira, 2009:**

Buse JB, Rosenstock J, Sesti G, Schmidt WE, Montanya E, Brett JH, Zychma M, Blonde L Liraglutide once a day versus exenatide twice a day for type 2 diabetes: a 26-week randomised, parallel-group, multinational, open-label trial (LEAD-6). *Lancet* 2009;374:39-47 [[19515413](#)] [10.1016/S0140-6736\(09\)60659-0](#)

**Feinglos, 2005:**

Feinglos MN, Saad MF, Pi-Sunyer FX, An B, Santiago O Effects of liraglutide (NN2211), a long-acting GLP-1 analogue, on glycaemic control and bodyweight in subjects with Type 2 diabetes. *Diabet Med* 2005;22:1016-23 [[16026367](#)] [10.1111/j.1464-5491.2005.01567.x](#)

**DURATION-2 (Bergental) (vs pioglitazone), 2010:**

Wysham C, Bergental R, Malloy J, Yan P, Walsh B, Malone J, Taylor K DURATION-2: efficacy and safety of switching from maximum daily sitagliptin or pioglitazone to once-weekly exenatide. *Diabet Med* 2011 Mar 22;: [[21434995](#)] [10.1111/j.1464-5491.2011.03301.x](#)

**BC21893, 0:**

**LEAD-1 SU (1.2 vs rosiglitazone), 2009:**

**LEAD-1 SU (1.8 vs rosiglitazone), 2009:**

Marre M, Shaw J, Brndle M, Bebakar WM, Kamaruddin NA, Strand J, Zdravkovic M, Le Thi TD, Colagiuri S Liraglutide, a once-daily human GLP-1 analogue, added to a sulphonylurea over 26 weeks produces greater improvements in glycaemic and weight control compared with adding rosiglitazone or placebo in subjects with Type 2 diabetes (LEAD-1 SU). *Diabet Med* 2009;26:268-78 [[19317822](#)] [10.1111/j.1464-5491.2009.02666.x](#)

**DeFronzo (EXE vs ROSI), 2010:**

DeFronzo RA, Triplitt C, Qu Y, Lewis MS, Maggs D, Glass LC Effects of exenatide plus rosiglitazone on beta-cell function and insulin sensitivity in subjects with type 2 diabetes on metformin. *Diabetes Care* 2010;33:951-7 [[20107105](#)] [10.2337/dc09-1521](#)

**DURATION-2 (Bergental) (vs sitagliptin), 2010:**

Bergental RM, Wysham C, Macconell L, Malloy J, Walsh B, Yan P, Wilhelm K, Malone J, Porter LE Efficacy and safety of exenatide once weekly versus sitagliptin or pioglitazone as an adjunct to metformin for treatment of type 2 diabetes (DURATION-2): a randomised trial. *Lancet* 2010 Jun 25;: [[20580422](#)] [10.1016/S0140-6736\(10\)60590-9](#)

**Pratley 1.2mg, 2010:**

Pratley RE, Nauck M, Bailey T, Montanya E, Cuddihy R, Filetti S, Thomsen AB, Sndergaard RE, Davies M Liraglutide versus sitagliptin for patients with type 2 diabetes who did not have adequate glycaemic control with metformin: a 26-week, randomised, parallel-group, open-label trial. *Lancet* 2010 Apr 24;375:1447-1456 [[20417856](#)] [10.1016/S0140-6736\(10\)60307-8](#)

Pratley R, Nauck M, Bailey T, Montanya E, Cuddihy R, Filetti S, Garber A, Thomsen AB, Hartvig H, Davies M One year of liraglutide treatment offers sustained and more effective glycaemic control and weight reduction compared with sitagliptin, both in combination with metformin, in patients with type 2 diabetes: a randomised, parallel-group, open-label trial. *Int J Clin Pract* 2011;65:397-407 [21355967] [10.1111/j.1742-1241.2011.02656.x](https://doi.org/10.1111/j.1742-1241.2011.02656.x)

**Pratley 1.8mg, 2010:**

Pratley RE, Nauck M, Bailey T, Montanya E, Cuddihy R, Filetti S, Thomsen AB, Sndergaard RE, Davies M Liraglutide versus sitagliptin for patients with type 2 diabetes who did not have adequate glycaemic control with metformin: a 26-week, randomised, parallel-group, open-label trial. *Lancet* 2010 Apr 24;375:1447-1456 [20417856] [10.1016/S0140-6736\(10\)60307-8](https://doi.org/10.1016/S0140-6736(10)60307-8)

**DeFronzo, 2008:**

DeFronzo RA, Okerson T, Viswanathan P, Guan X, Holcombe JH, MacConell L Effects of exenatide versus sitagliptin on postprandial glucose, insulin and glucagon secretion, gastric emptying, and caloric intake: a randomized, cross-over study. *Curr Med Res Opin* 2008;24:2943-52 [18786299] [10.1185/03007990802418851](https://doi.org/10.1185/03007990802418851)

**Drucker, 2008:**

Drucker DJ, Buse JB, Taylor K, Kendall DM, Trautmann M, Zhuang D, Porter L Exenatide once weekly versus twice daily for the treatment of type 2 diabetes: a randomised, open-label, non-inferiority study. *Lancet* 2008 Oct 4;372:1240-50 [18782641]

Buse JB, Drucker DJ, Taylor KL, Kim T, Walsh B, Hu H, Wilhelm K, Trautmann M, Shen LZ, Porter LE DURATION-1: exenatide once weekly produces sustained glycemic control and weight loss over 52 weeks. *Diabetes Care* 2010;33:1255-61 [20215461] [10.2337/dc09-1914](https://doi.org/10.2337/dc09-1914)

## 4 lixisenatide

Trial	Treatments	Patients	Trials design and methods
<b>lixisenatide vs control</b>			
<b>GETGOAL-MONO Japan</b> <i>LTS ongoing</i> [NCT00905255] n=NA follow-up: 24-76 weeks	AVE0010 once daily treatment in monotherapy versus usual care	type 2 diabetes patients	Parallel groups open Japan
<b>lixisenatide vs placebo</b>			
<b>ELIXA (EFC11319)</b> <i>ongoing</i> [NCT01147250] n=NA follow-up:	Lixisenatide 20 g one daily versus placebo	type 2 diabetic patients who experienced an acute coronary syndrome	double-blind USA
<b>GETGOAL-MONO</b> <i>ongoing</i> [NCT00688701] n=NA follow-up: 12 weeks	AVE0010 in monotherapy versus lixisenatide	Type 2 diabetes mellitus not treated with any antidiabetic agent	Parallel groups double-blind USA
<b>lixisenatide vs placebo (add on basal insulin)</b>			
<b>GETGOAL-L</b> <i>unpublished</i> [NCT00715624] n=328/167 follow-up: 24 weeks	AVE0010 (10,15 and 20 g) in association with basal insulin, with or without metformin versus placebo on top basal insulin	Type 2 diabetes mellitus insufficiently controlled with basal insulin with or without metformin	double-blind USA
<b>GetGoal Duo1</b> <i>ongoing</i> [NCT00975286] n=NA follow-up: 24 weeks	Lixisenatide as an add-on treatment to insulin glargine and metformin versus placebo	patients with type 2 diabetes insufficiently controlled with insulin glargine and metformin	Parallel groups double-blind USA

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>GETGOAL-L-ASIA</b> <i>ongoing</i> [NCT00866658] n=NA follow-up: 24 weeks	24 weeks of AVE0010 versus placebo on Top of Basal Insulin +/- Sulfonylurea	Patients With Type 2 Diabetes Insufficiently Controlled With Basal Insulin With or Without Sulfonylurea	double-blind Japan
<b>lixisenatide vs placebo (add on MET)</b>			
<b>Ratner DRI6012 , 2010</b> [NCT00299871] n=433/109 follow-up: 13 weeks	subcutaneous lixisenatide doses of 5, 10, 20 or 30 microg once daily or twice daily versus placebo	patients with Type 2 diabetes inadequately controlled with metformin (>= 1000 mg/day)	Parallel groups double-blind (nature not volume) multinational
<b>GETGOAL-F1</b> <i>ongoing</i> [NCT00763451] n=NA follow-up: 24 weeks	AVE0010 in association with metformin versus placebo	Type 2 diabetes mellitus insufficiently controlled with metformin	double-blind USA
<b>GETGOAL-M</b> <i>ongoing</i> [NCT00712673] n=NA follow-up: 24 weeks	-	Type 2 diabetes mellitus insufficiently controlled with metformin	Parallel groups double-blind USA
<b>lixisenatide vs placebo (add on MET +/-SU)</b>			
<b>GetGoal-M-As</b> <i>ongoing</i> [NCT01169779] n=NA follow-up: 24 weeks	Lixisenatide (Titration phase: 10 g maintenance phase: 20 g, add-on treatment to metformin with or without sulfonylurea versus add-on treatment to metformin with or without sulfonylurea	-	double-blind China
<b>lixisenatide vs placebo (add on SU +/-MET)</b>			
<b>GETGOAL-S</b> <i>ongoing</i> [NCT00713830] n=NA follow-up: 24 weeks	AVE0010 in association with sulfonylurea without or with metformin versus placebo	patients with type 2 diabetes not adequately controlled with sulfonylurea	double-blind USA
<b>lixisenatide vs placebo (add on TZD +/-MET)</b>			
<b>GETGOAL-P</b> <i>ongoing</i> [NCT00763815] n=NA follow-up: 24 weeks	AVE0010 in association with pioglitazone with or without metformin versus placebo	Type 2 diabetes mellitus insufficiently controlled with pioglitazone with or without metformin	Parallel groups double-blind USA
<b>lixisenatide vs exenatide</b>			
<b>GetGoal-X</b> <i>ongoing</i> [NCT00707031] n=NA follow-up: 24 weeks	AVE0010 in association with metformin versus exenatide in association with metformin	type 2 diabetes	parallel groups open USA
<b>lixisenatide vs sitagliptin (add on MET)</b>			
<b>EFC10780 , 2010</b> <i>ongoing</i> [NCT00976937] n=NA follow-up: 24 weeks	Lixisenatide titrated 15-20 g once daily versus Sitagliptin (add-on to Metformin)	Obese Type 2 Diabetic Patients Younger Than 50	Parallel groups double-blind WW

## References

**GETGOAL-MONO Japan LTS, 0:**

**ELIXA (EFC11319), 0:**

**GETGOAL-MONO, 0:**

Fonseca VA, Alvarado-Ruiz R, Raccach D, Boka G, Miossec P, Gerich JE Efficacy and Safety of the Once-Daily GLP-1 Receptor Agonist Lixisenatide in Monotherapy: A randomized, double-blind, placebo-controlled trial in patients with type 2 diabetes (GetGoal-Mono). *Diabetes Care* 2012;; [22432104] 10.2337/dc11-1935

**GETGOAL-L, 0:**

**GetGoal Duo1, 0:**

10.2337/dc12-2462

**GETGOAL-L-ASIA, 0:**

**Ratner DRI6012, 2010:**

Ratner RE, Rosenstock J, Boka G Dose-dependent effects of the once-daily GLP-1 receptor agonist lixisenatide in patients with Type 2 diabetes inadequately controlled with metformin: a randomized, double-blind, placebo-controlled trial. *Diabet Med* 2010;27:1024-32 [20722676] 10.1111/j.1464-5491.2010.03020.x

**GETGOAL-F1, 0:**

**GETGOAL-M, 0:**

**GetGoal-M-As, 0:**

**GETGOAL-S, 0:**

**GETGOAL-P, 0:**

**GetGoal-X, 0:**

**EFC10780, 2010:**

## 5 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.