

Clinical trials of direct oral anticoagulant (DAO) for DVT prophylaxis in elective hip replacement

TrialResults-center www.trialresultscenter.org

1 direct factor Xa inhibitors

Trial	Treatments	Patients	Trials design and methods
apixaban vs enoxaparin			
ADVANCE 3 , 2010 [NCT00423319] n=2708/2699 follow-up: 35 days (+60)	apixaban 2.5mg twice daily for 35 days versus enoxaparin 40mg once daily for 35 days	patients undergoing elective total hip replacement surgery	Parallel groups double blind 21 countries
rivaroxaban vs enoxaparin			
RECORD 1 , 2008 [NCT00329628] n=2266/2275 follow-up: 36 days (range 30-42)	rivaroxaban 10mg once daily for 35 days versus enoxaparin 40mg subcutaneous once daily for 31-39 days	patients undergoing total hip arthroplasty	Parallel groups double blind 27 countries worldwide
edoxaban vs enoxaparin (short duration)			
STARS J-V [NCT01181167] n=255/248 follow-up:	edoxaban 30 mg once daily for 11 to 14 days versus subcutaneous enoxaparin 2,000 IU, equivalent to 20 mg, twice daily (BID) for 11 to 14 days	total hip arthroplasty	Parallel groups double-blind japan
rivaroxaban vs enoxaparin (short duration)			
ODIXa-HIP 10mg , 2006 n=142/157 follow-up: 5-9 days	rivaroxaban 10mg daily for 59 days versus once-daily subcutaneous enoxaparin dose of 40 mg for 59 days	patients undergoing elective total hip replacement	Parallel groups double blind Europe, Israel
rivaroxaban (long duration) vs enoxaparin (short duration)			
RECORD 2 , 2008 [NCT00332020] n=1252/1257 follow-up: 30-42 days	extended thromboprophylaxis with rivaroxaban 10mg once daily for 31-39 days versus enoxaparin 40mg subcutaneous once daily for 10-14 days	patients undergoing elective total hip replacement	Parallel groups double blind 21 countries worldwide

References

ADVANCE 3, 2010:

Lassen MR, Gallus A, Raskob GE, Pineo G, Chen D, Ramirez LM Apixaban versus Enoxaparin for Thromboprophylaxis after Hip Replacement. N Engl J Med 2010;363:2487-2498 [21175312] [10.1056/NEJMoa1006885](https://doi.org/10.1056/NEJMoa1006885)

RECORD 1, 2008:

Eriksson BI, Borris LC, Friedman RJ, Haas S, Huisman MV, Kakkar AK, Bandel TJ, Beckmann H, Muehlhofer E, Misselwitz F, Geerts W Rivaroxaban versus enoxaparin for

thromboprophylaxis after hip arthroplasty N Engl J Med 2008;358:2765-75 [18579811] 10.1056/NEJMoa0800374

STARS J-V, 0:

ODIXa-HIP 10mg, 2006:

Eriksson BI, Borris L, Dahl OE, Haas S, Huisman MV, Kakkar AK, Misselwitz F, Klebo P Oral, direct Factor Xa inhibition with BAY 59-7939 for the prevention of venous thromboembolism after total hip replacement. J Thromb Haemost 2006 Jan;4:121-8 [16409461]

Eriksson BI, Borris LC, Dahl OE, Haas S, Huisman MV, Kakkar AK, Muehlhofer E, Dierig C, Misselwitz F, Klebo P A once-daily, oral, direct Factor Xa inhibitor, rivaroxaban (BAY 59-7939), for thromboprophylaxis after total hip replacement. Circulation 2006 Nov 28;114:2374-81 [17116766]

RECORD 2, 2008:

Kakkar AK, Brenner B, Dahl OE, Eriksson BI, Mouret P, Muntz J, Soglian AG, Pap AF, Misselwitz F, Haas S Extended duration rivaroxaban versus short-term enoxaparin for the prevention of venous thromboembolism after total hip arthroplasty: a double-blind, randomised controlled trial. Lancet 2008 Jun 24;: [18582928]

2 oral direct thrombin inhibitor

Trial	Treatments	Patients	Trials design and methods
dabigatran 150mg vs enoxaparin			
RE-NOVATE (150mg) , 2007 [NCT00168818] n=1174/1162 follow-up: 28-35 days, median 33d	dabigatran etexilate 150 mg q.d. 28-35 days versus Enoxaparin 40 mg q.d. for 28-25 days	Total hip replacement	double blind Europe, Australia, South Africa
dabigatran 220mg vs enoxaparin			
RE-NOVATE 2 unpublished [NCT00657150] n=1010/1003 follow-up: 28-35 days (mean 32d)	dabigatran 220mg once daily for 28-35 Days versus enoxaparin 40mg subcutaneous once daily for 28-35 Days	patients undergoing total hip-replacement surgery	Parallel groups double-blind
RE-NOVATE (220mg) , 2007 [NCT00168818] n=1157/1162 follow-up: 28-35 days, median 33d	dabigatran etexilate 220 mg q.d. for 28-35 days versus Enoxaparin 40 mg q.d. for 23-35 days	Total hip replacement	Parallel groups double blind Europe, Australia, South Africa

References

RE-NOVATE (150mg), 2007:

Eriksson BI, Dahl OE, Rosencher N, Kurth AA, van Dijk CN, Frostick SP, Prins MH, Hettiarachchi R, Hantel S, Schnee J, Bller HR Dabigatran etexilate versus enoxaparin for prevention of venous thromboembolism after total hip replacement: a randomised, double-blind, non-inferiority trial. Lancet 2007;370:949-56 [17869635]

RE-NOVATE 2, 0:

RE-NOVATE (220mg), 2007:

Eriksson BI, Dahl OE, Rosencher N, Kurth AA, van Dijk CN, Frostick SP, Prins MH, Hettiarachchi R, Hantel S, Schnee J, Bller HR Dabigatran etexilate versus enoxaparin for prevention of venous thromboembolism after total hip replacement: a randomised, double-blind, non-inferiority trial. Lancet 2007;370:949-56 [17869635]

3 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.