

# Clinical trials of antithrombotics for thrombosis prevention in hip Fracture

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 Low molecular weight heparin

Trial	Treatments	Patients	Trials design and methods
<b>semuloparin vs enoxaparin</b>			
<b>SAVE-HIP 2 , 2012</b> [NCT00721760] n=500/503 follow-up:	Semuloparin 20 mg once-daily versus Enoxaparin 40 mg once-daily	hip fracture surgery	Parallel groups
<b>dalteparin vs placebo</b>			
<b>Jorgensen , 1989</b> n=30/38 follow-up: 9 days	dalteparin 5000 x1 versus Placebo	Hip fracture	double blind
<b>nadroparin vs placebo</b>			
<b>Sourmelis , 1995</b> n=72/78 follow-up: 10-12 days	nadroparin 3075x1 preop, 6150x1 post op versus Placebo	Hip fracture	double blind
<b>certoparin + DHE vs Unfractionated heparin</b>			
<b>Lassen , 1989</b> n=68/71 follow-up: 6 days	certoparin 3000+0.5mg DHE x1 versus placebo	Hip fracture	double blind

## References

### SAVE-HIP 2, 2012:

Lassen MR, Fisher W, Mouret P, Agnelli G, George D, Kakkar A, Mismetti P, Turpie AG Semuloparin for prevention of venous thromboembolism after major orthopedic surgery: results from three randomized clinical trials, SAVE-HIP1, SAVE-HIP2 and SAVE-KNEE. J Thromb Haemost 2012 May;10:822-32 [22429800] [10.1111/j.1538-7836.2012.04701.x](https://doi.org/10.1111/j.1538-7836.2012.04701.x)

### Jorgensen, 1989:

### Sourmelis, 1995:

### Lassen, 1989:

Lassen MR, Borris LC, Christiansen HM, Moller-Larsen F, Knudsen VE, Boris P, Nehen AM, Jurik AG, de Carvalho A, Nielsen BW Prevention of thromboembolism in hip-fracture patients. Comparison of low-dose heparin and low-molecular-weight heparin combined with dihydroergotamine. Arch Orthop Trauma Surg 1989;108:10-3 [2913977]

## 2 platelet aggregation inhibitors

Trial	Treatments	Patients	Trials design and methods
<b>aspirin vs placebo</b>			

continued...

Trial	Treatments	Patients	Trials design and methods
PEP hip-fracture , 2000 n=6679/6677 follow-up: 35 days	aspirin 160mg/d started preoperatively and continued for 35 days versus placebo	patients undergoing surgery for hip fracture	Parallel groups Double blind Australia, New Zealand, South Africa,

## References

### PEP hip-fracture, 2000:

Prevention of pulmonary embolism and deep vein thrombosis with low dose aspirin: Pulmonary Embolism Prevention (PEP) trial. Lancet 2000 Apr 15;355:1295-302 [[10776741](#)]

## 3 synthetic oligosaccharide

Trial	Treatments	Patients	Trials design and methods
<b>fondaparinux vs enoxaparin</b>			
PENTHIFRA (Eriksson) , 2001 n=831/840 follow-up: 11 days	fondaparinux 2.5-mg once-daily subcutaneous, starting 6 hours after surgery versus enoxaprin 40mg once daily	hip fracture surgery	Parallel groups double blind 21 countries
<b>extended prophylaxis vs standard prophylaxis</b>			
PENTHIFRAPLUS (Eriksson) , 2003 n=656 follow-up: 19-23 days	25-31 days of fondaparinux 2.5-mg once-daily versus 6-8 days of fondaparinux 2.5-mg once-daily	patients undergoing hip fracture surgery	Parallel groups double blind

## References

### PENTHIFRA (Eriksson), 2001:

Eriksson BI, Bauer KA, Lassen MR, Turpie AG Fondaparinux compared with enoxaparin for the prevention of venous thromboembolism after hip-fracture surgery. N Engl J Med 2001 Nov 1;345:1298-304 [[11794148](#)]

### PENTHIFRAPLUS (Eriksson), 2003:

Eriksson BI, Lassen MR Duration of prophylaxis against venous thromboembolism with fondaparinux after hip fracture surgery: a multicenter, randomized, placebo-controlled, double-blind study. Arch Intern Med 2003 Jun 9;163:1337-42 [[12796070](#)]

## 4 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.