

Clinical trials of antithrombotics for thrombosis prevention in patients with immobilization of the lower extremities

TrialResults-center www.trialresultscenter.org

1 Low molecular weight heparin

Trial	Treatments	Patients	Trials design and methods
certoparin vs control			
Kock , 1995 n=176/163 follow-up: 15 days	Certoparin 3000 IU versus no prophylaxis	patients with minor injuries treated with plaster-cast immobilisation of the leg	Parallel groups open
nadroparin vs control			
Kujath , 1993 n=126/126 follow-up: 65279;16 days	Nadroparin 2850 IU versus no prophylaxis	patients with injuries of the lower limb immobilized by a plaster cast	Parallel groups open
tinzaparin vs control			
Jorgensen , 2002 n=99/106 follow-up: 38 days	Tinzaparin 3500 IU versus no prophylaxis	patients over 18 years of age with planned plaster cast on a lower extremity of at least 3 weeks	Parallel groups open, assessor-blinded
nadroparin vs no treatment			
PROTECT (nadroparin) <i>ongoing</i> [NCT00881088] n=NA follow-up: 6 weeks	nadroparin 0,3 cc daily during immobilization versus no treatment	patients with a nonsurgical fracture of the lower extremity requiring immobilisation in a below-knee plaster cast	Parallel groups single blind
dalteparin vs placebo			
D-KAf (Selby) , 2007 [NCT00187408] n=134/131 follow-up:	dalteparin 5000U daily versus placebo	below-knee fractures repaired surgically	
Lapidus , 2007 n=47/44 follow-up: 43 days	Dalteparin 5000 IU versus Placebo	patients surgically treated for Achilles tendon rupture	Parallel groups double-blind
Lapidus , 2007 n=101/96 follow-up: 44 days	Dalteparin 5000 IU versus Placebo	patients undergoing ankle fracture surgery	Parallel groups double-blind
reviparin vs placebo			
Lassen , 2002 n=183/188 follow-up: 43 days	Reviparin 1750 IU versus Placebo	patients who required immobilization in a plaster cast or brace for at least five weeks after a leg fracture or rupture of the Achilles tendon	Parallel groups double-blind

References

Kock, 1995:

Kock HJ, Schmit-Neuerburg KP, Hanke J, Rudofsky G, Hirche H Thromboprophylaxis with low-molecular-weight heparin in outpatients with plaster-cast immobilisation of the leg. *Lancet* 1995;346:459-61 [[7637478](#)]

Kujath, 1993:

Kujath P, Spannagel U, Habscheid W Incidence and prophylaxis of deep venous thrombosis in outpatients with injury of the lower limb. *Haemostasis* 1993;23 Suppl 1:20-6 [[8388353](#)]

Jorgensen, 2002:

Jrgensen PS, Warming T, Hansen K, Paltved C, Vibeke Berg H, Jensen R, Kirchhoff-Jensen R, Kjaer L, Kerbouche N, Leth-Espensen P, Narvestad E, Rasmussen SW, Sloth C, Trholm C, Wille-Jrgensen P Low molecular weight heparin (Innohep) as thromboprophylaxis in outpatients with a plaster cast: a venografic controlled study. *Thromb Res* 2002;105:477-80 [[12091045](#)]

PROTECT (nadroparin), :

D-KAf (Selby), 2007:

Geerts WH, Pineo GF, Heit JA, Bergqvist D, Lassen MR, Colwell CW, Ray JG Prevention of venous thromboembolism: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004;126:338S-400S [[15383478](#)] [10.1378/chest.126.3_suppl.338S](#)

Lapidus, 2007:

Lapidus LJ, Rosfors S, Ponzer S, Levander C, Elvin A, Lrfars G, de Bri E Prolonged thromboprophylaxis with dalteparin after surgical treatment of achilles tendon rupture: a randomized, placebo-controlled study. *J Orthop Trauma* 2007;21:52-7 [[17211270](#)] [10.1097/01.bot.0000250741.65003.14](#)

Lapidus, 2007:

Lapidus LJ, Ponzer S, Elvin A, Levander C, Lrfars G, Rosfors S, de Bri E Prolonged thromboprophylaxis with Dalteparin during immobilization after ankle fracture surgery: a randomized placebo-controlled, double-blind study. *Acta Orthop* 2007;78:528-35 [[17966008](#)] [10.1080/17453670710014185](#)

Lassen, 2002:

Lassen MR, Borris LC, Nakov RL Use of the low-molecular-weight heparin reviparin to prevent deep-vein thrombosis after leg injury requiring immobilization. *N Engl J Med* 2002;347:726-30 [[12213943](#)] [10.1056/NEJMoa011327](#)

2 synthetic oligosaccharide

Trial	Treatments	Patients	Trials design and methods
fondaparinux vs no treatment			
PROTECT (fundaparinux) ongoing [NCT00881088] n=NA follow-up:	fondaparinux 2,5 mg daily group during immobilization versus no treatment	Patients with a nonsurgical fracture of the lower extremity immobilised in a below-knee plaster cast	Parallel groups single blind
fondaparinux vs nadroparin			
FONDACAST ongoing [NCT00843492] n=NA follow-up: 5 weeks	subcutaneously, once daily, fondaparinux 2.5 mg for at least 21 Days, up to complete mobilization, with a maximal duration of treatment of 45 days versus daily nadroparin 2850 anti-Xa IU (0.3 mL) for at least 21 Days, up to complete mobilization	patients requiring rigid or semi-rigid immobilization for at least 21 days and up to 45 days because of isolated non-surgical below-knee injury	Parallel groups open Europe

References

PROTECT (fundaparinux), :

FONDACAST, :

3 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.