

Clinical trials of angiotensin renin system blockade for diabetes type 2 in all type of patients

TrialResults-center www.trialresultscenter.org

1 angiotensin receptor blocker

Trial	Treatments	Patients	Trials design and methods
irbesartan vs placebo			
IDNT (irbesartan vs pbo) , 2001 n=579/569 follow-up: 2.6 years	Irbesartan 300 mg daily versus placebo	hypertensive patients with nephropathy due to type 2 diabetes	Parallel groups double blind Worldwide
IPDM (150mg) , 2001 n=195/201 follow-up: 2 years	irbesartan 150 mg daily versus placebo	hypertensive patients with type 2 diabetes and microalbuminuria	Parallel groups double-blind Worldwide
losartan vs placebo			
RENAAL , 2001 n=751/762 follow-up: 3.4 y	losartan 50 to 100 mg once daily versus placebo	patients with type 2 diabetes and nephropathy	Parallel groups double-blind America, Europe, Asia
olmesartan vs placebo			
ROADMAP , 2010 [NCT00185159] n=2232/2215 follow-up: 3.2 y	olmesartan at 40 mg/day versus placebo	patients with diabetes and at least one additional cardiovascular risk factor, but no evidence of renal dysfunction	Parallel groups double-blind Europe (19 countries)
ORIENT [NCT00141453] n=282/284 follow-up:	olmesartan versus placebo	patients with diabetic Nephropathy and overt proteinuria secondary to type 2 diabetes mellitus	Parallel groups double-blind Japan, Hong Kong
irbesartan vs amlodipine			
IDNT (irbesartan vs amlodipine) , 2001 n=579/567 follow-up: 2.6 years	Irbesartan 300 mg daily versus amlodipine 10 mg daily	hypertensive patients with nephropathy due to type 2 diabetes	Parallel groups double blind Worldwide
valsartan vs amlodipine			
NAGOYA HEART , 2011 <i>unpublished</i> [NCT00129233] n=575/575 follow-up: 3.2 y median	blood-pressure-lowering therapy based on valsartan; blood-pressure goal of <130/80 mm Hg versus blood-pressure-lowering therapy based on amlodipine; blood-pressure goal of <130/80 mm Hg	patients with hypertension with type 2 diabetes or impaired glucose tolerance	Parallel groups open Japan

References

IDNT (irbesartan vs pbo), 2001:

Lewis EJ, Hunsicker LG, Clarke WR, Berl T, Pohl MA, Lewis JB, Ritz E, Atkins RC, Rohde R, Raz I Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001;345:851-60 [[11565517](#)]

IPDM (150mg), 2001:

Parving HH, Lehnert H, Brchner-Mortensen J, Gomis R, Andersen S, Arner P The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. *N Engl J Med* 2001;345:870-8 [[11565519](#)]

RENAAL, 2001:

Brenner BM, Cooper ME, de Zeeuw D, Keane WF, Mitch WE, Parving HH, Remuzzi G, Snapinn SM, Zhang Z, Shahinfar S Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001;345:861-9 [[11565518](#)]

ROADMAP, 2010:

Ritz E, Viberti GC, Ruilope LM, Rabelink AJ, Izzo JL Jr, Katayama S, Ito S, Mimran A, Menne J, Rump LC, Januszewicz A, Haller H Determinants of urinary albumin excretion within the normal range in patients with type 2 diabetes: the Randomised Olmesartan and Diabetes Microalbuminuria Prevention (ROADMAP) study. *Diabetologia* 2010;53:49-57 [[19876613](#)] [10.1007/s00125-009-1577-3](#)

Haller H, Ito S, Izzo JL Jr, Januszewicz A, Katayama S, Menne J, Mimran A, Rabelink TJ, Ritz E, Ruilope LM, Rump LC, Viberti G Olmesartan for the delay or prevention of microalbuminuria in type 2 diabetes. *N Engl J Med* 2011 Mar 10;364:907-17 [[21388309](#)]

ORIENT, :

IDNT (irbesartan vs amlodipine), 2001:

Lewis EJ, Hunsicker LG, Clarke WR, Berl T, Pohl MA, Lewis JB, Ritz E, Atkins RC, Rohde R, Raz I Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001;345:851-60 [[11565517](#)]

NAGOYA HEART, 2011:

Matsushita K, Muramatsu T, Kondo T, Maeda K, Shintani S, Murohara T Rationale and design of the NAGOYA HEART Study: comparison between valsartan and amlodipine regarding morbidity and mortality in patients with hypertension and glucose intolerance. *J Cardiol* 2010;56:111-7 [[20409690](#)] [10.1016/j.jjcc.2010.03.004](#)

2 angiotensin-converting enzyme inhibitors

Trial	Treatments	Patients	Trials design and methods
captopril or atenolol vs control			
UKPDS 38 , 1998 n=758/390 follow-up: 8.4y (median)	tight control of blood pressure aiming at a BP <150/85 (with the use of captopril or atenolol as main treatment, other treatment were added if the control criteria were not met) versus less tight control aiming at a blood pressure of <180/105 (avoiding treatment with ACE inhibitors or beta-blockers)	hypertensive patients with type 2 diabetes	Parallel groups open UK
ACE inhibitors vs placebo			

continued...

Trial	Treatments	Patients	Trials design and methods
HOPE (diabetic subgroup) , 2000 n=1808/1759 follow-up: 4.5 years	ramipril 10 mg once per day orally versus placebo	patients with diabetes (sub group), aged 55 years or older, who had a previous cardiovascular event or at least one other cardiovascular risk factor, no clinical proteinuria, heart failure, or low ejection fraction	Factorial plan double-blind North, South america, Europe
perindopril and indapamide vs placebo			
ADVANCE , 2007 [NCT00145925] n=NA follow-up:	fixed combination of perindopril and indapamide versus placebo	patients with type 2 diabetes irrespective of initial blood pressure levels or the use of other blood pressure lowering drugs	
captopril vs atenolol			
UKPDS 39 , 1998 n=400/358 follow-up: ND	captopril 25 mg/d aiming at a BP <150/85 versus atenolol 50mg/d aiming at a BP <150/85	hypertensive patients with type 2 diabetes	Parallel groups open UK
ACE inhibitor vs calcium-channel blocker			
STOP-2 (ACEI vs CCB) (diabetic subgroup) , 2000 n=235/231 follow-up: 5.03y	ACE inhibitor versus calcium antagonists	diabetic (subgroup) elderly patients aged 70-84 years	open with blind assessment Sweden
lisinopril vs chlorthalidone			
ALLHAT (lisi vs chlor, diabetic subgroup) , 2002 n=2431/4498 follow-up: 4.9 y	lisinopril 10 to 40 mg/d versus chlorthalidone 12.5 to 25 mg/d	diabetic (subgroup) participants aged 55 years or older with hypertension	Parallel groups double-blind
captopril vs diuretic and/or beta-blockers			
CAPP (diabetic subgroup) , 1999 n=309/263 follow-up: 6.1 year	Captopril initial dose of 50 mg daily given in one or two doses versus thiazide diuretic or beta-blocker	Patients aged 25-66 years with a measured diastolic blood pressure of 100 mm Hg or more on two occasions; subgroup of diabetic patients	Parallel groups open with blinded assessment Sweden, Finland
ACE inhibitor vs diuretic or beta-blocker			
STOP-2 (ACEI, diabetic subgroup) , 2000 n=235/253 follow-up: 5.03y	ACE inhibitor versus conventional treatment (diuretic or beta-blocker)	diabetic (subgroup) elderly patients aged 70-84 years with hypertension	Parallel groups open with blind assessment Sweden

References

UKPDS 38, 1998:

Efficacy of atenolol and captopril in reducing risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 39. UK Prospective Diabetes Study Group. *BMJ* 1998;317:713-20 [9732338]

Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. UK Prospective Diabetes Study Group. *BMJ* 1998;317:703-13 [9732337]

HOPE (diabetic subgroup), 2000:

Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. Heart Outcomes Prevention Evaluation Study Investigators. *Lancet* 2000;355:253-9 [10675071]

Yusuf S, Sleight P, Pogue J, Bosch J, Davies R, Dagenais G Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. The Heart Outcomes Prevention Evaluation Study Investigators. *N Engl J Med* 2000;342:145-53 [10639539]

ADVANCE, 2007:

Rationale and design of the ADVANCE study: a randomised trial of blood pressure lowering and intensive glucose control in high-risk individuals with type 2 diabetes mellitus. Action in Diabetes and Vascular Disease: PreterAx and DiamicroN Modified-Release Controlled Evaluation. *J Hypertens Suppl* 2001 Nov;19:S21-8 [11848259]

Patel A, MacMahon S, Chalmers J, Neal B, Woodward M, Billot L, Harrap S, Poulter N, Marre M, Cooper M, Glasziou P, Grobbee DE, Hamet P, Heller S, Liu LS, Mancia G, Mogensen CE, Pan CY, Rodgers A, Williams B Effects of a fixed combination of perindopril and indapamide on macrovascular and microvascular outcomes in patients with type 2 diabetes mellitus (the ADVANCE trial): a randomised controlled trial. *Lancet* 2007 Sep 8;370:829-40 [17765963]

UKPDS 39, 1998:

Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. UK Prospective Diabetes Study Group. *BMJ* 1998;317:703-13 [9732337]

Efficacy of atenolol and captopril in reducing risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 39. UK Prospective Diabetes Study Group. *BMJ* 1998;317:713-20 [9732338]

STOP-2 (ACEI vs CCB) (diabetic subgroup), 2000:

Lindholm LH, Hansson L, Ekblom T, Dahlb B, Lanke J, Linjer E, Scherstn B, Wester PO, Hedner T, de Faire U Comparison of antihypertensive treatments in preventing cardiovascular events in elderly diabetic patients: results from the Swedish Trial in Old Patients with Hypertension-2. STOP Hypertension-2 Study Group. *J Hypertens* 2000 Nov;18:1671-5 [11081782]

Hansson L, Lindholm LH, Ekblom T, Dahlb B, Lanke J, Scherstn B, Wester PO, Hedner T, de Faire U Randomised trial of old and new antihypertensive drugs in elderly patients: cardiovascular mortality and morbidity the Swedish Trial in Old Patients with Hypertension-2 study. *Lancet* 1999;354:1751-6 [10577635]

ALLHAT (lisi vs chlor, diabetic subgroup), 2002:

Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium channel blocker vs diuretic: The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *JAMA* 2002;288:2981-97 [12479763]

Whelton PK, Barzilay J, Cushman WC, Davis BR, Iamathi E, Kostis JB, Leenen FH, Louis GT, Margolis KL, Mathis DE, Moloo J, Nwachuku C, Panebianco D, Parish DC, Pressel S, Simmons DL, Thadani U Clinical outcomes in antihypertensive treatment of type 2 diabetes, impaired fasting glucose concentration, and normoglycemia: Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *Arch Intern Med* 2005;165:1401-9 [15983290] 10.1001/archinte.165.12.1401

CAPP (diabetic subgroup), 1999:

Hansson L, Lindholm LH, Niskanen L, Lanke J, Hedner T, Niklason A, Luomanmki K, Dahlb B, de Faire U, Mrlin C, Karlberg BE, Wester PO, Bjrcck JE Effect of angiotensin-converting-enzyme inhibition compared with conventional therapy on cardiovascular morbidity and mortality in hypertension: the Captopril Prevention Project (CAPPP) randomised trial. *Lancet* 1999;353:611-6 [10030325]

Niklason A, Hedner T, Niskanen L, Lanke J Development of diabetes is retarded by ACE inhibition in hypertensive patients—a subanalysis of the Captopril Prevention Project (CAPPP). *J Hypertens* 2004;22:645-52 [15076172]

Niskanen L, Hedner T, Hansson L, Lanke J, Niklason A Reduced cardiovascular morbidity and mortality in hypertensive diabetic patients on first-line therapy with an ACE inhibitor compared with a diuretic/beta-blocker-based treatment regimen: a subanalysis of the Captopril Prevention Project. *Diabetes Care* 2001;24:2091-6 [11723089]

STOP-2 (ACEI, diabetic subgroup), 2000:

Lindholm LH, Hansson L, Ekblom T, Dahlb B, Lanke J, Linjer E, Scherstn B, Wester PO, Hedner T, de Faire U Comparison of antihypertensive treatments in preventing cardiovascular events in elderly diabetic patients: results from the Swedish Trial in Old Patients with Hypertension-2. STOP Hypertension-2 Study Group. *J Hypertens* 2000;18:1671-5 [11081782]

Hansson L, Lindholm LH, Ekblom T, Dahlb B, Lanke J, Scherstn B, Wester PO, Hedner T, de Faire U Randomised trial of old and new antihypertensive drugs in elderly patients: cardiovascular mortality and morbidity the Swedish Trial in Old Patients with Hypertension-2 study. *Lancet* 1999;354:1751-6 [10577635]

Hansson L, Hedner T, Dahlb B Prospective randomized open blinded end-point (PROBE) study. A novel design for intervention trials. Prospective Randomized Open Blinded End-Point. *Blood Press* 1992;1:113-9 [1366259]

Entry terms: candesartan, candesartan cilexetil, TCV 116, TCV-116, Atacand, Blopress, Kenzen, Amias, Parapres, candesartan, CV 11974, CV11974, CV-11974, , irbesartan, Aprovel, Avapro, Karvea, losartan, Losartan, Cozaar, MK-954, MK 954, MK954, DuP-753, DuP 753, DuP753, Losartan Potassium, Losartan Monopotassium Salt, , valsartan, valsartan, Diovan, Tareg, KalpressMiten, Provas, Vals, walsartan, CGP 48933, Nisis, Aventis brand of valsartan,

3 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.