

Clinical trials of antiplatelets drug for thrombosis prevention in elective orthopedic surgery

TrialResults-center www.trialresultscenter.org

1 platelet aggregation inhibitors

Trial	Treatments	Patients	Trials design and methods
Aspirin vs no treatment			
Pasteyer , 1977 n=20/20 follow-up: 2 weeks	Aspirin 1000mg daily + Hep versus control (Hep alone)	Elective orthopaedic surgery	Parallel groups
Rocha , 1986 n=60/30 follow-up: 1 weeks	Aspirin 250mg or 1000mg daily versus control (combination of heparin plus dihydroergotamine)	total hip replacement	Parallel groups open
aspirin + dipyridamol vs no treatment			
Lyon-I , 1975 n=20/20 follow-up: 2 weeks	Aspirin 1500 mg daily + Dipyridamole versus control	Elective orthopaedic surgery	
Aspirin vs placebo			
Stockholm-I , 1975 n=26/25 follow-up: 2 weeks	Aspirin 2000mg daily versus placebo	elective surgery of the hip	double blind
Harris-I , 1977 n=58/59 follow-up: 1 weeks	Aspirin 1200mg daily versus placebo	patients over 40 years of age, who had undergone total hip replacement	Parallel groups double-blind
McKenna-I , 1980 n=24/12 follow-up: 2 weeks	Aspirin 975mg or 3900mg daily versus placebo	total knee replacement	Parallel groups double-blind
Sautter , 1983 n=68/77 follow-up: 3 weeks	Aspirin 900mg daily + sulfipyrazone versus placebo	patient with total hip replacement	Parallel groups
McBride , 1983 n=21/22 follow-up: 1 weeks	A1800+Dipyridamole versus placebo	Elective orthopaedic surgery	
aspirin + dipyridamol vs placebo			
Hamburg , 1976 n=21/11 follow-up: 3 weeks	A+Dipyridamole,A1000 versus placebo	Elective orthopaedic surgery	
Hydroxychloroquine vs placebo			

continued...

Trial	Treatments	Patients	Trials design and methods
Massachusetts-II , 1976 n=51/51 follow-up: 3 weeks	Hydroxychloroquine 600mg daily versus placebo	fractures or orthopaedic operations involving the skeleton between the knee and the pelvis	
Cooke , 1977 n=25/25 follow-up: 2 weeks	Hydroxychloroquine versus placebo	elective surgery on the hip	Parallel groups double-blind
Hume-A , 1977 n=20/20 follow-up: 2 weeks	Hydroxychloroquine versus placebo	total hip replacement	
Stockholm-II , 1981 n=18/17 follow-up: 2 weeks	Hydroxychloroquine versus placebo	total hip replacement	
Ticlopidine vs placebo			
McKenna-II , 1983 n=29/29 follow-up: 2 weeks	Ticlopidine versus placebo	Elective orthopaedic surgery	
Lyon-II , 3000 n=20/20 follow-up: 3 weeks	Ticlopidine versus placebo	Elective orthopaedic surgery	
Gardecki , 3000 n=48/46 follow-up: 2 weeks	Ticlopidine versus placebo	Elective orthopaedic surgery	

2

References

Pasteyer, 1977:

Flicoteaux H, Kher A, Jean N, Blery M, Judet T, Honnart F, et al. Comparison of low dose heparin and low dose heparin combined with aspirin in prevention of deep vein thrombosis after total hip replacement. *Pathol Biol (Paris)* 1977;25(suppl):55-8.

Rocha, 1986:

Alfaro MJ, Pramo JA, Rocha E Prophylaxis of thromboembolic disease and platelet-related changes following total hip replacement: a comparative study of aspirin and heparin-dihydroergotamine. *Thromb Haemost* 1986;56:53-6 [3535158]

Lyon-I, 1975:

Dechavanne M, Ville D, Viala JJ, Kher A, Faivre J, Pousset MB, Dejour H Controlled trial of platelet anti-aggregating agents and subcutaneous heparin in prevention of postoperative deep vein thrombosis in high risk patients. *Haemostasis* 1975;4:94-100 [1205340]

Stockholm-I, 1975:

Soreff J, Johnsson H, Diener L, Gransson L Acetylsalicylic acid in a trial to diminish thromboembolic complications after elective hip surgery. *Acta Orthop Scand* 1975;46:246-55 [1096521]

Harris-I, 1977:

Harris WH, Salzman EW, Athanasoulis CA, Waltman AC, DeSanctis RW Aspirin prophylaxis of venous thromboembolism after total hip replacement. *N Engl J Med* 1977;297:1246-9 [335247]

McKenna-I, 1980:

McKenna R, Galante J, Bachmann F, Wallace DL, Kaushal PS, Meredith P Prevention of venous thromboembolism after total knee replacement by high-dose aspirin or intermittent calf and thigh compression. *Br Med J* 1980;280:514-7 [6989432]

Sautter, 1983:

Sautter RD, Koch EL, Myers WO, Ray JR 3rd, Mazza JJ, Larson DE, Chen HM, Milbauer JP, Treuhaf PS, Plotka ED Aspirin-sulfinpyrazone in prophylaxis of deep venous thrombosis in total hip replacement. JAMA 1983;250:2649-54 [6355542]

McBride, 1983:

McBride JA, Turpie AG, Kraus V, Hiltz C. Failure of aspirin and dipyridamole to influence the incidence of leg scan detected venous thrombosis after elective hip surgery Thrombosis et Diathesis Haemorrhagica 975;34:abstract 204.

Hamburg, 1976:

Boehringer Ingelheim DVT nach Hirntumoroperationen Boehringer Ingelheim, 1976. (Internal report.)

Massachusetts-II, 1976:

Chrisman OD, Snook GA, Wilson TC, Short JY Prevention of venous thromboembolism by administration of hydroxychloroquine. A preliminary report. J Bone Joint Surg Am 1976;58:918-20 [789380]

Cooke, 1977:

Cooke ED, Dawson MH, Ibbotson RM, Bowcock SA, Ainsworth ME, Pilcher MF Failure of orally administered hydroxychloroquine sulphate to prevent venous thromboembolism following elective hip operations. J Bone Joint Surg Am 1977;59:496-500 [325009]

Hume-A, 1977:

Hume M, Bierbaum B, Kuriakose TX, Surprenant, J Prevention of postoperative thrombosis by aspirin. Am J Surg 1977;133:420-2 [322520]

Hume M, Donaldson WR, Surprenant J Sex, aspirin, and venous thrombosis. Orthop Clin North Am 1978;9:761-7 [358040]

Stockholm-II, 1981:

Johansson E, Forsberg K, Johnsson H Clinical and experimental evaluation of the thromboprophylactic effect of hydroxychloroquine sulfate after total hip replacement. Haemostasis 1981;10:89-96 [7007179]

McKenna-II, 1983:

McKenna R, Galante J, Molony B, Kamm B. Failure of ticlopidine hydrochloride to prevent DVT in orthopedic patients Blood 1983;62(suppl 1):304A

Lyon-II, 3000:**Gardecki, 3000:**

Entry terms: ticlopidine, Ticlopidine, Ticlopidine Hydrochloride, Ticlodix, Ticlodone, 53-32C, 53 32C, 5332C, Ticlid,

2 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.