

Clinical trials of mechanical devices for thromboprophylaxis for DVT prophylaxis in general non orthopaedic surgery

TrialResults-center www.trialresultscenter.org

1 GCS (adjunctive therapy)

Trial	Treatments	Patients	Trials design and methods
GCS + dextran vs dextran			
*Bergqvist , 1984 n=80/80 follow-up:	one leg encased in a graded compression stocking versus other leg unstockinged	patients undergoing Abdominal surgery	
GCS +IPC vs IPC			
Mellbring , 1986 n=NA follow-up:	graduated static compression + peroperative intermittent pneumatic calf compression versus peroperative intermittent pneumatic calf compression	patients undergoing major abdominal surgery	Factorial plan
Scurr (GCS+IPC vs IPC) , 1987 n=78/78 follow-up:	simultaneous use of graduated compression stockings and intermittent sequential pneumatic compression versus intermittent sequential pneumatic compression	patients undergoing Abdominal surgery	parallel groups
GCS + UFH vs UFH			
*Rasmussen (adj) , 1988 n=NA follow-up:	subcutaneous heparin and graduated compression stockings to the knee (TED stockings), versus subcutaneous heparin	patients (age more than 40 yrs) admitted for major abdominal surgery	Parallel groups open
Wille-Jorgensen , 1985 n=86/90 follow-up:	low-dose heparin treatment with graded compression stockings versus low-dose heparin treatment (5000 units twice daily subcutaneously)	patients undergoing Abdominal surgery	parallel groups
Wille-Jorgensen , 1991 n=83/83 follow-up:	low dose heparin and graded compression stockings versus low dose heparin	patients undergoing Abdominal surgery	parallel groups

References

*Bergqvist, 1984:

Bergqvist D, Lindblad B The thromboprophylactic effect of graded elastic compression stockings in combination with dextran 70. Arch Surg 1984;119:1329-31 [6208877]

Mellbring, 1986:

Mellbring G, Palmr K Prophylaxis of deep vein thrombosis after major abdominal surgery. Comparison between dihydroergotamine-heparin and intermittent pneumatic calf compression and evaluation of added graduated static compression. Acta Chir Scand 1986;152:597-600 [3544626]

Scurr (GCS+IPC vs IPC), 1987:

Scurr JH, Coleridge-Smith PD, Hasty JH Regimen for improved effectiveness of intermittent pneumatic compression in deep venous thrombosis prophylaxis. Surgery 1987;102:816-20 [3672322]

***Rasmussen (adj), 1988:**

Rasmussen A, Hansen PT, Lindholt J, Poulsen TD, Toftdahl DB, Gram J, Toftgaard C, Jespersen J Venous thrombosis after abdominal surgery. A comparison between subcutaneous heparin and antithrombotic stockings, or both. J Med 1988;19:193-201 [2972790]

Wille-Jorgensen, 1985:

Wille-Jrgensen P, Thorup J, Fischer A, Holst-Christensen J, Flamsholt R Heparin with and without graded compression stockings in the prevention of thromboembolic complications of major abdominal surgery: a randomized trial. Br J Surg 1985;72:579-81 [4016545]

Wille-Jorgensen, 1991:

Wille-Jrgensen P, Hauch O, Dimo B, Christensen SW, Jensen R, Hansen B Prophylaxis of deep venous thrombosis after acute abdominal operation. Surg Gynecol Obstet 1991;172:44-8 [1702235]

2 Graduated compression stockings

2

Trial	Treatments	Patients	Trials design and methods
GCS vs no prophylaxis			
*Inada , 1983 n=NA follow-up:	graduated compression stocking on one leg versus no GCS on the other leg serving as a control.	patients undergoing major surgery	
*Scurr , 1977 n=70/70 follow-up:	graduated static compression stockings on one leg versus other leg being used as a control	patients undergoing 65279;Abdominal surgery	open
Turner , 1984 n=104/92 follow-up:	graduated compression stockings versus not wear the stockings	patients undergoing Gynecologic surgery	parallel groups
Allan , 1983 n=97/103 follow-up:	graduated compression stockings versus control	patients undergoing Abdominal surgery	parallel groups
Turpie (GCS vs ctrl) , 1989 n=80/81 follow-up:	graduated compression stockings versus untreated control	patients undergoing Neurosurgery surgery	
Holford , 1976 n=48/47 follow-up:	Graded compression versus control	patients undergoing major operations	parallel groups
Knee length GCS vs Thigh length GCS			

continued...

Trial	Treatments	Patients	Trials design and methods
Porteous , 1989 n=58/58 follow-up:	Knee length GCS versus Thigh length GCS	65279;General surgical patients	parallel groups
Williams , 1988 n=44/44 follow-up:	Knee length GCS versus Thigh length GCS	General surgical patients	parallel groups
Howard , 2004 n=99/195 follow-up:	Knee length GCS versus Thigh length GCS	Breast surgery, oncology, ENT, urology, vascular, neurosurgery. And gastrointestinal surgery	parallel groups
GCS vs UFH			
Hansberry (vs UFH) , 1991 n=NA follow-up: 6 days	thromboembolic stockings versus heparin plus dihydroergotamine	patients undergoing a major urological operation	Parallel groups
Rasmussen (GCS vs UFH) , 1998 n=74/85 follow-up:	graduated compression stockings to the knee (TED stockings) versus subcutaneous heparin	patients (age more than 40 yrs) admitted for major abdominal surgery	Parallel groups open

References

*Inada, 1983:

Inada K, Shirai N, Hayashi M, Matsumoto K, Hirose M Postoperative deep venous thrombosis in Japan. Incidence and prophylaxis. Am J Surg 1983;145:775-9 [6859416]

*Scurr, 1977:

Scurr JH, Ibrahim SZ, Faber RG, Le Quesne LP The efficacy of graduated compression stockings in the prevention of deep vein thrombosis. Br J Surg 1977;64:371-3 [861482]

Turner, 1984:

Turner GM, Cole SE, Brooks JH The efficacy of graduated compression stockings in the prevention of deep vein thrombosis after major gynaecological surgery. Br J Obstet Gynaecol 1984;91:588-91 [6733063]

Allan, 1983:

Allan A, Williams JT, Bolton JP, Le Quesne LP The use of graduated compression stockings in the prevention of postoperative deep vein thrombosis. Br J Surg 1983;70:172-4 [6338992]

Turpie (GCS vs crt1), 1989:

Turpie AG, Hirsh J, Gent M, Julian D, Johnson J Prevention of deep vein thrombosis in potential neurosurgical patients. A randomized trial comparing graduated compression stockings alone or graduated compression stockings plus intermittent pneumatic compression with control. Arch Intern Med 1989;149:679-81 [2645846]

Holford, 1976:

Holford CP Graded compression for preventing deep venous thrombosis. Br Med J 1976;2:969-70 [788855]

Porteous, 1989:

Porteous MJ, Nicholson EA, Morris LT, James R, Negus D Thigh length versus knee length stockings in the prevention of deep vein thrombosis. Br J Surg 1989;76:296-7 [2720328]

Williams, 1988:

Williams JT, Palfrey SM Cost effectiveness and efficacy of below knee against above knee graduated compression stockings in the prevention of deep vein thrombosis. Phlebologie 1988;41:809-11 [3150063]

Howard, 2004:

Howard A, Zaccagnini D, Ellis M, Williams A, Davies AH, Greenhalgh RM Randomized clinical trial of low molecular weight heparin with thigh-length or knee-length antiembolism stockings for patients undergoing surgery. Br J Surg 2004;91:842-7 [15227689]

Hansberry (vs UFH), 1991:

Hansberry KL, Thompson IM Jr, Bauman J, Deppe S, Rodriguez FR A prospective comparison of thromboembolic stockings, external sequential pneumatic compression stockings and heparin sodium/dihydroergotamine mesylate for the prevention of thromboembolic complications in urological surgery. J Urol 1991;145:1205-8 [2033693]

Rasmussen (GCS vs UFH), 1998:

Rasmussen A, Hansen PT, Lindholt J, Poulsen TD, Toftdahl DB, Gram J, Toftgaard C, Jespersen J Venous thrombosis after abdominal surgery. A comparison between subcutaneous heparin and antithrombotic stockings, or both. J Med 1988;19:193-201 [2972790]

3 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.