

# Clinical trials of monoclonal antibody for early breast cancer in primary operable breast cancer

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 aaa

Trial	Treatments	Patients	Trials design and methods
<b>trastuzumab during chemotherapy vs control</b>			
NSABP B-31 n=864/872 follow-up: 2.9 years	same regimen plus 52 weeks of trastuzumab beginning with the first dose of paclitaxel versus doxorubicin and cyclophosphamide followed by paclitaxel every 3 weeks	Early-stage, node-positive invasive breast cancer	

## References

### NSABP B-31, :

Romond EH, Perez EA, Bryant J, Suman VJ, Geyer CE Jr, Davidson NE, Tan-Chiu E, Martino S, Paik S, Kaufman PA, Swain SM, Pisansky TM, Fehrenbacher L, Kutteh LA, Vogel VG, Visscher DW, Yothers G, Jenkins RB, Brown AM, Dakhil SR, Mamounas EP, Lingle WL, Klei Trastuzumab plus adjuvant chemotherapy for operable HER2-positive breast cancer. *N Engl J Med* 2005;353:1673-84 [[16236738](#)] [10.1056/NEJMoa052122](#)

## 2 after chemotherapy

Trial	Treatments	Patients	Trials design and methods
<b>trastuzumab vs</b>			
HERA (BIG 01-01) 2-years , 2005 [NCT00045032] n=NA follow-up: 2 years	two years of trastuzumab given every three weeks versus observation	patients with HER2-positive and either node-negative or node-positive breast cancer who had completed locoregional therapy and at least four cycles of neoadjuvant or adjuvant chemotherapy	Parallel groups open
<b>trastuzumab vs control</b>			
HERA (BIG 01-01) 1-year , 2005 [NCT00045032] n=1694/1693 follow-up: 2 years	one years of trastuzumab given every three weeks versus observation	patients with HER2-positive and either node-negative or node-positive breast cancer who had completed locoregional therapy and at least four cycles of neoadjuvant or adjuvant chemotherapy	Parallel groups open

continued...

Trial	Treatments	Patients	Trials design and methods
NCCTG N9831(C vs A) n=808/807 follow-up: 2.9 years	same regimen plus 52 weeks of trastuzumab initiated concomitantly with paclitaxel versus doxorubicin and cyclophosphamide followed by weekly paclitaxel	Early-stage invasive breast cancer, node positive or high-risk node negative (tumor 1 cm if HR negative or 2 cm if HR positive)	

## References

### HERA (BIG 01-01) 2-years, 2005:

Smith I, Procter M, Gelber RD, Guillaume S, Feyereislova A, Dowsett M, Goldhirsch A, Untch M, Mariani G, Baselga J, Kaufmann M, Cameron D, Bell R, Bergh J, Coleman R, Wardley A, Harbeck N, Lopez RI, Mallmann P, Gelmon K, Wilcken N, Wist E, Snchez Rovira 2-year follow-up of trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer: a randomised controlled trial. *Lancet* 2007;369:29-36 [17208639] [10.1016/S0140-6736\(07\)60028-2](https://doi.org/10.1016/S0140-6736(07)60028-2)

Cameron D, Piccart-Gebhart MJ, Gelber RD, Procter M, Goldhirsch A, de Azambuja E, Castro G Jr, Untch M, Smith I, Gianni L, Baselga J, Al-Sakaff N, Lau 11 years' follow-up of trastuzumab after adjuvant chemotherapy in HER2-positive early breast cancer: final analysis of the HERceptin Adjuvant (HERA) trial. *Lancet* 2017;: [28215665]

Cameron D, Piccart-Gebhart MJ, Gelber RD, Procter M, Goldhirsch A, de Azambuja E, Castro G Jr, Untch M, Smith I, Gianni L, Baselga J, Al-Sakaff N, Lau 11 years' follow-up of trastuzumab after adjuvant chemotherapy in HER2-positive early breast cancer: final analysis of the HERceptin Adjuvant (HERA) trial. *Lancet* 2017;: [28215665]

### HERA (BIG 01-01) 1-year, 2005:

Smith I, Procter M, Gelber RD, Guillaume S, Feyereislova A, Dowsett M, Goldhirsch A, Untch M, Mariani G, Baselga J, Kaufmann M, Cameron D, Bell R, Bergh J, Coleman R, Wardley A, Harbeck N, Lopez RI, Mallmann P, Gelmon K, Wilcken N, Wist E, Snchez Rovira 2-year follow-up of trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer: a randomised controlled trial. *Lancet* 2007;369:29-36 [17208639] [10.1016/S0140-6736\(07\)60028-2](https://doi.org/10.1016/S0140-6736(07)60028-2)

Piccart-Gebhart MJ, Procter M, Leyland-Jones B, Goldhirsch A, Untch M, Smith I, Gianni L, Baselga J, Bell R, Jackisch C, Cameron D, Dowsett M, Barrios CH, Steger G, Huang CS, Andersson M, Inbar M, Lichinitser M, Lng I, Nitz U, Iwata H, Thomssen C, Lohris Trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer. *N Engl J Med* 2005;353:1659-72 [16236737] [10.1056/NEJMoa052306](https://doi.org/10.1056/NEJMoa052306)

### NCCTG N9831(C vs A), :

Romond EH, Perez EA, Bryant J, Suman VJ, Geyer CE Jr, Davidson NE, Tan-Chiu E, Martino S, Paik S, Kaufman PA, Swain SM, Pisansky TM, Fehrenbacher L, Kutteh LA, Vogel VG, Visscher DW, Yothers G, Jenkins RB, Brown AM, Dakhil SR, Mamounas EP, Lingle WL, Klei Trastuzumab plus adjuvant chemotherapy for operable HER2-positive breast cancer. *N Engl J Med* 2005;353:1673-84 [16236738] [10.1056/NEJMoa052122](https://doi.org/10.1056/NEJMoa052122)

## 3 during chemotherapy

Trial	Treatments	Patients	Trials design and methods
<b>trastuzumab vs control</b>			
HERA (BIG 01-01) 1-year , 2005 [NCT00045032] n=1694/1693 follow-up: 2 years	one years of trastuzumab given every three weeks versus observation	patients with HER2-positive and either node-negative or node-positive breast cancer who had completed locoregional therapy and at least four cycles of neoadjuvant or adjuvant chemotherapy	Parallel groups open

continued...

Trial	Treatments	Patients	Trials design and methods
NCCTG N9831(C vs A) n=808/807 follow-up: 2.9 years	same regimen plus 52 weeks of trastuzumab initiated concomitantly with paclitaxel versus doxorubicin and cyclophosphamide followed by weekly paclitaxel	Early-stage invasive breast cancer, node positive or high-risk node negative (tumor 1 cm if HR negative or 2 cm if HR positive)	

## References

### HERA (BIG 01-01) 1-year, 2005:

Smith I, Procter M, Gelber RD, Guillaume S, Feyereislova A, Dowsett M, Goldhirsch A, Untch M, Mariani G, Baselga J, Kaufmann M, Cameron D, Bell R, Bergh J, Coleman R, Wardley A, Harbeck N, Lopez RI, Mallmann P, Gelmon K, Wilcken N, Wist E, Snchez Rovira 2-year follow-up of trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer: a randomised controlled trial. *Lancet* 2007;369:29-36 [[17208639](#)] [10.1016/S0140-6736\(07\)60028-2](#)

Piccart-Gebhart MJ, Procter M, Leyland-Jones B, Goldhirsch A, Untch M, Smith I, Gianni L, Baselga J, Bell R, Jackisch C, Cameron D, Dowsett M, Barrios CH, Steger G, Huang CS, Andersson M, Inbar M, Lichinitser M, Lng I, Nitz U, Iwata H, Thomssen C, Lohris Trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer. *N Engl J Med* 2005;353:1659-72 [[16236737](#)] [10.1056/NEJMoa052306](#)

### NCCTG N9831(C vs A), :

Romond EH, Perez EA, Bryant J, Suman VJ, Geyer CE Jr, Davidson NE, Tan-Chiu E, Martino S, Paik S, Kaufman PA, Swain SM, Pisansky TM, Fehrenbacher L, Kutteh LA, Vogel VG, Visscher DW, Yothers G, Jenkins RB, Brown AM, Dakhil SR, Mamounas EP, Lingle WL, Klei Trastuzumab plus adjuvant chemotherapy for operable HER2-positive breast cancer. *N Engl J Med* 2005;353:1673-84 [[16236738](#)] [10.1056/NEJMoa052122](#)

3

## 4 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.