

Clinical trials of intensive blood pressure control for hypertension in patients with chronic kidney disease

TrialResults-center www.trialresultscenter.org

1 Treatment blood pressure target

Trial	Treatments	Patients	Trials design and methods
more intensive blood pressure lowering strategie vs less intensive blood pressure lowering strategie			
AASK , 2002 n=540/554 follow-up: (range 3-6.4y)	arterial pressure goal of 92 mm Hg or lower versus usual mean arterial pressure goal of 102 to 107 mm Hg/pj	African-Americans,with diastolic blood pressure higher than 94mmHg and a glomerular filtration rate between 20 and 65 ml/min per 1.73 m2	Parallel groups open USA
REIN-2 , 2005 n=169/169 follow-up: 36 months	intensified (systolic/diastolic <130/80 mm Hg) blood-pressure control versus conventional (diastolic <90 mm Hg) blood-pressure control	patients with non-diabetic proteinuric nephropathies receiving background treatment with the ACE inhibitor ramipril	open
MDRD , 1994 n=840 follow-up: 2.2 y	low target blood pressure (mean arterial pressure <92 mm Hg) versus usual target blood pressure (mean arterial pressure <107 mm Hg)	patients with predominantly nondiabetic kidney disease and a glomerular filtration rate of 13 to 55 mL/min per 1.73 m2	open
Toto , 1995 n=42/35 follow-up:	strict blood pressure control (DBP 65 to 80 mm Hg) versus usual blood pressure control (DBP 85 to 95 mm Hg)	non-diabetic patients (age 25 to 73) with long-standing hypertension (DBP \geq 95 mm Hg), chronic renal insufficiency (GFR \leq 70 mL/min/1.73 m2) and a normal urine sediment	open

References

AASK, 2002:

Wright JT Jr, Bakris G, Greene T, Agodoa LY, Appel LJ, Charleston J, Cheek D, Douglas-Baltimore JG, Gassman J, Glassock R, Hebert L, Jamerson K, Lewis J, Phillips RA, Toto RD, Middleton JP, Rostand SG Effect of blood pressure lowering and antihypertensive drug class on progression of hypertensive kidney disease: results from the AASK trial. *JAMA* 2002;288:2421-31 [[12435255](#)]

REIN-2, 2005:

Ruggenti P, Perna A, Loriga G, Ganeva M, Ene-Iordache B, Turturro M, Lesti M, Perticucci E, Chakarski IN, Leonardis D, Garini G, Sessa A, Basile C, Alpa M, Scanziani R, Sorba G, Zoccali C, Remuzzi G Blood-pressure control for renoprotection in patients with non-diabetic chronic renal disease (REIN-2): multicentre, randomised controlled trial. *Lancet* 2005;365:939-46 [[15766995](#)] [10.1016/S0140-6736\(05\)71082-5](#)

Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. UK Prospective Diabetes Study Group. *BMJ* 1998;317:703-13 [[9732337](#)]

MDRD, 1994:

Klahr S, Levey AS, Beck GJ, Caggiula AW, Hunsicker L, Kusek JW, Striker G The effects of dietary protein restriction and blood-pressure control on the progression of chronic renal disease. Modification of Diet in Renal Disease Study Group. N Engl J Med 1994;330:877-84 [8114857]

Lazarus JM, Bourgoignie JJ, Buckalew VM, Greene T, Levey AS, Milas NC, Paronandi L, Peterson JC, Porush JG, Rauch S, Soucie JM, Stollar C Achievement and safety of a low blood pressure goal in chronic renal disease. The Modification of Diet in Renal Disease Study Group. Hypertension 1997;29:641-50 [9040451]

Peterson JC, Adler S, Burkart JM, Greene T, Hebert LA, Hunsicker LG, King AJ, Klahr S, Massry SG, Seifter JL Blood pressure control, proteinuria, and the progression of renal disease. The Modification of Diet in Renal Disease Study. Ann Intern Med 1995;123:754-62 [7574193]

Sarnak MJ, Greene T, Wang X, Beck G, Kusek JW, Collins AJ, Levey AS The effect of a lower target blood pressure on the progression of kidney disease: long-term follow-up of the modification of diet in renal disease study. Ann Intern Med 2005;142:342-51 [15738453]

Toto, 1995:

Toto RD, Mitchell HC, Smith RD, Lee HC, McIntire D, Pettinger WA "Strict" blood pressure control and progression of renal disease in hypertensive nephrosclerosis. Kidney Int 1995;48:851-9 [7474675]

2 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.