

Clinical trials of HDL increasing drugs for cardiovascular prevention in all type of patients

TrialResults-center www.trialresultscenter.org

1 CETP inhibitor

Trial	Treatments	Patients	Trials design and methods
dalcetrapib vs			
NCT01516541 <i>ongoing</i> [NCT01516541] n=2220 follow-up:	dalcetrapib 600 mg orally daily versus placebo	patients with stable coronary heart disease (CHD), with CHD risk equivalents or at elevated risk for cardiovascular disease	
NCT01059682 <i>ongoing</i> [NCT01059682] n=936 follow-up:	dalcetrapib 600 mg orally once a day versus placebo	subjects undergoing coronary angiography who have coronary artery disease	
anacetrapib vs placebo			
REVEAL HPS-3 TIMI-55 , 2017 [NCT01252953] n=30624 follow-up: median 4 years	anacetrapib 100mg daily versus placebo	high risk patients already taking statins	Parallel groups double-blind
REALIZE , 2015 [NCT01524289] n=204/102 follow-up: 52 weeks	oral anacetrapib 100 mg for 52 weeks versus placebo	patients aged 18-80 years with a genotype-confirmed or clinical diagnosis of heterozygous familial hypercholesterolaemia, on optimum lipid-lowering treatment for at least 6 weeks, and with an LDL-C concentration of 259 mmol/L or higher without cardiovascular disease or 181 mmol/L or higher with cardiovascular disease	Parallel groups double-blind
DEFINE , 2010 [NCT00685776] n=811/812 follow-up:	anacetrapib 100mg fr 18 months versus placebo	patients with coronary heart disease or at high risk for coronary heart disease	Parallel groups double-blind 20 countries
dalcetrapib vs placebo			
dal-VESSEL , 2011 n=NA follow-up: 12 weeks	dalcetrapib 600 mg daily versus placebo	men and women with coronary heart disease or coronary heart disease risk equivalents with HDL-cholesterol levels <50 mg/dL	Parallel groups double-blind
dal-OUTCOMES , 2012 [NCT00658515] n=7938/7933 follow-up: 31 montsh (median)	dalcetrapib 600 mg daily beginning 4 to 12 weeks after an index ACS event versus placebo	patients with recent acute coronary syndrome	Parallel groups double-blind 27 countries

continued...

Trial	Treatments	Patients	Trials design and methods
evacetrapib vs placebo			
ACCELERATE , 2017 [NCT01687998] n=6038/6054 follow-up:	evacetrapib at adose of 130 mg versus placebo	Patients at a High-Risk for Vascular Outcomes who had at least one of the following conditions: an acutecoronary syndrome within the previous 30 to 365 days, cerebrovascular atheroscleroticdisease, peripheral vascular arterial disease, or diabetes mellitus with coronaryartery disease	Parallel groups double-blind 37 countries
torcetrapib vs placebo			
RADIANCE 1 , 2007 [NCT00136981] n=450/454 follow-up: 24 months	atorvastatin combined with 60 mg of torcetrapib versus atorvastatin monotherapy	patients with heterozygous familial hypercholesterolemia	Parallel groups open
ILLUMINATE , 2007 [NCT00134264] n=7533/7534 follow-up: 1.52y	torcetrapib 60mg daily plus atorvastatin (at a dose established during the runinperiod) versus atorvastatin alone	patients at highcardiovascular risk	Parallel groups double blind 7 countries
RADIANCE 2 , 2007 n=377/375 follow-up: 24 months	torcetrapib 60mg daily (on top of atorvastatin attitrated dose) versus placebo +atorvastatin attitrated dose	patients with mixed dyslipidaemia	Parallel groups double blind North America and Europe
ILLUSTRATE , 2007 [NCT00134173] n=591/597 follow-up: 24 months	atorvastatin plus 60 mg of torcetrapib daily versus atorvastatin monotherapy	patients with coronary disease	Parallel groups open North America and Europe

References

NCT01516541, :

NCT01059682, :

REVEAL HPS-3 TIMI-55, 2017:

Effects of Anacetrapib in Patients with Atherosclerotic Vascular Disease. N Engl J Med 2017;; [28847206]

REALIZE, 2015:

Kastelein JJ, Besseling J, Shah S, Bergeron J, Langslet G, Hovingh GK, Al-Saady N, Koeijvoets M, Hunter J, Johnson-Levonas AO, Fable J, Sapre A, Mitchel Y Anacetrapib as lipid-modifying therapy in patients with heterozygous familial hypercholesterolaemia (REALIZE): a randomised, double-blind, placebo-controlled, phase 3 study. Lancet 2015 Mar 2;; [25743173] 10.1016/S0140-6736(14)62115-2

DEFINE, 2010:

Cannon CP, Shah S, Dansky HM, Davidson M, Brinton EA, Gotto AM, Stepanavage M, Liu SX, Gibbons P, Ashraf TB, Zafarino J, Mitchel Y, Barter P Safety of Anacetrapib in Patients with or at High Risk for Coronary Heart Disease. N Engl J Med 2010 Nov 17;; [21082868] 10.1056/NEJMoa1009744

dal-VESSEL, 2011:

dal-OUTCOMES, 2012:

Schwartz GG, Olsson AG, Ballantyne CM, Barter PJ, Holme IM, Kallend D, Leiter LA, Leitersdorf E, McMurray JJ, Shah PK, Tardif JC, Chaitman BR, Duttlinger-Maddux R, Mathieson J Rationale and design of the dal-OUTCOMES trial: efficacy and safety of dalcetrapib in patients with recent acute coronary syndrome. Am Heart J 2009 Dec;158:896-901.e3 [19958854]

Schwartz GG, Olsson AG, Abt M, Ballantyne CM, Barter PJ, Brumm J, Chaitman BR, Holme IM, Kallend D, Leiter LA, Leitersdorf E, McMurray JJ, Mundl H, Nicholls SJ, Shah PK, Tardif JC, Wright RS Effects of Dalcetrapib in Patients with a Recent Acute Coronary Syndrome. N Engl J Med 2012 Nov 5;: [23126252] 10.1056/NEJMoa1206797

ACCELERATE, 2017:

Lincoff AM, Nicholls SJ, Riesmeyer JS, Barter PJ, Brewer HB, Fox KAA, Gibson CM, Granger C, Menon V, Montalescot G, Rader D, Tall AR, McErlean E, Wolski K, Ruotolo G, Vangerow B, Weerakkody G, Goodman SG, Conde D, McGuire DK, Nicolau JC, Leiva-Pons JL, Pe Evacetrapib and Cardiovascular Outcomes in High-Risk Vascular Disease. N Engl J Med 2017;376:1933-1942 [28514624]

RADIANCE 1, 2007:

Kastelein JJ, van Leuven SI, Burgess L, Evans GW, Kuivenhoven JA, Barter PJ, Revkin JH, Grobbee DE, Riley WA, Shear CL, Duggan WT, Bots ML Effect of torcetrapib on carotid atherosclerosis in familial hypercholesterolemia. N Engl J Med 2007 Apr 19;356:1620-30 [17387131] 10.1056/NEJMoa071359

ILLUMINATE, 2007:

Barter PJ, Caulfield M, Eriksson M, Grundy SM, Kastelein JJ, Komajda M, Lopez-Sendon J, Mosca L, Tardif JC, Waters DD, Shear CL, Revkin JH, Buhr KA, Fisher MR, Tall AR, Brewer B Effects of torcetrapib in patients at high risk for coronary events. N Engl J Med 2007 Nov 22;357:2109-22 [17984165] 10.1056/NEJMoa0706628

RADIANCE 2, 2007:

Bots ML, Visseren FL, Evans GW, Riley WA, Revkin JH, Tegeler CH, Shear CL, Duggan WT, Vicari RM, Grobbee DE, Kastelein JJ Torcetrapib and carotid intima-media thickness in mixed dyslipidaemia (RADIANCE 2 study): a randomised, double-blind trial. Lancet 2007;370:153-60 [17630038]

ILLUSTRATE, 2007:

Nissen SE, Tardif JC, Nicholls SJ, Revkin JH, Shear CL, Duggan WT, Ruzyllo W, Bachinsky WB, Lasala GP, Tuzcu EM Effect of torcetrapib on the progression of coronary atherosclerosis. N Engl J Med 2007;356:1304-16 [17387129]

2 fibrates

3

Trial	Treatments	Patients	Trials design and methods
bezafibrate vs placebo			
BECAIT , 1996 n=47/45 follow-up: 5.0 years	bezafibrate 200 mg three times daily versus placebo	dyslipidaemic male survivors of myocardial infarction who were younger than 45 years at the time of the event	Parallel groups double blind Sweden
BIP , 2000 n=1548/1542 follow-up: 6.2 y	bezafibrate 400 mg/d versus placebo	patients with a previous myocardial infarction or stable angina, total cholesterol of 180 to 250 mg/dL, HDL-C <or =45 mg/dL, triglycerides <or =300 mg/dL, and low-density lipoprotein cholesterol <or =180 mg/dL	Parallel groups double blind Israel
LEADER , 2002 n=783/785 follow-up: 4.6y	bezafibrate 400 mg daily versus placebo	men with lower extremity arterial disease	Parallel groups double-blind UK
SENDCAP , 1998 n=81/83 follow-up: 3.0 years	bezafibrate 400 mg daily versus placebo	type 2 diabetic subjects without a history of clinical cardiovascular	Parallel groups double blind UK
clofibrate vs placebo			

continued...

Trial	Treatments	Patients	Trials design and methods
Acheson , 1972 n=NA follow-up: 6 years	clofibrate versus placebo	cerebral vascular disease	Parallel groups NA UK
Begg , 1971 n=76/79 follow-up: 3.5 y	clofibrate versus placebo	peripheral arteriopathy	Parallel groups
CDP Clofibrate , 1975 n=1103/2789 follow-up: 6.2 years	clofibrate 1.8 mg/d versus placebo	men, 30-64 y	Parallel groups double blind USA
Cullen , 1974 n=20/20 follow-up: 2 years	clofibrate versus placebo		Parallel groups
Hanefeld , 1991 n=379/382 follow-up: 5 years	clofibrilic acid 1.6 g/day versus placebo	newly diagnosed middle-aged (30- to 55-yr-old) patients with non-insulin-dependent diabetes mellitus	Parallel groups double-blind Germany
Harrold , 1969 n=30/33 follow-up: 1 years	clofibrate versus placebo	diabetic retinopathy	Parallel groups double-blind
Newcastle , 1971 n=244/253 follow-up: 3.6 y	clofibrate 1.5-2 g daily versus placebo	Hommes et femmes <65 ans	Parallel groups double blind UK
Scottish , 1971 n=350/367 follow-up: 3.4 years	clofibrate 1.6-2 g daily versus placebo	Hommes et femmes, de 40 69 ans	Parallel groups double blind Scotland
VA Neurology Section , 1974 n=268/264 follow-up: 1.8 years	clofibrate versus placebo	treatment of cerebrovascular disease	Parallel groups USA
WHO clofibrate , 1978 n=5331/5296 follow-up: 5.3 years	clofibrate 1.6 g daily versus olive oil	primary prevention, Hommes, de 30 59 ans	Parallel groups double blind Scotland, Hungary, Czech Republic
etofibrate vs placebo			
Emmerich , 2009 n=NA follow-up: 12 months	etofibrate 1g/j versus placebo	patients with type 2 diabetes mellitus and concomitant diabetic retinopathy	Parallel groups double-blind Germany
fenofibrate vs placebo			
DAIS , 2001 n=207/211 follow-up: 3.3 years	fenofibrate 200 mg/day versus placebo	men and women with type 2 diabetes and coronary atherosclerosis	Parallel groups double-blind Canada, Finland, France, Sweden

continued...

Trial	Treatments	Patients	Trials design and methods
FIELD , 2005 [ISRCTN64783481] n=4895/4900 follow-up: 5 years	fenofibrate 200mg/d versus Placebo	participants aged 50-75 years, with type 2 diabetes mellitus, and not taking statin therapy at study entry	Parallel groups double blind Australia, New Zealand, Finland
gemfibrozil vs placebo			
Helsinki (HHS) , 1987 n=2046/2035 follow-up: 5 years	gemfibrozil 1,2 g/d versus placebo	asymptomatic middle-aged men (40 to 55 years of age) with primary dyslipidemia (non-HDL cholesterol greater than or equal to 200 mg per deciliter [5.2 mmol per liter])	Parallel groups double blind Finland
HHS (Frick)(secondary prev subgroup) , 1993 n=311/317 follow-up: 5.0 years	gemfibrozil 600 mg twice daily versus placebo	individuals who exhibited symptoms and signs of possible coronary heart disease	Parallel groups double blind Sweden
LOCAT , 1997 n=197/198 follow-up: 32 months	gemfibrozil 1200 mg/d versus placebo	post-coronary bypass men, who had an HDL cholesterol concentration \leq 1.1 mmol/L and LDL cholesterol \leq 4.5 mmol/L	Parallel groups double blind Germany
VA-HIT , 1999 [NCT00283335] n=1264/1267 follow-up: 5.1 years	gemfibrozil 1.2g daily versus placebo	men with coronary heart disease, an HDL cholesterol level of 40 mg per deciliter (1.0 mmol per liter) or less, and an LDL cholesterol level of 140 mg per deciliter (3.6 mmol per liter) or less	Parallel groups double blind USA
fenofibrate vs placebo (on top simvastatine)			
ACCORD lipid , 2010 [NCT00000620] n=2765/2753 follow-up: 4.7y	fenofibrate on top simvastatin versus placebo (on top simvastatine)	high-risk patients with type 2 diabetes	Factorial plan double-blind United States and Canada

References

BECAIT, 1996:

Ruotolo G, Ericsson CG, Tettamanti C, Karpe F, Grip L, Svane B, Nilsson J, de Faire U, Hamsten A Treatment effects on serum lipoprotein lipids, apolipoproteins and low density lipoprotein particle size and relationships of lipoprotein variables to progression of coronary artery disease in the Bezafibrate Coronary Atherosclerosis Intervention Trial (BECAIT). J Am Coll Cardiol 1998;32:1648-56 [9822092]

Ericsson CG Results of the Bezafibrate Coronary Atherosclerosis Intervention Trial (BECAIT) and an update on trials now in progress. Eur Heart J 1998;19 Suppl H:H37-41 [9717064]

Ericsson CG, Hamsten A, Nilsson J, Grip L, Svane B, de Faire U Angiographic assessment of effects of bezafibrate on progression of coronary artery disease in young male postinfarction patients. Lancet 1996;347:849-53 [8622389]

de Faire U, Ericsson CG, Hamsten A, Nilsson J Design features of a five-year Bezafibrate Coronary Atherosclerosis Intervention Trial (BECAIT). Drugs Exp Clin Res 1995;21:105-24 [7555614]

BIP, 2000:

, Secondary prevention by raising HDL cholesterol and reducing triglycerides in patients with coronary artery disease: the Bezafibrate Infarction Prevention (BIP) study. Circulation 2000; 102:21-7 [10880410]

Goldenberg I, Boyko V, Tennenbaum A, Tanne D, Behar S, Guetta V Long-term benefit of high-density lipoprotein cholesterol-raising therapy with bezafibrate: 16-year mortality

follow-up of the bezafibrate infarction prevention trial. Arch Intern Med 2009;169:508-14 [19273782]

LEADER, 2002:

Meade T, Zuhrie R, Cook C, Cooper J Bezafibrate in men with lower extremity arterial disease: randomised controlled trial. BMJ 2002;325:1139 [12433762]

SENDCAP, 1998:

Elkeles RS, Diamond JR, Poulter C, Dhanjil S, Nicolaidis AN, Mahmood S, Richmond W, Mather H, Sharp P, Feher MD Cardiovascular outcomes in type 2 diabetes. A double-blind placebo-controlled study of bezafibrate: the St. Mary's, Ealing, Northwick Park Diabetes Cardiovascular Disease Prevention (SENDCAP) Study. Diabetes Care 1998;21:641-8 [9571357]

Acheson, 1972:

Acheson J, Hutchinson EC Controlled trial of clofibrate in cerebral vascular disease. Atherosclerosis 1972;15:177-83 [4579955]

Begg, 1971:

Begg TB, Rifkind BM [Evaluation of clofibrate therapy in peripheral arteriopathy] Minerva Med 1971;62:3469-75 [5097219]

CDP Clofibrate, 1975:

, Clofibrate and niacin in coronary heart disease. JAMA 1975; 231:360-81 [1088963]

Cullen, 1974:

Hanefeld, 1991:

Hanefeld M, Fischer S, Schmechel H, Rothe G, Schulze J, Dude H, Schwanebeck U, Julius U Diabetes Intervention Study. Multi-intervention trial in newly diagnosed NIDDM. Diabetes Care 1991;14:308-17 [2060433]

Harrold, 1969:

Harrold BP, Marmion VJ, Gough KR A double-blind controlled trial of clofibrate in the treatment of diabetic retinopathy. Diabetes 1969;18:285-91 [4894161]

Newcastle, 1971:

, Trial of clofibrate in the treatment of ischaemic heart disease. Five-year study by a group of physicians of the Newcastle upon Tyne region. Br Med J 1971; 4:767-75 [4943605]

Scottish, 1971:

, Ischaemic heart disease: a secondary prevention trial using clofibrate. Report by a research committee of the Scottish Society of Physicians. Br Med J 1971; 4:775-84 [4943606]

VA Neurology Section, 1974:

The treatment of cerebrovascular disease with clofibrate. Final report of the Veterans Administration Cooperative Study of Atherosclerosis, Neurology Section. Stroke 1973;4:684-93 [4723698]

WHO clofibrate, 1978:

, WHO cooperative trial on primary prevention of ischaemic heart disease with clofibrate to lower serum cholesterol: final mortality follow-up. Report of the Committee of Principal Investigators. Lancet 1984; 2:600-4 [6147641]

Heady JA, Morris JN, Oliver MF, WHO clofibrate/cholesterol trial: clarifications. Lancet 1992; 340:1405-6 [1360101]

Emmerich, 2009:

Emmerich KH, Poritis N, Stelmane I, Klindzane M, Erbler H, Goldsteine J, Grtelmeyer R [Efficacy and safety of etofibrate in patients with non-proliferative diabetic retinopathy] Klin Monbl Augenheilkd 2009;226:561-7 [19644802] 10.1055/s-0028-1109516

DAIS, 2001:

Effect of fenofibrate on progression of coronary-artery disease in type 2 diabetes: the Diabetes Atherosclerosis Intervention Study, a randomised study. Lancet 2001;357:905-10 [11289345]

FIELD, 2005:

The need for a large-scale trial of fibrate therapy in diabetes: the rationale and design of the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study. [IS-RCTN64783481]. Cardiovasc Diabetol 2004 Dec 1;3:9 [15571637]

Scott R, Best J, Forder P, Taskinen MR, Simes J, Barter P, Keech A Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study: baseline characteristics and short-term effects of fenofibrate [ISRCTN64783481]. Cardiovasc Diabetol 2005 Aug 22;4:13 [16111499]

Lancet 2005

Keech A, Simes RJ, Barter P, Best J, Scott R, Taskinen MR, Forder P, Pillai A, Davis T, Glasziou P, Drury P, Kesaniemi YA, Sullivan D, Hunt D, Colman P, d'Emden M, Whiting M, Ehnholm C, Laakso M Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): randomised controlled trial. *Lancet* 2005 Nov 26;366:1849-61 [[16310551](#)]

Hiukka A, Westerbacka J, Leinonen ES, Watanabe H, Wiklund O, Hulten LM, Salonen JT, Tuomainen TP, Yki-Jrvinen H, Keech AC, Taskinen MR Long-term effects of fenofibrate on carotid intima-media thickness and augmentation index in subjects with type 2 diabetes mellitus. *J Am Coll Cardiol* 2008 Dec 16;52:2190-7 [[19095138](#)] [10.1016/j.jacc.2008.09.049](#)

Hiukka A, Westerbacka J, Leinonen ES, Watanabe H, Wiklund O, Hulten LM, Salonen JT, Tuomainen TP, Yki-Jrvinen H, Keech AC, Taskinen MR Long-term effects of fenofibrate on carotid intima-media thickness and augmentation index in subjects with type 2 diabetes mellitus. *J Am Coll Cardiol* 2008 Dec 16;52:2190-7 [[19095138](#)]

Rajamani K, Colman PG, Li LP, Best JD, Voysey M, D'Emden MC, Laakso M, Baker JR, Keech AC Effect of fenofibrate on amputation events in people with type 2 diabetes mellitus (FIELD study): a prespecified analysis of a randomised controlled trial. *Lancet* 2009 May 23;373:1780-8 [[19465233](#)] [10.1016/S0140-6736\(09\)60698-X](#)

Helsinki (HHS), 1987:

Manninen V, Elo MO, Frick MH, Haapa K, Heinonen OP, Heinsalmi P, Helo P, Huttunen JK, Kaitaniemi P, Koskinen P, et al, Lipid alterations and decline in the incidence of coronary heart disease in the Helsinki Heart Study. *JAMA* 1988; 260:641-51 [[3164788](#)]

Frick MH, Elo O, Haapa K, Heinonen OP, Heinsalmi P, Helo P, Huttunen JK, Kaitaniemi P, Koskinen P, Manninen V Helsinki Heart Study: primary-prevention trial with gemfibrozil in middle-aged men with dyslipidemia. Safety of treatment, changes in risk factors, and incidence of coronary heart disease. *N Engl J Med* 1987;317:1237-45 [[3313041](#)]

HHS (Frick)(secondary prev subgroup), 1993:

Frick MH, Heinonen OP, Huttunen JK, Koskinen P, Mnttri M, Manninen V Efficacy of gemfibrozil in dyslipidaemic subjects with suspected heart disease. An ancillary study in the Helsinki Heart Study frame population. *Ann Med* 1993;25:41-5 [[8435186](#)]

LOCAT, 1997:

Frick MH, Syvne M, Nieminen MS, Kauma H, Majahalme S, Virtanen V, Kesniemi YA, Pasternack A, Taskinen MR Prevention of the angiographic progression of coronary and vein-graft atherosclerosis by gemfibrozil after coronary bypass surgery in men with low levels of HDL cholesterol. Lipid Coronary Angiography Trial (LOCAT) Study Group. *Circulation* 1997;96:2137-43 [[9337181](#)]

VA-HIT, 1999:

Rubins HB, Robins SJ, Collins D, Fye CL, Anderson JW, Elam MB, Faas FH, Linares E, Schaefer EJ, Schectman G, Wilt TJ, Wittes J, Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol. Veterans Affairs High-Density Lipoprotein Cholesterol Intervention Trial Study Group. *N Engl J Med* 1999; 341:410-8 [[10438259](#)]

Adabag AS, Mithani S, Al Aloul B, Collins D, Bertog S, Bloomfield HE Efficacy of gemfibrozil in the primary prevention of atrial fibrillation in a large randomized controlled trial. *Am Heart J* 2009 May;157:913-8 [[19376321](#)]

ACCORD lipid, 2010:

Effects of Combination Lipid Therapy in Type 2 Diabetes Mellitus. *N Engl J Med* 2010 Mar 14;: [[20228404](#)] [10.1056/NEJMoa1001282](#)

Effects of Medical Therapies on Retinopathy Progression in Type 2 Diabetes. *N Engl J Med* 2010 Jun 29;: [[20587587](#)] [10.1056/NEJMoa1001288](#)

Ismail-Beigi F, Craven T, Banerji MA, Basile J, Calles J, Cohen RM, Cuddihy R, Cushman WC, Genuth S, Grimm RH Jr, Hamilton BP, Hoogwerf B, Karl D, Katz L, Krikorian A, O'Connor P, Pop-Busui R, Schubart U, Simmons D, Taylor H, Thomas A, Weiss D, Hramiak I Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the ACCORD randomised trial. *Lancet* 2010 Jun 29;: [[20594588](#)] [10.1016/S0140-6736\(10\)60576-4](#)

3 niacin

Trial	Treatments	Patients	Trials design and methods
niacin vs control			

continued...

Trial	Treatments	Patients	Trials design and methods
VA drugs , 1968 n=77/143 follow-up: 3.2 years	-	-	Parallel groups double blind
niacin vs placebo			
CDP niacin , 1975 n=1119/2789 follow-up: 6.2 years	niacin 3 mg/d versus placebo	Hombres, de 30 64 ans	Parallel groups double blind
niacin vs ezetimibe			
ARBITER 6-HALTS (niacin vs ezetimibe) , 2009 [NCT00397657] n=97/111 follow-up: 14 months	extended-release niacin 1 g/d, titrated to max tolerable dose up to 2 g/d (HDL-focused strategy) versus ezetimibe 10 mg/d (LDL-focused strategy)	patients with known coronary or vascular disease or coronary risk equivalents	Parallel groups open US

References

VA drugs, 1968:

Schoch HK. The US Veterans Administration Cardiology drug lipid study: an interim report Adv Exp Med Biol. 1968;4:405-420

CDP niacin, 1975:

, Clofibrate and niacin in coronary heart disease. JAMA 1975; 231:360-81 [1088963]

ARBITER 6-HALTS (niacin vs ezetimibe), 2009:

Taylor AJ, Villines TC, Stanek EJ, Devine PJ, Griffen L, Miller M, Weissman NJ, Turco M Extended-release niacin or ezetimibe and carotid intima-media thickness. N Engl J Med 2009 Nov 26;361:2113-22 [19915217]

Villines TC, Stanek EJ, Devine PJ, Turco M, Miller M, Weissman NJ, Griffen L, Taylor AJ The ARBITER 6-HALTS Trial (Arterial Biology for the Investigation of the Treatment Effects of Reducing Cholesterol 6-HDL and LDL Treatment Strategies in Atherosclerosis) Final Results and the Impact of Medication Adherence, Dose, and Treatment Duration. J Am Coll Cardiol 2010 Apr 8;: [20399059] 10.1016/j.jacc.2010.03.017

4 niacin (on top statin)

Trial	Treatments	Patients	Trials design and methods
niacin vs placebo (on top statin)			
AIM-HIGH , 2011 [NCT00120289] n=1718/1691 follow-up: 32 months	high-dose, extended-release niacin in gradually increasing doses up to 2000 mg daily (+ simvastatin) versus placebo	patients with a history of cardiovascular disease, high triglycerides, and low levels of HDL cholesterol	Parallel groups double blind US, Canada
HPS 2-Thrive [NCT00461630] n=12838/12835 follow-up: 3.9y (median)	2 g of extended-release niacin and 40 mg of laropiprant versus placebo	patients with vascular disease	Parallel groups double blind UK, Scandinavia, China

continued...

Trial	Treatments	Patients	Trials design and methods
Oxford Niaspan Study , 2009 [NCT00232531] n=35/36 follow-up: 1 year	niacin 2g daily (added to statin therapy) versus placebo (statins alone)	patients with low HDL-C (<40 mg/dl) and either a type 2 diabetes with coronary heart disease or a carotid/peripheral atherosclerosis	Parallel groups double blind USA
ARBITER 2 , 2009 n=87/80 follow-up: 1 y	long-acting niacin target dose of 1 g/day (added to statin therapy) versus placebo	patients with known coronary artery disease and well controlled on statin therapy	Parallel groups double blind USA
HATS , 2001 n=73/73 follow-up: 3 y	simvastatin plus niacin versus placebo	patients with coronary disease, low HDL cholesterol levels and normal LDL cholesterol levels	Factorial plan double blind USA, Canada

References

AIM-HIGH, 2011:

Boden WE, Probstfield JL, Anderson T, Chaitman BR, Desvignes-Nickens P, Koprowicz K, McBride R, Teo K, Weintraub W Niacin in patients with low HDL cholesterol levels receiving intensive statin therapy. *N Engl J Med* 2011;365:2255-67 [22085343] [10.1056/NEJMoa1107579](https://doi.org/10.1056/NEJMoa1107579)

HPS 2-Thrive, :

Landray MJ, Haynes R, Hopewell JC, Parish S, Aung T, Tomson J, Wallendszus K, Craig M, Jiang L, Collins R, Armitage J Effects of extended-release niacin with laropiprant in high-risk patients. *N Engl J Med* 2014 Jul 17;371:203-12 [25014686] [10.1056/NEJMoa1300955](https://doi.org/10.1056/NEJMoa1300955)

Oxford Niaspan Study, 2009:

Lee JM, Robson MD, Yu LM, Shirodaria CC, Cunningham C, Kylintireas I, Digby JE, Bannister T, Handa A, Wiesmann F, Durrington PN, Channon KM, Neubauer S, Choudhury RP Effects of high-dose modified-release nicotinic Acid on atherosclerosis and vascular function a randomized, placebo-controlled, magnetic resonance imaging study. *J Am Coll Cardiol* 2009;54:1787-94 [19874992]

ARBITER 2, 2009:

Taylor AJ, Sullenberger LE, Lee HJ, Lee JK, Grace KA Arterial Biology for the Investigation of the Treatment Effects of Reducing Cholesterol (ARBITER) 2: a double-blind, placebo-controlled study of extended-release niacin on atherosclerosis progression in secondary prevention patients treated with statins. *Circulation* 2004;110:3512-7 [15537681]

HATS, 2001:

Brown BG, Zhao XQ, Chait A, Fisher LD, Cheung MC, Morse JS, Dowdy AA, Marino EK, Bolson EL, Alaupovic P, Frohlich J, Albers JJ Simvastatin and niacin, antioxidant vitamins, or the combination for the prevention of coronary disease. *N Engl J Med* 2001 Nov 29;345:1583-92 [11757504]

5 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.