

Clinical trials of antidiabetic drugs for diabetes type 2 in all types of patients

TrialResults-center www.trialresultscenter.org

1 insulin therapy

Trial	Treatments	Patients	Trials design and methods
insulin vs control			
UGDP n=414/205 follow-up:	-	-	
UKPDS 33 n=911/896 follow-up:	-	-	

References

UGDP, :

Effects of hypoglycemic agents on vascular complications in patients with adult-onset diabetes. VIII. Evaluation of insulin therapy: final report. Diabetes 1982 Nov;31 Suppl 5:1-81 [6757026]

UKPDS 33, :

Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK Prospective Diabetes Study (UKPDS) Group. Lancet 1998 Sep 12;352:837-53 [9742976]

2 meglitinides (glinides)

Trial	Treatments	Patients	Trials design and methods
repaglinide vs ???			
YSRE0001 [NCT00336310] n=NA follow-up: 12 weeks	Repaglinide versus NA	-	double-blind Taiwan
repaglinide vs control (add on MET)			
Moses , 1999 n=27/27 follow-up: 12 weeks	prestudy dose of metformin with the addition of repaglinide versus prestudy dose of metformin	patients with type 2 diabetes who had inadequate glycemic control (HbA1c >7.1%) when receiving the antidiabetic agent metformin	Parallel groups open
repaglinide vs glibenclamide			

continued...

Trial	Treatments	Patients	Trials design and methods
Landgraf , 1999 n=NA follow-up: 14 weeks	repaglinide, administered preprandially three times daily versus glibenclamide, given preprandially once or twice daily	-	double-blind
Marbury , 1999 n=NA	-	-	
Wolffenbittel , 1999 n=211/109 follow-up: 1 year	repaglinide (0.5-4 mg t.i.d.) versus glyburide (1.75-10.5 mg daily)	-	double-blind
nateglinide vs glibenclamide (add on MET)			
Derosa , 2009 n=124/124 follow-up: 12 months	nateglinide versus glibenclamide	nave type 2 diabetic patients treated with metformin	double-blind
repaglinide vs gliclazide			
AGEE-3783 [NCT01022762] n=NA follow-up:	repaglinide (1 mg repaglinide twice daily (weeks 0-4), titrated versus gliclazide (80 mg gliclazide once daily (weeks 0-4), titrated	Chinese subjects with type 2 diabetes who never have been treated with oral anti-diabetic drugs	China
repaglinide vs glipizide			
Madsbad , 2001 n=256 follow-up: 1 year	repaglinide, 1-4 mg at mealtimes versus glipizide, 5-15 mg daily	-	double-blind
nateglinide vs glyburide (add on MET)			
PRESERVE-beta n=NA follow-up:	-	-	
repaglinide + insulin vs insulin			
AGEE-1524 [NCT00799448] n=NA follow-up:	repaglinide combined with insulin NPH versus biphasic human insulin 30 alone	type 2 diabetics inadequately controlled with sulfonylurea (SU) +/- biguanide therapy	open Greece
AGEE-3020 n=NA	-	-	
mitiglinide+voglibose vs insulin glargine			
GLORIA [NCT00663884] n=NA follow-up: 16 weeks	combination therapy of 10 mg mitiglinide or 0.2mg voglibose versus insulin glargine	diabetic patients whose glycemic control were not enough despite administration of oral antidiabetic drug or insulin glargine	open Korea
nateglinide + metformin vs metformin			

continued...

Trial	Treatments	Patients	Trials design and methods
Horton DOUBLON , 2000 n=172/178 follow-up: 24 weeks	nateglinide (120 mg, ac) and metformin (500 mg, tid) versus 500 mg metformin three times a day	-	
repaglinide vs metformin			
ReMet [NCT00118950] n=NA follow-up:	Repaglinide versus Metformin	Non-Obese Type 2 Diabetic Patients Uncontrolled by Diet	double-blind Denmark
Lund , 2007 n=NA follow-up:	repaglinide 2 mg thrice daily versus metformin 1 g twice daily	non-obese patients with type 2 diabetes	Cross over double-blind
repaglinide + metformin vs metformin			
AGEE-1411 [NCT01465152] n=NA follow-up:	-	-	open Spain
repaglinide vs Metformin (add on insulin)			
Reform [NCT00118963] n=NA follow-up:	Repaglinide + BIAsp30 versus Metformin + BIAsp30	non-obese patients with type-2-diabetes, uncontrolled on diet alone	double-blind
mitiglinide vs nateglinide			
Gao [NCT00461617] n=291 follow-up: 20 weeks	mitiglinide 10 - 20 mg three times daily versus nateglinide 120 mg three times daily	Chinese type 2 diabetes mellitus patients	Parallel groups double-blind
mitiglinide vs on top insulin glargine			
Kumashiro , 2007 n=NA follow-up:	mitiglinide versus on top of once daily insulin glargine	-	
repaglinide vs on top pioglitazone			
Raskin , 2001 n=NA	-	-	
repaglinide vs on top rosiglitazone			
Raskin , 2001 n=NA	-	-	
repaglinide vs on top troglitazone			
Raskin , 2000 n=256 follow-up: 22 weeks	repaglinide (0.54.0 mg at meals), versus combination of repaglinide (14 mg at meals) and troglitazone (200600 mg once daily)	Patients with type 2 diabetes who had inadequate glycemic control (HbA1c 7.0%) during previous monotherapy	open
nateglinide vs placebo			

continued...

Trial	Treatments	Patients	Trials design and methods
CDJN608AUS13 n=NA	-	-	
CDJN608AUS13 n=NA	-	-	
CDJN608A ES03 <i>unpublished</i> n=NA follow-up:	-	-	
NAVIGATOR n=NA follow-up:	-	patients with impaired glucose tolerance (IGT)	
Schwarz , 2008 n=66 follow-up: 12 weeks	nateglinide monotherapy (120 mg, before meals) versus placebo	drug-naive patients with T2DM aged ≥ 65 years	double-blind
Schwarz (study 2) , 2008 n=NA follow-up: 12 weeks	nateglinide monotherapy (120 mg, before meals) versus	drug-naive patients with T2DM aged ≥ 65 years	double-blind
Hanefeld , 1990 n=229/60 follow-up: 12 weeks	nateglinide at doses of 30 mg, 60 mg, 120 mg, or 180 mg versus placebo	-	Parallel groups double-blind
Horton , 2000 n=179/172 follow-up:	120 mg nateglinide before meals versus placebo	patients with an HbA1c level between 6.8 and 11.0% during a 4-week placebo run-in	Parallel groups double-blind
Mari , 2005 n=108 follow-up: 24 weeks	30, 60, or 120 mg nateglinide versus placebo	mild type 2 diabetic men and women (fasting glucose 7.0-8.3 mmol/l) on diet treatment	Parallel groups double-blind
Marre , 2002 n=NA follow-up:	nateglinide 60 mg, 120 mg before three meals versus placebo	metformin-treated patients with HbA1c between 6.8% and 11%	Parallel groups double-blind
Moses , 2001 n=408 follow-up: 16 weeks	0.5 mg repaglinide at mealtimes (increased to 1 mg after 4 weeks depending on blood glucose response) versus placebo	patients with type 2 diabetes considered poorly controlled by diet, but without a history of previous antidiabetic medication	Parallel groups double-blind
Saloranta , 2002 n=675 follow-up: 24 weeks	nateglinide (30, 60, or 120 mg, with meals). versus placebo	patients with type 2 diabetes but only moderately elevated fasting plasma glucose (FPG = 7.0-8.3 mmol/liter)	Parallel groups double-blind
repaglinide vs placebo			
Goldberg , 1998 n=66/33 follow-up: 18 weeks	repaglinide versus placebo	patients with type 2 diabetes	Parallel groups double-blind

continued...

Trial	Treatments	Patients	Trials design and methods
Jovanovic , 2000 n=286/75 follow-up: 24 weeks	repaglinide 1 mg (n = 140), or repaglinide 4 mg (n = 146) versus placebo	-	double-blind
Chuang , 1999 n=NA	-	-	
Bech , 2003 n=253 follow-up: 16 weeks	repaglinide initiated at 0.5 mg per meal, increased to 1 mg after 4 weeks if fasting plasma glucose exceeded 7.8 mmol/l. versus placebo	pharmacotherapy-naive patients with Type 2 diabetes	Parallel groups double-blind
Goldberg , 1998 n=66/33 follow-up:	repaglinide versus placebo	type 2 diabetes	Parallel groups double-blind
Jovanovic , 2000 n=286/75 follow-up: 24 weeks	repaglinide 1 mg or repaglinide 4 mg versus placebo	-	Parallel groups double-blind
nateglinide vs placebo (add on insulin)			
Dashora , 2007 n=55 follow-up: 16 weeks	nateglinide before meals versus placebo	-	double-blind
nateglinide vs placebo (add on insulin+MET)			
Juurinen , 2009 n=NA follow-up: 24 weeks	nateglinide (120 mg three times daily) before main meals versus placebo (add on insulin+MET)	Type 2 diabetes treated with the combination of basal insulin and metformin	Parallel groups double-blind
mitiglinide vs placebo (add on MET)			
NCT01037842 [NCT01037842] n=NA follow-up: 16 weeks	mitiglinide versus placebo	patients with type 2 diabetes who show inadequate glycemic control with metformin monotherapy	double-blind Korea
EX-1510-CT-003 [NCT00519142] n=NA follow-up: 24 weeks	metformin + mitiglinide three times a day with meals versus (metformin + placebo for mitiglinide)	patients with Type 2 diabetes mellitus not well controlled with metformin alone	double-blind US
nateglinide vs placebo (add on standard treatment)			
NCT00402909 [NCT00402909] n=NA follow-up:	-	patients with type 2 diabetes who are not achieving glycemic control with glargine, metformin and/or thiazolidinedione only	double-blind
nateglinide vs placebo (add on TZD)			
026-CL-004 <i>ongoing</i> [NCT00189774] n=NA follow-up:	nateglinide versus placebo (on top pioglitazone)	inadequately controlled type 2 diabetic patients with pioglitazone treatment	double-blind Japan

continued...

Trial	Treatments	Patients	Trials design and methods
repaglinide vs placebo (on top bedtime NPH-insulin)			
Landin-Olsson , 1999 n=NA	-	-	
mitiglinide vs placebo (on top pioglitazone)			
Kaku , 2009 n=NA follow-up: 16 weeks	additional mitiglinide 5 or 10 mg tid versus placebo on top pioglitazone	Japanese type 2 diabetic patients who are insufficiently controlled by pioglitazone monotherapy	Parallel groups multicenter
metformin + repaglinide vs repaglinide			
AGEE-1411 [NCT01465152] n=NA follow-up: 24 weeks	metformin and repaglinide versus repaglinide	subjects with type 2 diabetes in which diet and exercise have failed	open Spain
nateglinide vs repaglinide			
Rosenstock , 2004 n=74/76 follow-up: 16 week	nateglinide monotherapy versus repaglinide monotherapy	type 2 diabetic patients previously treated with diet and exercise	open
Li , 2009 n=NA follow-up:	Nateglinide versus repaglinide	-	
Li , 2007 n=115/115 follow-up: 12 weeks	nateglinide 90 mg three times daily versus repaglinide 1.0 mg three times daily	Chinese patients with type 2 diabetes	Parallel groups
repaglinide + metformin vs repaglinide			
AGEE-3705 [NCT00819741] n=NA follow-up:	repaglinide plus metformin versus repaglinide alone	Chinese subjects with type 2 diabetes having an HbA1c (glycosylated haemoglobin A1c) over 8.5 % and who never have taken oral sugar-lowering drugs before	open China
AGEE-3018 n=NA follow-up:	-	-	
repaglinide + metformin vs rosiglitazone + metformin			
Raskin , 2009 [NCT00399711] n=NA follow-up: 26 weeks	repaglinide and metformin fixed dose combination tablet given as twice daily versus twice daily rosiglitazone and metformin fixed dose combination	subjects with type 2 diabetes currently on monotherapy	open USA
repaglinide + metformin vs SU or MET			
AGEE-3017 [NCT00568984] n=NA follow-up:	combination therapy of repaglinide and metformin versus conventional treatment with a sulphonylurea or metformin in monotherapy	-	China
nateglinide vs gliclazide (add on MET)			

continued...

Trial	Treatments	Patients	Trials design and methods
Ristic , 2006 n=133/129 follow-up: 24 weeks	nateglinide plus metformin versus gliclazide plus metformin	Patients with inadequate glucose control on maximal doses of metformin	Parallel groups double-blind

References

YSRE0001, 0:

Moses, 1999:

Moses R, Slobodniuk R, Boyages S, Colagiuri S, Kidson W, Carter J, Donnelly T, Moffitt P, Hopkins H Effect of repaglinide addition to metformin monotherapy on glycemic control in patients with type 2 diabetes. *Diabetes Care* 1999;22:119-24 [[10333912](#)]

Moses R Repaglinide in combination therapy with metformin in Type 2 diabetes. *Exp Clin Endocrinol Diabetes* 1999;107 Suppl 4:S136-9 [[10522839](#)] [10.1055/s-0029-1212169](#)

Landgraf, 1999:

Landgraf R, Bilo HJ, Mller PG A comparison of repaglinide and glibenclamide in the treatment of type 2 diabetic patients previously treated with sulphonylureas. *Eur J Clin Pharmacol* 1999;55:165-71 [[10379630](#)]

Marbury, 1999:

Marbury T, Huang WC, Strange P, Lebovitz H Repaglinide versus glyburide: a one-year comparison trial. *Diabetes Res Clin Pract* 1999;43:155-66 [[10369424](#)]

Wolffenbittel, 1999:

Wolffenbittel BH, Landgraf R A 1-year multicenter randomized double-blind comparison of repaglinide and glyburide for the treatment of type 2 diabetes. Dutch and German Repaglinide Study Group. *Diabetes Care* 1999;22:463-7 [[10097930](#)]

Derosa , 2009:

Derosa G, D'Angelo A, Fogari E, Salvadeo S, Gravina A, Ferrari I, Cicero AF Nateglinide and glibenclamide metabolic effects in nave type 2 diabetic patients treated with metformin. *J Clin Pharm Ther* 2009;34:13-23 [[19125899](#)] [10.1111/j.1365-2710.2008.00984.x](#)

Derosa G, D'Angelo A, Fogari E, Salvadeo S, Gravina A, Ferrari I, Cicero AF Effects of nateglinide and glibenclamide on prothrombotic factors in nave type 2 diabetic patients treated with metformin: a 1-year, double-blind, randomized clinical trial. *Intern Med* 2007;46:1837-46 [[18025765](#)]

AGEE-3783, 0:

Madsbad, 2001:

Madsbad S, Kilhovd B, Lager I, Mustajoki P, Dejgaard A Comparison between repaglinide and glipizide in Type 2 diabetes mellitus: a 1-year multicentre study. *Diabet Med* 2001;18:395-401 [[11472451](#)]

PRESERVE-beta, :

Gerich J, Raskin P, Jean-Louis L, Purkayastha D, Baron MA PRESERVE-beta: two-year efficacy and safety of initial combination therapy with nateglinide or glyburide plus metformin. *Diabetes Care* 2005;28:2093-9 [[16123472](#)]

AGEE-1524, 0:

AGEE-3020, :

GLORIA, 0:

Horton DOUBLON, 2000:

Horton ES, Foley JE, Shen SG, Baron MA Efficacy and tolerability of initial combination therapy with nateglinide and metformin in treatment-naive patients with type 2 diabetes. *Curr Med Res Opin* 2004;20:883-9 [[15200747](#)] [10.1185/030079903125003881](#)

Horton ES, Clinkingbeard C, Gatlin M, Foley J, Mallows S, Shen S Nateglinide alone and in combination with metformin improves glycemic control by reducing mealtime glucose levels in type 2 diabetes. *Diabetes Care* 2000;23:1660-5 [[11092289](#)]

ReMet, 0:

Lund, 2007:

Lund SS, Tarnow L, Stehouwer CD, Schalkwijk CG, Frandsen M, Smidt UM, Pedersen O, Parving HH, Vaag A Targeting hyperglycaemia with either metformin or repaglinide in non-obese patients with type 2 diabetes: results from a randomized crossover trial. *Diabetes Obes Metab* 2007 May;9:394-407 [[17391168](#)]

AGEE-1411, :

Reform, 0:

Lund SS, Tarnow L, Frandsen M, Nielsen BB, Hansen BV, Pedersen O, Parving HH, Vaag AA Combining insulin with metformin or an insulin secretagogue in non-obese patients with type 2 diabetes: 12 month, randomised, double blind trial. *BMJ* 2009;339:b4324 [[19900993](#)]

Gao, 0:

Gao X, , Multicentre, double-blind, randomized study of mitiglinide compared with nateglinide in type 2 diabetes mellitus patients in China. *J Int Med Res* ;37:812-21. [[19589264](#)]

Kumashiro, 2007:

Kumashiro N, Yoshihara T, Kanazawa Y, Shimizu T, Watada H, Tanaka Y, Fujitani Y, Kawamori R, Hirose T, Long-term effect of combination therapy with mitiglinide and once daily insulin glargine in patients who were successfully switched from intensive insulin therapy in short-term study. *Endocr J* 2007;54:163-6. [[17185877](#)]

Raskin, 2001:

Jovanovic L, Jain R, Greco S, et al. Repaglinide/pioglitazone combination therapy of type 2 diabetes *Diabetes* 2001; 50 Suppl. 2: A439

Raskin, 2001:

Raskin P, McGill J, Hale P, et al. Repaglinide/rosiglitazone combination therapy of type 2 diabetes *Diabetes* 2001; 50 Suppl. 2: A128

Raskin, 2000:

Raskin P, Jovanovic L, Berger S, Schwartz S, Woo V, Ratner R Repaglinide/troglitazone combination therapy: improved glycemic control in type 2 diabetes. *Diabetes Care* 2000;23:979-83 [[10895850](#)]

CDJN608AUS13 , :

CDJN608AUS13 , :

CDJN608A ES03, :

NAVIGATOR, :

Holman RR, Haffner SM, McMurray JJ, Bethel MA, Holzhauer B, Hua TA, Belenkov Y, Boolell M, Buse JB, Buckley BM, Chacra AR, Chiang FT, Charbonnel B, Chow CC, Davies MJ, Deedwania P, Diem P, Einhorn D, Fonseca V, Fulcher GR, Gaciong Z, Gaztambide S, Giles T Effect of nateglinide on the incidence of diabetes and cardiovascular events. *N Engl J Med* 2010 Apr 22;362:1463-76 [[20228402](#)] [10.1056/NEJMoa1001122](#)

Califf RM, Boolell M, Haffner SM, Bethel MA, McMurray J, Duggal A, Holman RR Prevention of diabetes and cardiovascular disease in patients with impaired glucose tolerance: rationale and design of the Nateglinide And Valsartan in Impaired Glucose Tolerance Outcomes Research (NAVIGATOR) Trial. *Am Heart J* 2008;156:623-32 [[18946890](#)]

Schwarz , 2008:

Schwarz SL, Gerich JE, Marcellari A, Jean-Louis L, Purkayastha D, Baron MA Nateglinide, alone or in combination with metformin, is effective and well tolerated in treatment-naïve elderly patients with type 2 diabetes. *Diabetes Obes Metab* 2008;10:652-60 [[17941876](#)] [10.1111/j.1463-1326.2007.00792.x](#)

Schwarz (study 2), 2008:

Schwarz SL, Gerich JE, Marcellari A, Jean-Louis L, Purkayastha D, Baron MA Nateglinide, alone or in combination with metformin, is effective and well tolerated in treatment-naïve elderly patients with type 2 diabetes. *Diabetes Obes Metab* 2008;10:652-60 [[17941876](#)] [10.1111/j.1463-1326.2007.00792.x](#)

Hanefeld, 1990:

Hanefeld M, Bouter KP, Dickinson S, Guitard C Rapid and short-acting mealtime insulin secretion with nateglinide controls both prandial and mean glycemia. *Diabetes Care* 2000;23:202-7 [[10868832](#)]

Horton, 2000:

Horton ES, Clinkingbeard C, Gatlin M, Foley J, Mallows S, Shen S Nateglinide alone and in combination with metformin improves glycemic control by reducing mealtime glucose levels in type 2 diabetes. *Diabetes Care* 2000;23:1660-5 [[11092289](#)]

Mari, 2005:

Mari A, Gastaldelli A, Foley JE, Pratley RE, Ferrannini E Beta-cell function in mild type 2 diabetic patients: effects of 6-month glucose lowering with nateglinide. *Diabetes Care* 2005;28:1132-8 [[15855578](#)]

Marre, 2002:

Marre M, Van Gaal L, Usadel KH, Ball M, Whatmough I, Guitard C Nateglinide improves glycaemic control when added to metformin monotherapy: results of a randomized trial with type 2 diabetes patients. *Diabetes Obes Metab* 2002;4:177-86 [[12047396](#)]

Moses, 2001:

Moses RG, Gomis R, Frandsen KB, Schlienger JL, Dedov I Flexible meal-related dosing with repaglinide facilitates glycemic control in therapy-naive type 2 diabetes. *Diabetes Care* 2001;24:11-5 [[11194214](#)]

Saloranta, 2002:

Saloranta C, Hershon K, Ball M, Dickinson S, Holmes D Efficacy and safety of nateglinide in type 2 diabetic patients with modest fasting hyperglycemia. *J Clin Endocrinol Metab* 2002;87:4171-6 [[12213867](#)]

Goldberg, 1998:

Goldberg RB, Einhorn D, Lucas CP, Rendell MS, Damsbo P, Huang WC, Strange P, Brodows RG A randomized placebo-controlled trial of repaglinide in the treatment of type 2 diabetes. *Diabetes Care* 1998;21:1897-903 [[9802740](#)]

Jovanovic, 2000:

Jovanovic L, Dailey G 3rd, Huang WC, Strange P, Goldstein BJ Repaglinide in type 2 diabetes: a 24-week, fixed-dose efficacy and safety study. *J Clin Pharmacol* 2000;40:49-57 [[10631622](#)]

Chuang, 1999:

Chuang LM, Tai TY, Juang JH, et al. Effect of a prandial glucose regulator (NovoNorm) at two doses (0.5mg and 2.0mg) on glycemic control in type 2 diabetes in Taiwan *JAMA SE Asia* 1999; 51 (1): 22-5

Bech, 2003:

Bech P, Moses R, Gomis R The effect of prandial glucose regulation with repaglinide on treatment satisfaction, wellbeing and health status in patients with pharmacotherapy nave Type 2 diabetes: a placebo-controlled, multicentre study. *Qual Life Res* 2003;12:413-25 [[12797714](#)]

Goldberg, 1998:

Goldberg RB, Einhorn D, Lucas CP, Rendell MS, Damsbo P, Huang WC, Strange P, Brodows RG A randomized placebo-controlled trial of repaglinide in the treatment of type 2 diabetes. *Diabetes Care* 1998;21:1897-903 [[9802740](#)]

Jovanovic, 2000:

Jovanovic L, Dailey G 3rd, Huang WC, Strange P, Goldstein BJ Repaglinide in type 2 diabetes: a 24-week, fixed-dose efficacy and safety study. *J Clin Pharmacol* 2000;40:49-57 [[10631622](#)]

Dashora , 2007:

Dashora UK, Sibal L, Ashwell SG, Home PD Insulin glargine in combination with nateglinide in people with Type 2 diabetes: a randomized placebo-controlled trial. *Diabet Med* 2007;24:344-9 [[17298589](#)] [10.1111/j.1464-5491.2007.02094.x](#)

Juurinen , 2009:

Juurinen L, Tiikkainen M, Saltevo J, Nikkil K, Lanki H, Leppvuori E, Kock T, Teikari-Myyr T, Kauppinen-Mkelin R, Kotronen A, Yki-Jrvinen H Nateglinide combination therapy with basal insulin and metformin in patients with Type 2 diabetes. *Diabet Med* 2009;26:409-15 [[19388972](#)] [10.1111/j.1464-5491.2009.02691.x](#)

NCT01037842, :

EX-1510-CT-003, 0:

NCT00402909, :

026-CL-004, 0:

Landin-Olsson, 1999:

Landin-Olsson M, Brogard JMM, Eriksson J, et al. The efficacy of repaglinide administered in combination with bedtime NPH-insulin in patients with type 2 diabetes. A randomized, semi-blinded, parallel-group, multi-centre trial [abstract] Diabetes 1999 May; 48 Suppl. 1: A117

Kaku, 2009:

Kaku K, Tanaka S, Origasa H, Kikuchi M, Akanuma Y, Addition of mitiglinide to pioglitazone monotherapy improves overall glycemic control in Japanese patients with type 2 diabetes: a randomized double blind trial. Endocr J 2009;56:657-64. [19352048]

AGEE-1411, 0:

Rosenstock , 2004:

Rosenstock J, Hassman DR, Madder RD, Brazinsky SA, Farrell J, Khutoryansky N, Hale PM Repaglinide versus nateglinide monotherapy: a randomized, multicenter study. Diabetes Care 2004;27:1265-70 [15161773]

Li, 2009:

Li C, Xia J, Zhang G, Wang S, Wang L Nateglinide versus repaglinide for type 2 diabetes mellitus in China. Acta Diabetol 2009 Dec;46:325-33 [19183841]

Li, 2007:

Li J, Tian H, Li Q, Wang N, Wu T, Liu Y, Ni Z, Yu H, Liang J, Luo R, Li Y, Huang L Improvement of insulin sensitivity and beta-cell function by nateglinide and repaglinide in type 2 diabetic patients - a randomized controlled double-blind and double-dummy multicentre clinical trial. Diabetes Obes Metab 2007 Jul;9:558-65 [17587398]

AGEE-3705, 0:

AGEE-3018, :

Raskin , 2009:

Raskin P, Lewin A, Reinhardt R, Lyness W Twice-daily dosing of a repaglinide/metformin fixed-dose combination tablet provides glycaemic control comparable to rosiglitazone/metformin tablet. Diabetes Obes Metab 2009 Sep;11:865-73 [19476470] 10.1111/j.1463-1326.2009.01062.x

AGEE-3017, 0:

Ristic, 2006:

Ristic S, Collober-Maugeais C, Pecher E, Cressier F Comparison of nateglinide and gliclazide in combination with metformin, for treatment of patients with Type 2 diabetes mellitus inadequately controlled on maximum doses of metformin alone. Diabet Med 2006;23:757-62 [16842480] 10.1111/j.1464-5491.2006.01914.x

Ristic S, Collober-Maugeais C, Cressier F, Tang P, Pecher E Nateglinide or gliclazide in combination with metformin for treatment of patients with type 2 diabetes mellitus inadequately controlled on maximum doses of metformin alone: 1-year trial results. Diabetes Obes Metab 2007;9:506-11 [17587393] 10.1111/j.1463-1326.2006.00632.x

3 sulfonylureas G2

Trial	Treatments	Patients	Trials design and methods
vs			
Cefalu , 1998 n=NA	-	-	
Hermann , 1994 n=NA	-	-	
Charpentier , 2001 n=NA	-	-	
glipizide vs glyburide			
Rosenstock , 1993 n=139 follow-up: 4 months	glipizide, 2.5 or 5 mg/day versus glyburide, 1.25 or 2.5 mg/day	elderly patients with NIDDM that was controlled for at least 3 months with oral sulfonylurea therapy	Parallel groups open

continued...

Trial	Treatments	Patients	Trials design and methods
Birkeland , 1994 n=NA follow-up:	glipizide versus glyburide	NIDDM patients	
Birkeland , 1994 n=NA follow-up: 15 months	glipizide versus glyburide	NIDDM patients	Parallel groups double-blind
glipizide vs placebo			
Simonson , 1997 n=NA follow-up: 4+8 weeks	once-daily doses of 5, 20, 40, or 60 mg glipizide GITS versus placebo	NIDDM patients	Parallel groups double-blind
Testa , 1998 n=377/192 follow-up: 12 weeks	5 to 20 mg of glipizide gastrointestinal therapeutic system (GITS) versus placebo	patients with type 2 diabetes mellitus	Parallel groups USA
glyburide vs placebo			
Garber , 2002 n=NA follow-up:	glyburide 2.5 mg versus placebo	patients with type 2 diabetes who had failed diet and exercise	Parallel groups double-blind
Vray , 1995 n=NA follow-up:	glibenclamide (2.5 mg X 3/d) versus placebo	type 2 diabetic outpatients, 40-70 years of age, treated by diet alone or oral anti-diabetic drugs	Factorial plan double-blind China

11

References

Cefalu, 1998:

Cefalu WT, Bell-Farrow A, Wang ZQ, McBride D, Dagleish D, Terry JG. Effect of glipizide GITS on insulin sensitivity, glycemic indices, and abdominal fat composition in NIDDM. *Drug Dev. Res.* 1998; 44(1):1-7.

Hermann, 1994:

Hermann LS, Scherstin B, Bitzn PO, Kjellstrm T, Lindgrde F, Melander A Therapeutic comparison of metformin and sulfonylurea, alone and in various combinations. A double-blind controlled study. *Diabetes Care* 1994;17:1100-9 [7821128]

Charpentier, 2001:

Charpentier G, Vaur L, Halimi S, Fleury F, Derobert E, Grimaldi A, Oriol V, Etienne S, Altman JJ Predictors of response to glimepiride in patients with type 2 diabetes mellitus. *Diabetes Metab* 2001;27:563-71 [11694855]

Rosenstock, 1993:

Rosenstock J, Corrao PJ, Goldberg RB, Kilo C Diabetes control in the elderly: a randomized, comparative study of glyburide versus glipizide in non-insulin-dependent diabetes mellitus. *Clin Ther* 1993;15:1031-40 [8111800]

Birkeland, 1994:

Birkeland KI, Furueth K, Melander A, Mowinckel P, Vaaler S Long-term randomized placebo-controlled double-blind therapeutic comparison of glipizide and glyburide. Glycemic control and insulin secretion during 15 months. *Diabetes Care* 1994;17:45-9 [8112188]

Birkeland, 1994:

Birkeland KI, Furueth K, Melander A, Mowinckel P, Vaaler S Long-term randomized placebo-controlled double-blind therapeutic comparison of glipizide and glyburide. Glycemic control and insulin secretion during 15 months. *Diabetes Care* 1994;17:45-9 [8112188]

Simonson, 1997:

Simonson DC, Kourides IA, Feinglos M, Shamoon H, Fischette CT Efficacy, safety, and dose-response characteristics of glipizide gastrointestinal therapeutic system on glycemic control and insulin secretion in NIDDM. Results of two multicenter, randomized, placebo-controlled clinical trials. The Glipizide Gastrointestinal Therapeutic System Study Group. Diabetes Care 1997;20:597-606 [9096986]

Testa, 1998:

Testa MA, Simonson DC Health economic benefits and quality of life during improved glycemic control in patients with type 2 diabetes mellitus: a randomized, controlled, double-blind trial. JAMA 1998;280:1490-6 [9809729]

Garber, 2002:

Garber AJ, Larsen J, Schneider SH, Piper BA, Henry D Simultaneous glyburide/metformin therapy is superior to component monotherapy as an initial pharmacological treatment for type 2 diabetes. Diabetes Obes Metab 2002;4:201-8 [12047399]

Vray, 1995:

Vray M, Attali JR Randomized study of glibenclamide versus traditional Chinese treatment in type 2 diabetic patients. Chinese-French Scientific Committee for the Study of Diabetes. Diabete Metab 1995;21:433-9 [8593925]

4 sulfonylureas G2 add on MET

Trial	Treatments	Patients	Trials design and methods
glibenclamide vs c (add on MET)			
Hermann , 1991 n=NA follow-up: 6 months	metformin + glibenclamide versus metformin	patients with non-insulin-dependent diabetes mellitus	Parallel groups
glyburide vs c (add on MET)			
DeFronzo , 1995 n=NA follow-up: 29 weeks	metformin and glyburide versus metformin	patients with non-insulin-dependent diabetes mellitus	double-blind USA
Erle , 1999 n=NA follow-up:	low-dose glyburide plus metformin versus high-dose glyburide alone	-	Cross over
glibenclamide vs control (add on MET)			
Marre (ass) , 2002 n=NA follow-up: 16 weeks	metformin-glibenclamide 500 mg/2.5 mg or metformin-glibenclamide 500 mg/5 mg, titrated with the intention to achieve fasting plasma glucose (FPG) <or = 7 mmol/l versus metformin 500 mg,	patients with Type 2 diabetes mellitus inadequately controlled by metformin monotherapy	Parallel groups double-blind
Tosi , 2003 n=NA follow-up: 6 months	metformin 400 to 2,400 mg/d + glibenclamide 2.5 to 15 mg/d versus metformin (500 to 3,000 mg/d),	-	Cross over double-blind
glipizide vs control (add on MET)			

continued...

Trial	Treatments	Patients	Trials design and methods
Goldstein n=NA follow-up:	glipizide/metformin 5/500 mg tablets versus metformin 500-mg	patients with type 2 DM that is uncontrolled by at least half the maximum labeled daily dose of a sulfonylurea	Cross over open
glyburide vs control (add on MET)			
Blonde , 2002 n=NA follow-up: 16 weeks	glyburide/metformin 2.5 mg/500 mg (n = 160); or glyburide/metformin 5 mg/500 mg (n = 162) versus metformin 500 mg	patients with inadequate glycaemic control on at least half-maximal dose of sulphonylurea	Parallel groups double-blind
Garber , 2003 n=NA follow-up:	glyburide/metformin versus metformin	patients with type 2 diabetes who had inadequate glycemic control [glycosylated hemoglobin A(1C) (A1C), >7% and <12%) with diet and exercise alone	Parallel groups

References

Hermann, 1991:

Hermann LS, Bitzn PO, Kjellstrm T, Lindgrde F, Scherstn B Comparative efficacy of metformin and glibenclamide in patients with non-insulin-dependent diabetes mellitus. *Diabete Metab* 1991;17:201-8 [[1936477](#)]

DeFronzo, 1995:

DeFronzo RA, Goodman AM Efficacy of metformin in patients with non-insulin-dependent diabetes mellitus. The Multicenter Metformin Study Group. *N Engl J Med* 1995;333:541-9 [[7623902](#)] [10.1056/NEJM199508313330902](#)

Erle, 1999:

Erle G, Lovise S, Stocchiero C, Lora L, Coppini A, Marchetti P, Merante D A comparison of preconstituted, fixed combinations of low-dose glyburide plus metformin versus high-dose glyburide alone in the treatment of type 2 diabetic patients. *Acta Diabetol* 1999;36:61-5 [[10436254](#)]

Marre (ass), 2002:

Marre M, Howlett H, Lehert P, Allavoine T Improved glycaemic control with metformin-glibenclamide combined tablet therapy (Glucovance) in Type 2 diabetic patients inadequately controlled on metformin. *Diabet Med* 2002;19:673-80 [[12147149](#)]

Tosi, 2003:

Tosi F, Muggeo M, Brun E, Spiazzi G, Perobelli L, Zanolin E, Gori M, Coppini A, Moghetti P Combination treatment with metformin and glibenclamide versus single-drug therapies in type 2 diabetes mellitus: a randomized, double-blind, comparative study. *Metabolism* 2003;52:862-7 [[12870162](#)]

Goldstein, :

Goldstein BJ, Pans M, Rubin CJ Multicenter, randomized, double-masked, parallel-group assessment of simultaneous glipizide/metformin as second-line pharmacologic treatment for patients with type 2 diabetes mellitus that is inadequately controlled by a sulfonylurea. *Clin Ther* 2003;25:890-903 [[12852706](#)]

Blonde, 2002:

Blonde L, Rosenstock J, Mooradian AD, Piper BA, Henry D Glyburide/metformin combination product is safe and efficacious in patients with type 2 diabetes failing sulphonylurea therapy. *Diabetes Obes Metab* 2002;4:368-75 [[12406033](#)]

Garber, 2003:

Garber AJ, Donovan DS Jr, Dandona P, Bruce S, Park JS Efficacy of glyburide/metformin tablets compared with initial monotherapy in type 2 diabetes. *J Clin Endocrinol Metab* 2003;88:3598-604 [[12915642](#)]

5 sulfonylureas G3 add on insulin

Trial	Treatments	Patients	Trials design and methods
glimepiride vs placebo (add on insulin)			
Riddle , 1994 <i>unpublished</i> n=72/73 follow-up:	Glimepiride (16 mg/day) plus insulin versus insulin plus placebo	obese patients with type 2 diabetes insufficiently controlled by full dosages of sulphonylureas (glimepiride titrated up to 8mg twice daily and with laboratory-monitored FPG of 10 to 16 mmol/L (180 to 300 mg/dl))	

References

Riddle, 1994:

Riddle M, Schneider J, Glimepiride CG. Glimepiride (HOE490) combined with insulin for NIDDM secondary failures to sulfonylurea monotherapy: results of a multicenter trial [abstract]. 15th Int Diab Fed Congr 1994: 418

6 sulfonylureas G3 add on MET

Trial	Treatments	Patients	Trials design and methods
glimepiride vs placebo (add on MET)			
Charpentier , 2001 n=NA follow-up:	metformin and glimepiride versus metformin	Type 2 diabetic patients aged 35-70 years inadequately controlled by metformin monotherapy 2550 mg daily	double-blind France

References

Charpentier, 2001:

Charpentier G, Fleury F, Kabir M, Vaur L, Halimi S Improved glycaemic control by addition of glimepiride to metformin monotherapy in type 2 diabetic patients. Diabet Med 2001;18:828-34 [[11678974](#)]

Charpentier G, Fleury F, Kabir M, Vaur L, Halimi S, Improved glycaemic control by addition of glimepiride to metformin monotherapy in type 2 diabetic patients. Diabet Med 2001;18:828-34. [[11678974](#)]

7 sulfonylureas G3 monotherapy

Trial	Treatments	Patients	Trials design and methods
glimepiride vs placebo			

continued...

Trial	Treatments	Patients	Trials design and methods
Kaneko , 1993 n=62/31 follow-up:	glimepiride 0.25mg od, 0.5mg od versus placebo	-	
Luis Bautista , 2003 n=NA follow-up: 14 weeks	glimepiride with titration to 2 mg and 4 mg for FPG levels >120 mg/dL versus placebo	Mexican American Patients with type 2 diabetes mellitus	Parallel groups double-blind Mexique
Rosenstock , 1996 n=416 follow-up: 14 weeks	glimepiride 8 mg q.d., 4 mg b.i.d., 16 mg q.d., or 8 mg b.i.d versus placebo	previously treated NIDDM patients	Parallel groups double-blind
Schade , 1998 n=123/126 follow-up:	glimepiride at individually determined optimal dose (1-8 mg of glimepiride) for 10+12 weeks versus placebo	patients with type 2 diabetes mellitus for whom diet therapy is unsuccessful	Parallel groups double-blind
Study 201 (Goldberg) , 1996 n=304 follow-up: 14 weeks	glimepiride, 1, 4, or 8 mg once daily versus placebo	patients with NIDDM	Parallel groups double-blind
Study 202 n=122/125 follow-up:	glimepiride 1-8mg od versus placebo	-	
glimepiride vs glibenclamide			
Draeger , 1996 n=524/520 follow-up:	glimepiride 1 mg daily versus 2.5 mg glibenclamide	type 2 diabetic patients stabilised on glibenclamide	Parallel groups double-blind
Protocol 311 n=427/425 follow-up:	glimepiride 1-8mg od versus glibenclamide 1.75-14 mg/day (od or bid)	-	
glimepiride vs gliclazide			
Charpentier (301F) n=96/107 follow-up:	glimepiride 1-4mg od versus gliclazide 80-320 mg/day (od or bid)	-	
glimepiride vs gliclazide or glibenclamide			
Inukai , 2005 n=172 follow-up: 6 months	glimepiride versus gliclazide or glibenclamide	Japanese type 2 diabetic patients (HbA1C >or = 7.0%), maintained on a conventional SU	Parallel groups open Japan
glimepiride od vs glimepiride bid			
Sonnenberg , 1997 n=50/48 follow-up:	glimepirid e6mg od versus glimepiride 3mg bid	-	Cross over
glimepiride vs glipizide			

continued...

Trial	Treatments	Patients	Trials design and methods
Clark (301) , 1997 n=444/208 follow-up:	glimepiride 1-16 mg/day (od or bid) versus glipizide 2.5-40 mg/day (od or bid)	-	
glimepiride vs glyburide			
Dills , 1996 n=289/288 follow-up:	glimepiride 1-16mg od versus non-micronized glyburide 1.25-20mg od	patients with non-insulin dependent diabetes	Parallel groups double-blind

References

Kaneko, 1993:

Kaneko T, Kaku K, Sakamoto N, et al. Study on minimum effective dose of glimepiride (HOE490) for non-insulin dependent diabetes mellitus patients. Multi center double blind trial compared to placebo [in Japanese] Rinsho Iyaku 1993; 9 (4): 827-48

Luis Bautista, 2003:

Luis Bautista J, Bugos C, Dirnberger G, Atherton T Efficacy and safety profile of glimepiride in Mexican American Patients with type 2 diabetes mellitus: a randomized, placebo-controlled study. Clin Ther 2003;25:194-209 [12637120]

Rosenstock, 1996:

Rosenstock J, Samols E, Muchmore DB, Schneider J Glimepiride, a new once-daily sulfonylurea. A double-blind placebo-controlled study of NIDDM patients. Glimepiride Study Group. Diabetes Care 1996;19:1194-9 [8908379]

Schade, 1998:

Schade DS, Jovanovic L, Schneider J A placebo-controlled, randomized study of glimepiride in patients with type 2 diabetes mellitus for whom diet therapy is unsuccessful. J Clin Pharmacol 1998;38:636-41 [9702849]

Study 201 (Goldberg), 1996:

Goldberg RB, Holvey SM, Schneider J A dose-response study of glimepiride in patients with NIDDM who have previously received sulfonylurea agents. The Glimepiride Protocol 201 Study Group. Diabetes Care 1996;19:849-56 [8842603]

Study 202, :

Hoechst-Roussel Pharmaceuticals Inc. USA. A placebo controlled, dose-titration study of HOE490 in patients with noninsulin- dependent diabetes mellitus (NIDDM) Protocol 202.

Draeger, 1996:

Draeger KE, Wernicke-Panten K, Lomp HJ, Schler E, Roskamp R Long-term treatment of type 2 diabetic patients with the new oral antidiabetic agent glimepiride (Amaryl): a double-blind comparison with glibenclamide. Horm Metab Res 1996;28:419-25 [8911976] 10.1055/s-2007-979830

Draeger KE, Wernicke-Panten K, Lomp HJ, Schler E, Roskamp R, Long-term treatment of type 2 diabetic patients with the new oral antidiabetic agent glimepiride (Amaryl): a double-blind comparison with glibenclamide. Horm Metab Res 1996;28:419-25. [8911976] 10.1055/s-2007-979830

Protocol 311, :

Draeger E, Roskamp R, Lomp H-J, et al. Multicenter clinical trial to study the effects of glimepiride during long-term treatment of type II diabetic patients; double-blind, parallel-group comparison of glimepiride and Euglucon N (micronised glibenclamide). Protocol 311. Hoechst AG.

Charpentier (301F), :

Charpentier G. Comparative double-blind study of the effective dosages of glimepiride and gliclazide in non-insulin-dependent diabetics Protocol 301F. Laboratoires Hoechst, France

Inukai, 2005:

Inukai K, Watanabe M, Nakashima Y, Sawa T, Takata N, Tanaka M, Kashiwabara H, Yokota K, Suzuki M, Kurihara S, Awata T, Katayama S Efficacy of glimepiride in Japanese type 2 diabetic subjects. Diabetes Res Clin Pract 2005;68:250-7 [15936468] 10.1016/j.diabres.2004.10.002

Inukai K, Watanabe M, Nakashima Y, Sawa T, Takata N, Tanaka M, Kashiwabara H, Yokota K, Suzuki M, Kurihara S, Awata T, Katayama S, Efficacy of glimepiride in Japanese type 2 diabetic subjects. *Diabetes Res Clin Pract* 2005;68:250-7. [[15936468](#)] [10.1016/j.diabres.2004.10.002](#)

Sonnenberg, 1997:

Sonnenberg GE, Garg DC, Weidler DJ, Dixon RM, Jaber LA, Bowen AJ, DeChemey GS, Mullican WS, Stonesifer LD Short-term comparison of once- versus twice-daily administration of glimepiride in patients with non-insulin-dependent diabetes mellitus. *Ann Pharmacother* 1997;31:671-6 [[9184703](#)]

Clark (301), 1997:

Clark Jr CM, Goldberg RB. Glimepiride dosing and efficacy: results of placebo-controlled, dose-regimen, and active-controlled trials *Postgrad Med* 1997 Jun Special Report: 45-56

Dills, 1996:

Dills DG, Schneider J Clinical evaluation of glimepiride versus glyburide in NIDDM in a double-blind comparative study. Glimepiride/Glyburide Research Group. *Horm Metab Res* 1996;28:426-9 [[8911977](#)] [10.1055/s-2007-979831](#)

8 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.