

Clinical trials of benazepril + amlodipine

TrialResults-center www.trialresultscenter.org

1 hypertension

Trial	Treatments	Patients	Trials design and methods
benazepril + amlodipine vs benazepril + hydrochlorothiazide			
ACCOMPLISH (diabetic subgroup) , 2010 [NCT00170950] n=1432/1410 follow-up: 36 months	benazepril, combined with amlodipine versus benazepril, combined with hydrochlorothiazide	patients with diabetes (subgroup) and hypertension at high risk of cardiovascular and related events	Parallel groups double-blind US, Norway, Denmark, Finland

More details and results :

- anti hypertensive agents for hypertension in diabetic patients at <http://www.trialresultscenter.org/go-Q10>

References

ACCOMPLISH (diabetic subgroup), 2010:

Weber MA, Bakris GL, Jamerson K, Weir M, Kjeldsen SE, Devereux RB, Velazquez EJ, Dahlf B, Kelly RY, Hua TA, Hester A, Pitt B Cardiovascular events during differing hypertension therapies in patients with diabetes. J Am Coll Cardiol 2010;56:77-85 [20620720] 10.1016/j.jacc.2010.02.046

Jamerson K, Weber MA, Bakris GL, Dahlf B, Pitt B, Shi V, Hester A, Gupte J, Gatlin M, Velazquez EJ Benazepril plus amlodipine or hydrochlorothiazide for hypertension in high-risk patients. N Engl J Med 2008;359:2417-28 [19052124] 10.1056/NEJMoa0806182

2 diabetes type 2

Trial	Treatments	Patients	Trials design and methods
benazepril + amlodipine vs benazepril + hydrochlorothiazide			
ACCOMPLISH (diabetic subgroup) , 2010 [NCT00170950] n=1432/1410 follow-up: 36 months	benazepril, combined with amlodipine versus benazepril, combined with hydrochlorothiazide	patients with diabetes (subgroup) and hypertension at high risk of cardiovascular and related events	Parallel groups double-blind US, Norway, Denmark, Finland

More details and results :

- anti hypertensive agents for diabetes type 2 in patients with hypertension at <http://www.trialresultscenter.org/go-Q83>
- anti hypertensive agents for diabetes type 2 in patients with or without hypertension at <http://www.trialresultscenter.org/go-Q414>

References

ACCOMPLISH (diabetic subgroup), 2010:

Weber MA, Bakris GL, Jamerson K, Weir M, Kjeldsen SE, Devereux RB, Velazquez EJ, Dahlf B, Kelly RY, Hua TA, Hester A, Pitt B Cardiovascular events during differing hypertension therapies in patients with diabetes. J Am Coll Cardiol 2010;56:77-85 [20620720] [10.1016/j.jacc.2010.02.046](https://doi.org/10.1016/j.jacc.2010.02.046)

Jamerson K, Weber MA, Bakris GL, Dahlf B, Pitt B, Shi V, Hester A, Gupte J, Gatlin M, Velazquez EJ Benazepril plus amlodipine or hydrochlorothiazide for hypertension in high-risk patients. N Engl J Med 2008;359:2417-28 [19052124] [10.1056/NEJMoa0806182](https://doi.org/10.1056/NEJMoa0806182)