

# Clinical trials of immediate systematic balloon angioplasty

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 acute myocardial infarction

Trial	Treatments	Patients	Trials design and methods
<b>immediate systematic balloon angioplasty vs no immediate angioplasty</b>			
ECSG , 1988 n=183/184 follow-up: 1 y	angioplasty as soon as possible (after rtPA) versus non-invasive strategy without immediate CA and PTCA	patients with acute myocardial infarction within 5 h after onset of symptoms	parallel group open Europe
Belenkie , 1991 n=50/39 follow-up: 4 months	immediate PTCA versus delayed PTCA (18-38h)	patients with a patent infarct-related artery after thrombolytic therapy suitable for angioplasty	parallel group open Canada
Ellis , 1994 n=78/73 follow-up:	balloon angioplasty supplemented by further thrombolytic therapy as needed versus conservative therapy	patients with first anterior wall infarction treated with any accepted intravenous thrombolytic regimen and angiographically demonstrated to have an occluded infarct vessel within 8 hours of chest pain onset	
Erbel , 1989 n=103/103 follow-up: 3 years	combined intravenous and intracoronary streptokinase with immediate coronary angioplasty versus combined intravenous and intracoronary streptokinase without immediate coronary angioplasty	patients with acute transmural myocardial infarction	Parallel groups
MERLIN (Sutton) , 2004 n=NA follow-up: 30 days	emergency coronary angiography with rescue PCI versus conservative treatment	patients with STEMI and failed fibrinolysis	Parallel groups
SHOCK (Hochman) , 1999 [NCT00000552] n=152/150 follow-up: 30 days (6y)	emergency revascularization versus initial medical stabilization	patients with cardiogenic shock complicating acute MI	Parallel groups open US

continued...

Trial	Treatments	Patients	Trials design and methods
<b>SWISS-SMASH , 1999</b> n=32/23 follow-up: 30 days (1y)	emergency angiography, followed immediately by revascularization when indicated versus initial medical management	Patients with acute myocardial infarction and early shock	Parallel groups open Europe
<b>TAMI 1 pilot , 1987</b> n=99/98 follow-up: in hospital	Angioplasty within 120 min (after rtPA) versus defered CA (7-10 days) and angioplasty if indicated	patients with acute myocardial infarction.	parallel group open USA
<b>TAMI-5 (Calif) , 1991</b> n=287/288 follow-up:	immediate catheterization with angioplasty for failed thrombolysis (90min after rtPA/urokinase) versus deferred predischarge catheterization on days 5-10, no PTCA planned	patient with acute myocardial infarction	Factorial plan
<b>TIMI 2A , 1988</b> n=195/194 follow-up: 21 days	CA within 120 min of the start of the rtPA infusion. PTCA whether the artery is open or closed versus CA within 18-48hrs. PTCA only if artery open (TIMI 2 or 3)	patient thrombolyzed for a AMI	parallel group open USA
<b>Topol , 1987</b> n=15/13 follow-up: in hospital	immediate PTCA versus no PTCA	patients with evolving transmural myocardial infarction	parallel group open USA

More details and results :

- myocardial revascularization for acute myocardial infarction in all type of patients at <http://www.trialresultscenter.org/go-Q129>
- PCI for acute myocardial infarction in all type of patients at <http://www.trialresultscenter.org/go-Q246>
- myocardial revascularization for acute myocardial infarction in patients in cardiogenic shock at <http://www.trialresultscenter.org/go-Q248>

## References

### ECSG, 1988:

Simoons ML, Arnold AE, Betriu A, de Bono DP, Col J, Dougherty FC, von Essen R, Lambertz H, Lubsen J, Meier B, et al, Thrombolysis with tissue plasminogen activator in acute myocardial infarction: no additional benefit from immediate percutaneous coronary angioplasty. Lancet 1988; 1:197-203 [2893037]

Arnold AE, Simoons ML, Van de Werf F, de Bono DP, Lubsen J, Tijssen JG, Serruys PW, Verstraete M Recombinant tissue-type plasminogen activator and immediate angioplasty in acute myocardial infarction. One-year follow-up. The European Cooperative Study Group. Circulation 1992 Jul;86:111-20 [1617763]

**Belenkie, 1991:**

Belenkie I, Knudtson ML, Roth DL, Hansen JL, Traboulsi M, Hall CA, Manyari D, Filipchuck NG, Schnurr LP, Rosenal TW Relation between flow grade after thrombolytic therapy and the effect of angioplasty on left ventricular function: a prospective randomized trial. Am Heart J 1991 Feb;121:407-16 [1990744]

**Ellis, 1994:**

Ellis SG, da Silva ER, Heyndrickx G, Talley JD, Cernigliaro C, Steg G, Spaulding C, Nobuyoshi M, Erbel R, Vassanelli C Randomized comparison of rescue angioplasty with conservative management of patients with early failure of thrombolysis for acute anterior myocardial infarction. Circulation 1994;90:2280-4 [7955184]

**Erbel, 1989:**

Erbel R, Pop T, Diefenbach C, Meyer J Long-term results of thrombolytic therapy with and without percutaneous transluminal coronary angioplasty. J Am Coll Cardiol 1989;14:276-85; discussion 286-8 [2526830]

**MERLIN (Sutton), 2004:**

Sutton AG, Campbell PG, Graham R, Price DJ, Gray JC, Grech ED, Hall JA, Harcombe AA, Wright RA, Smith RH, Murphy JJ, Shyam-Sundar A, Stewart MJ, Davies A, Linker NJ, de Belder MA A randomized trial of rescue angioplasty versus a conservative approach for failed fibrinolysis in ST-segment elevation myocardial infarction: the Middlesbrough Early Revascularization to Limit INfarction (MERLIN) trial. J Am Coll Cardiol 2004;44:287-96 [15261920]

**SHOCK (Hochman), 1999:**

Hochman JS, Sleeper LA, Webb JG, Sanborn TA, White HD, Talley JD, Buller CE, Jacobs AK, Slater JN, Col J, McKinlay SM, LeJemtel TH Early revascularization in acute myocardial infarction complicated by cardiogenic shock. SHOCK Investigators. Should We Emergently Revascularize Occluded Coronaries for Cardiogenic Shock. N Engl J Med 1999;341:625-34 [10460813]

Hochman JS, Sleeper LA, Webb JG, Dzavik V, Buller CE, Aylward P, Col J, White HD Early revascularization and long-term survival in cardiogenic shock complicating acute myocardial infarction. JAMA 2006 Jun 7;295:2511-5 [16757723]

Hochman JS, Sleeper LA, White HD, Dzavik V, Wong SC, Menon V, Webb JG, Steingart R, Picard MH, Menegus MA, Boland J, Sanborn T, Buller CE, Modur S, Forman R, Desvigne-Nickens P, Jacobs AK, Slater JN, LeJemtel TH One-year survival following early revascularization for cardiogenic shock. JAMA 2001 Jan 10;285:190-2 [11176812]

**SWISS-SMASH, 1999:**

Urban P, Stauffer JC, Bleed D, Khatchatrian N, Amann W, Bertel O, van den Brand M, Danchin N, Kaufmann U, Meier B, Machecourt J, Pfisterer M A randomized evaluation of early revascularization to treat shock complicating acute myocardial infarction. The (Swiss) Multicenter Trial of Angioplasty for Shock-(S)MASH. Eur Heart J 1999 Jul;20:1030-8 [10383377]

**TAMI 1 pilot, 1987:**

Topol EJ, Califf RM, George BS, Kereiakes DJ, Abbottsmith CW, Candela RJ, Lee KL, Pitt B, Stack RS, O'Neill WW A randomized trial of immediate versus delayed elective angioplasty after intravenous tissue plasminogen activator in acute myocardial infarction. N Engl J Med 1987 Sep 3;317:581-8 [2956516]

**TAMI-5 (Califf), 1991:**

Califf RM, Topol EJ, Stack RS, Ellis SG, George BS, Kereiakes DJ, Samaha JK, Worley SJ, Anderson JL, Harrelson-Woodlief L Evaluation of combination thrombolytic therapy and timing of cardiac catheterization in acute myocardial infarction. Results of thrombolysis and angioplasty in myocardial infarction-phase 5 randomized trial. TAMI Study Group. Circulation 1991;83:1543-56 [1902405]

**TIMI 2A, 1988:**

Immediate vs delayed catheterization and angioplasty following thrombolytic therapy for acute myocardial infarction. TIMI II A results. The TIMI Research Group.  
JAMA 1988 Nov 18;260:2849-58 [[2972848](#)]

**Topol, 1987:**

Topol EJ, O'Neill WW, Langburd AB, Walton JA Jr, Bourdillon PD, Bates ER, Grines CL, Schork AM, Kline E, Pitt B A randomized, placebo-controlled trial of intravenous recombinant tissue-type plasminogen activator and emergency coronary angioplasty in patients with acute myocardial infarction. Circulation 1987 Feb;75:420-8 [[2948735](#)]

Entry terms: PTCA