

Clinical trials of isradipine

TrialResults-center www.trialresultscenter.org

1 hypertension

Trial	Treatments	Patients	Trials design and methods
isradipine vs hydrochlorothiazide			
MIDAS , 1996 n=442/441 follow-up: 3y	isradipine 2.5-5.0 mg twice daily versus hydrochlorothiazide 12.5-25 mg Twice daily	HBP	

More details and results :

- anti hypertensive agents for hypertension in all type of patient at <http://www.trialresultscenter.org/go-Q13>
- anti hypertensive agents for hypertension in uncomplicated hypertension at <http://www.trialresultscenter.org/go-Q685>

1

References

MIDAS, 1996:

Borhani NO, Mercuri M, Borhani PA, Buckalew VM, Canossa-Terris M, Carr AA, Kappagoda T, Rocco MV, Schnaper HW, Sowers JR, Bond MG Final outcome results of the Multicenter Isradipine Diuretic Atherosclerosis Study (MIDAS). A randomized controlled trial. JAMA 1996;276:785-91 [8769587]

2 heart failure

Trial	Treatments	Patients	Trials design and methods
isradipine vs placebo			
Van den Toren , 1996 n=19 follow-up: 3 months	isradipine up to 5mg 3 times daily versus placebo	patients with congestive heart failure due to ischaemic heart disease, NYHA class II-III, FE<40%	Parallel groups Double blind The NetherlanA

More details and results :

- calcium channel blockers for heart failure in all type of patients at <http://www.trialresultscenter.org/go-Q70>

References

Van den Toren, 1996:

van den Toren EW, van Veldhuisen DJ, van Bruggen A, van den Broek SA, van Gilst WH, Lie KI Acute hemodynamic and long-term clinical effects of isradipine in patients with coronary artery disease and chronic heart failure. A double-blind, placebo-controlled study. Int J Cardiol 1996 Jan;53:37-43 [8776276]